

MARINE TURTLES

SUPPORTING INFORMATION

(Information collated from reports of the Working Party on Ecosystems and Bycatch and other sources as cited)

CONSERVATION AND MANAGEMENT MEASURES

Marine turtles in the Indian Ocean are currently subject to a number of Conservation and Management Measures adopted by the Commission:

- Resolution 15/01 *on the recording of catch and effort data by fishing vessels in the IOTC area of competence* requires numbers of marine turtles to be recorded or all gear types.
- Resolution 15/02 *mandatory statistical reporting requirements for IOTC CPCs* requires data on marine turtles to be reported as specified in Resolution 12/04. Forms for reporting logbook data on discards according to standard IOTC reporting procedures are located at: www.iotc.org/data/requested-statistics-and-submission-forms
- Resolution 12/04 *on the conservation of marine turtles* recognizes the threatened status of the populations of the six marine turtle species found in the Indian Ocean and that some tuna fishing operations carried out in the Indian Ocean can adversely impact marine turtles. This resolution makes mandatory the collection and provision of data on marine turtle interactions and the use of best handling practices to ensure the best chances of survival for any marine turtles returned to the sea after capture.
- Resolution 11/04 *on a Regional Observer Scheme* requires data on marine turtle interactions to be recorded by observers and reported to the IOTC within 150 days. The Regional Observer Scheme (ROS) started on 1st July 2010, and aims to collect scientific observer data on catch and bycatch on, at least, 5% of the fishing operations of vessel over 24 m and vessel under 24 m fishing outside their EEZ. The requirement under Resolution 11/04 in conjunction with the reporting requirements under Resolution 12/04, means that all CPCs should be reporting marine turtle interactions as part of their annual report to the Scientific Committee.

Extracts from Resolutions 11/04, 15/02 and 12/04

RESOLUTION 11/04 ON A REGIONAL OBSERVER SCHEME

Para. 10. Observers shall:

b) Observe and estimate catches as far as possible with a view to identifying catch composition and monitoring discards, by-catches and size frequency;

RESOLUTION 15/02 MANDATORY STATISTICAL REPORTING REQUIREMENTS FOR IOTC CONTRACTING PARTIES AND COOPERATING NON-CONTRACTING PARTIES (CPCS)

Para. 3. Concerning cetaceans, seabirds and marine turtles data should be provided as stated in Resolutions 13/04 *on Conservation of Cetaceans*, Resolution 12/06 *on reduction the incidental bycatch of seabirds in longline fisheries* and Resolution 12/04 *on the conservation of marine turtles* (or any subsequent superseding resolutions).

RESOLUTION 12/04 ON MARINE TURTLES

Para. 3. CPCs shall collect (including through logbooks and observer programs) and provide to the IOTC Secretariat no later than 30 June of the following year in accordance with Resolution 10/02 (or any subsequent revision), all data on their vessels' interactions with marine turtles. ***The data shall include the level of logbook or observer coverage and an estimation of total mortality of marine turtles incidentally caught in their fisheries.***

Para. 7. CPCs with gillnet vessels that fish for species covered by the IOTC Agreement shall:

- a) require that operators of such vessels record all incidents involving marine turtles during fishing operations in their logbooks¹ and report such incidents to the appropriate authorities of the CPC.

Para. 8. CPCs with longline vessels that fish for species covered by the IOTC Agreement shall:

...

- c) require that operators of such vessels record all incidents involving marine turtles during fishing operations in their logbooks¹ and report such incidents to the appropriate authorities of the CPC

Para. 9. CPCs with purse seine vessels that fish for species covered by the IOTC Agreement shall:

...

- c) require that operators of such vessels record all incidents involving marine turtles during fishing operations in their logbooks¹ and report such incidents to the appropriate authorities of the CPC

¹ This information should include where possible, details on species, location of capture, conditions, actions taken on board and location of release.

INDICATORS

Biology and ecology

Six species of marine turtles inhabit the Indian Ocean and likely interact with the fisheries for tuna and tuna-like species. The following section outlines some key aspects of their biology, distribution and historical exploitation.

Flatback turtle

The flatback turtle (*Natator depressus*) gets its name from its relatively flat, smooth shell, unlike other marine turtles which have a high domed shell. The flatback turtle does not have an oceanic dispersal phase for its small post-hatchlings. It has one of the smallest distribution for marine turtles and is restricted to the Australian continental shelf and adjacent countries of Australia, Indonesia and Papua New Guinea. **Table 1** outlines some of the key life history traits of flatback turtles.

TABLE 1. Biology of the flatback turtle (*Natator depressus*).

| Parameter | Description |
|---------------------------|--|
| Range and stock structure | Flatback turtles forage in northern coastal areas, from Western Australia's Ningaloo Coast across Northern Australia through Torres Strait and as south to Hervey Bay and across the Arafura Sea to Indonesia and the northern Coral Sea along southern Papua New Guinea. Nesting by flatback turtles is only known from Australia with five separate genetic stocks recognised. Small post-hatchling flatbacks feed on zooplankton (including <i>Velevella</i> , lobster larvae, and <i>Lepas</i>) at the surface in coastal waters. As large immature and adult turtles they feed on soft bottom habitats where they feed on soft-bodied prey such as sea pens, sea cucumbers, soft corals and jellyfish. |
| Longevity | Oldest individuals recorded to at least 50 years of age |
| Maturity (50%) | unknown |
| Spawning season | Many females nest every 1 to 5 years, one to four times a season (mean = 2.8), laying clutches of between 50 and 60 eggs. Populations nesting in western and eastern Australia breed during the summer while the populations nesting across northern Australia breed predominantly in the winter and spring. |
| Size (length and weight) | The flatback turtle is a medium-sized marine turtle, growing to up to 0.9 m long and weighing up to 70 kg. |

Sources: Mortimer 1984, FAO 1990; Limpus 2007

Green turtle

The green turtle (*Chelonia mydas*) is the largest of all the hard-shelled marine turtles and is one of the most widely distributed and commonest of the marine turtle species in the Indian Ocean. The Indian Ocean hosts some of the largest nesting populations of green turtles in the world, particularly on oceanic islands in the southwest Indian Ocean and on islands in South East Asia. Many of these populations are now recovering after intense exploitation in the last century that greatly reduced the populations; some populations are, however, still declining.

TABLE 2. Biology of the green turtle (*Chelonia mydas*).

| Parameter | Description |
|---------------------------|---|
| Range and stock structure | Globally distributed and generally found in tropical and subtropical waters along continental coasts and islands between 30°N and 30°S. Green turtles primarily use three types of habitat: open beaches (for nesting), convergence zones in the open ocean (oceanic stage juveniles and migratory adults for reproduction), and benthic feeding grounds in coastal areas (neritic stage juveniles and adults). Adults migrate from foraging areas to mainland or island nesting beaches and may travel hundreds or thousands of kilometres (up to 3500 km) each way. After emerging from the nest, hatchlings swim offshore, where they are believed to drift in major oceanic current systems and live for several years, feeding close to the surface on a variety of pelagic plants and animals. Once the juveniles reach a certain age/size range, they leave the pelagic habitat and travel to nearshore foraging grounds. Adult green turtles are unique among marine turtles in that they are herbivorous, feeding on seagrasses and algae beds. |
| Longevity | Unknown, but it is believed that they get up to 60-80 years old |
| Maturity (50%) | Exact age is unknown, it is believed that sexual maturity is reached between 25 and 30+ years |
| Spawning season | Females return to their natal beaches (i.e. the same beaches where they were born) every 2 to 4 years to nest, laying 3 to 5 clutches of about 125 eggs at roughly 14-day intervals during one nesting season. Nesting seasons can change throughout the year (i.e. winter vs summer) according to the nesting site locations in the Indian Ocean (e.g. Summer time south of the Mozambique Channel, winter time north of the Mozambique Channel).. |
| Size (length and weight) | The largest of all the hard-shelled marine turtles, with an average length of 110 cm and average weight of 145kg. |

Sources: Mortimer 1984, FAO 1990, Dalleau et al. 2012, Bourjea, 2015.

Hawksbill turtle

The hawksbill turtle (*Eretmochelys imbricata*) is a small to medium-sized marine turtle generally found in low concentrations, but widely distributed in the Indian Ocean. The keratinous (horn-like) scales covering the carapace and plastron are known as “tortoise shell,” have been regarded as a semi-precious material for centuries and sought after for manufacture of diverse articles on most continents. The global trade in hawksbill shell significantly reduced populations throughout the range. Fortunately, the species responds well to long term protection of nesting females and eggs at the nesting beach, and decades-of such protection at several sites in the Indian Ocean, have resulted in local population recovery

TABLE 3. Biology of the hawksbill turtle (*Eretmochelys imbricata*).

| Parameter | Description |
|---------------------------|---|
| Range and stock structure | Circumtropical, typically occurring from 20° throughout the range to 20°S latitude. Adult hawksbill turtles are capable of migrating long distances between nesting beaches and foraging areas, but on average make shorter migrations than do green and loggerhead turtles. Hawksbill turtles use a variety of habitats depending on the stages of their life cycle, but are most commonly associated with coral reefs. Post-hatchlings (oceanic stage juveniles) are believed to occupy the same pelagic environment as do post-hatchling green turtles. After a few years in the pelagic zone, small juveniles (carapace lengths of ~30 cm) recruit to coastal benthic foraging grounds. This shift in habitat also involves a shift in feeding strategies, from benthic opportunistically, primarily at the surface to feeding from a variety of substrates, primarily on sponges, anthozoans, tunicates, other relatively soft-bodied invertebrates and algae. Their narrow, pointed beaks allow them to prey selectively in crevices. |
| Longevity | Indo-Pacific hawksbills probably live for a total of about 60 to 70 years. |
| Maturity (50%) | Hawksbills can take 25 to 40 years to reach sexual maturity, and typically reproduce for another 10 to 30 years. |
| Spawning season | Female hawksbills return to their natal beaches to nest at intervals of 2–3 years in the Western Indian Ocean; but this can vary throughout the range. A female typically lays 3-5, egg clutches in a season, each with an average of 70-170 eggs depending on geographic location. Some of the largest national nesting populations of hawksbill turtles occur in or around the Indian Ocean especially in the Seychelles, Indonesia and Western Australia. Nesting generally takes place during the warmest months of the year. |
| Size (length and weight) | In the Indian Ocean, adults weigh around 60 kg but can grow to as large as 90 kg for an average size of 90cm. |

Sources: Mortimer 1984, FAO 1990, Mortimer & Donnelly 2008,

Leatherback turtle

The leatherback turtle (*Dermochelys coriacea*) is the largest turtle and the most widely distributed living reptile in the world. The leatherback turtle is the only marine turtle that has no hard shell: there are no large external keratinous scutes

and the underlying bony shell is composed of a mosaic of hundreds of tiny bones. **Table 5** outlines some of the key life history traits of leatherback turtles.

TABLE 4. Biology of the leatherback turtle (*Dermochelys coriacea*).

| Parameter | Description |
|---------------------------|--|
| Range and stock structure | The leatherback turtle regularly migrates enormous distances, e.g. between the Indian and south Atlantic Oceans. They are commonly found in pelagic areas, but they also forage in coastal waters in certain areas. The distribution and developmental habitats of juvenile leatherback turtles are poorly understood. While the leatherback turtle is not as common in the Indian Ocean as other species, important nesting populations are found in and around the Indian Ocean, including in Indonesia, South Africa, South Mozambique, Sri Lanka and India's Andaman and Nicobar Islands. Adults are capable of tolerating water temperatures well below tropical and subtropical conditions, and special physiological adaptations allow them to maintain body temperature above cool water temperatures. They specialise on soft bodied invertebrates found in the water column, particularly jelly fish and other sorts of "jellies." |
| Longevity | unknown |
| Maturity (50%) | Exact age is unknown, it is believed that sexual maturity is reached at around 15 years |
| Spawning season | Females lay clutches of approximately 100 eggs on sandy, tropical beaches. They nest 6–8 times during a nesting season. The nesting season is during the summer time in South Africa and Mozambique. |
| Size (length and weight) | Mature males and females have an average size of 1.7 m for an average weight of 450 kg (record at 918 kg). |

Sources: FAO 1990, Nel 2013.

Loggerhead turtle

The loggerhead turtle (*Caretta caretta*) is globally distributed and the species is known to be heavily impacted by longline fisheries worldwide. The hatchlings and juveniles are pelagic, living in the open ocean and have the ability to undertake long trans-hemispheric migrations from the South to the North Indian Ocean, assuming a development cycle within the entire Indian Ocean. Adults forage in coastal areas, near shallow sea mounts or in the open sea. Key nesting sites in the Indian Ocean are found in Oman, South Africa and West Australia. The loggerhead turtle is known to be the most impacted by longline fishing in the world. **Table 6** outlines some of the key life history traits of loggerhead turtles.

TABLE 5. Biology of the loggerhead turtle (*Caretta caretta*).

| Parameter | Description |
|---------------------------|---|
| Range and stock structure | Circumglobal, occurring throughout the temperate and tropical regions of the Atlantic, Pacific, and Indian Oceans. Studies in the Atlantic and Pacific Oceans show that loggerhead turtles can spend decades living on the high seas, crossing from one side of an ocean basin to another before taking up residence on benthic coastal waters. Adults are capable of migrating long distances between nesting beaches and foraging areas and late stage juveniles have also been shown to undertake extensive migrations. Their enormous heads and powerful jaws enable them to crush large marine molluscs, on which they specialise. |
| Longevity | unknown |
| Maturity (50%) | Exact age is unknown, it is believed that sexual maturity is reached between 12 and 30 years. Age at maturity was estimated at 21.6 years in Tongaland, South Africa, through tagging studies. |
| Spawning season | Many females nest every 2 to 3 year, three to four times a season, laying clutches of approximately 40 to 190 eggs. Loggerhead turtles nest in relatively few countries in the Indian Ocean and the number of nesting females is generally small, except on Masirah Island (Sultanate of Oman) which supports one of only two loggerhead turtles nesting beaches in the world that have greater than 10,000 females nesting per year. |
| Size (length and weight) | Mature males and females may grow to over one meter long and weigh around 110 kg or more. |

Sources: FAO 1990, Lewison et al., 2004, Rees et al. 2010, Dalleau et al. 2013, Hamann et al. 2013

Olive Ridley turtle

The olive Ridley turtle (*Lepidochelys olivacea*) is considered the most abundant marine turtle in the world, with an estimated 800,000 nesting females annually. The olive Ridley turtle has one of the most extraordinary nesting habits in the natural world. Large groups of turtles gather off shore of nesting beaches. Then, all at once, vast numbers of turtles come ashore and nest in what is known as an "arribada". During these arribadas, hundreds to thousands of females come ashore to lay their eggs at the same time. In the northern Indian Ocean, arribadas occur on three different beaches along the coast of Orissa, India. Gahirmatha used to be one of the largest arribada nesting sites in the world. However, arribada nesting events have been less frequent there in recent years and the average size of nesting females has been smaller, indicative of a declining population. Declines in solitary nesting of olive Ridley turtles have been recorded in

Bangladesh, Myanmar, Malaysia, and Pakistan. In particular, the number of nests in Terengganu, Malaysia has declined from thousands of nests to just a few dozen per year. Solitary nesting also occurs extensively throughout this species' range. Despite the enormous numbers of olive Ridley turtles that nest in Orissa, this species is not generally common throughout much of the Indian Ocean. **Table 7** outlines some of the key life history traits of olive Ridley turtles.

TABLE 6. Biology of the olive Ridley turtle (*Lepidochelys olivacea*).

| Parameter | Description |
|---------------------------|---|
| Range and stock structure | The olive Ridley turtle is globally distributed in the tropical regions of the South Atlantic, Pacific, and Indian Oceans. It is mainly a pelagic species, but it has been known to inhabit coastal areas, including bays and estuaries. Olive Ridley turtles often migrate great distances between feeding and breeding grounds. They have an annual migration from pelagic foraging, to coastal breeding and nesting grounds, back to pelagic foraging. They can dive to depths of about 150 m to forage. |
| Longevity | unknown |
| Maturity (50%) | Reach sexual maturity in around 15 years, a young age compared to some other marine turtle species. |
| Spawning season | Many females nest every year, once or twice a season, laying clutches of approximately 100 eggs. Arribadas occur at the beginning of each year in Indian, from January to March. |
| Size (length and weight) | Adults are relatively small, weighing on average around 70 kg at an average length of 90 cm.. As with other species of marine turtles, their size and morphology varies from region to region. |

Sources: Mortimer 1984, FAO 1990

Availability of information on the interactions between marine turtles and fisheries for tuna and tuna-like species in the Indian Ocean

IOTC CPCs are required to collect data on interactions with marine turtles either through logbooks¹ or onboard observers² (Resolution 12/04) to better understand the nature and extent of the interactions between fisheries for tuna and tuna-like species in the Indian Ocean and marine turtles. While ad hoc pieces of information from a number of sources have been collated as far as possible for this document, it is noted that data presented in various documents such as Working Party papers and National Reports are not considered to be formal data submissions to the IOTC. Formal submissions of data in an electronic and standardized format using the available IOTC templates will considerably improve the quality of data obtained and the type of regional analyses that these data can be used for. **Table 8** highlights which CPCs have provided some form of information to the IOTC on interactions with marine turtles, while Appendix II provides a summary of the observer data and discard data on marine turtle interactions that has been officially reported to the IOTC Secretariat.

Data from other sources and in other regions indicate that threats to marine turtles are highest from gillnets and longline gear, and to a lesser extent purse-seine gear, however, the majority of data reported to the IOTC are based on longline interactions (95%), followed by purse seine (3%) and gillnet (2%) fisheries, while the reported interactions for the pole and line fisheries were nil. Many turtles are still not identified to species level even when they are reported, but of those that are identified, the interactions were most frequently observed with Leatherback followed by Loggerhead and Olive Ridley turtles. There were also some unusual findings such as the presence of Kemp's ridley turtle, suggesting there may also be issues with species identification. Where data were available on the survival of marine turtles, 82% were reported to be released alive, while the remainder were discarded dead. **Table 8** highlights which CPCs have provided some form of information to the IOTC on interactions of marine turtles with fishing gear, while Appendix II provides a summary of the observer data and discard data on turtle interactions that have been officially reported to the IOTC Secretariat.

¹ www.iotc.org/data/requested-statistics-and-submission-forms

² www.iotc.org/science/regional-observer-scheme-science

TABLE 7. Contracting Parties and Cooperating non-Contracting Parties reporting marine turtle interactions to the IOTC (2008–2015), as of 24 November 2016

| CPCs | | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Sources/notes |
|---|----|--------------------------|----------------|----------------|-------------------------|-------------------------|---------------------------------|---------------------------------|-------------------------|---|
| Australia | | | | | | | | | | discard form |
| Belize | | | | | | | | | | No vessel operated in 2015 |
| China | | | | | | | | | | Observer data: submitted reports (2010, 2012,2013,2014,2015) |
| Taiwan,China | | | | | | | | | | Observer data: letter to the Secretariat (2012-2013), Report for SC16 (2008-2011) |
| Comoros | | | | | | | | | | |
| European Union* | LL | (ESP) | (ESP) | (ESP) | (ESP) (PRT) (FRA) | (ESP) (PRT) (FRA) | (ESP) (PRT) (UK) (FRA) | (ESP) (PRT) (UK) (FRA) | (PRT) (PRT) (FRA) | Observer data 2015: FRA,PRT |
| | PS | (ESP) (FRA) | (ESP) (FRA) | (ESP) (FRA) | (ESP) (FRA) | (FRA) | (FRA) | (ESP) (FRA) | (ESP) (FRA) | Discard forms 2015: FRA, ESP |
| Eritrea | | | | | | | | | | |
| France (territories)[1] | | See EU purse seine fleet | | | | 4 (OT, France) | as EU PS | | | See EU Purse Seine |
| Guinea | | | | | | | | | | |
| India | | | | | | | | | | |
| Indonesia | | | | | | | | | | Observer data-2014; IOTC-2016-SC19-NR10 (2015) |
| Iran, Islamic Republic of | | | | | | | | | | |
| Japan | LL | | | | | | | | ? | Observer data-2014 |
| | PS | | | | | | | | | |
| Kenya | | | | | | | | | | |
| Korea, Republic of | LL | | | | | | | | | discard form - 2015 |
| | PS | | | | | | | | | |
| Madagascar | | | | | | | | | | Observer data from foreign vessels |
| Malaysia | | | | | | | | | | |
| Maldives, Republic of | | | | | | | | | | discard form; IOTC-2016-SC19-NR17 Rev_1 |

Marine Turtles

Updated: December 2016

| | | | | | | | | | | |
|--|----|--|--|--|--|--|--|--|--|---|
| Mauritius | LL | | | | | | | | | |
| | PS | | | | | | | | | Observer data 2015 |
| Mozambique | | | | | | | | | | Discard form |
| Oman, Sultanate of | | | | | | | | | | |
| Pakistan | | | | | | | | | | |
| Philippines | | | | | | | | | | <i>No Vessel operated in 2015</i> |
| Seychelles | LL | | | | | | | | | |
| | PS | | | | | | | | | Observer data 2014-15 |
| Sierra Leone | | | | | | | | | | |
| Somalia | | | | | | | | | | |
| Sri Lanka | | | | | | | | | | Discard form |
| South Africa | | | | | | | | | | Discard forms (includes foreign fleets)/Observer data |
| Sudan | | | | | | | | | | |
| Tanzania | | | | | | | | | | <i>IOTC-2016-SC19-NR29</i> |
| Thailand | | | | | | | | | | |
| United Kingdom (OT) | | | | | | | | | | Discard forms for the recreational fishery |
| Vanuatu | | | | | | | | | | |
| Yemen | | | | | | | | | | |
| Cooperating Non-Contracting Parties | | | | | | | | | | |
| Djibouti | | | | | | | | | | |
| Senegal | | | | | | | | | | No fishing activity since 2007 |

Green = CPC reported level of marine turtle interactions; Red = CPC did not report level marine turtle interactions, Blue = no reported fleet activity

Purse seine

European Union observers (covering on average 5% of the operations annually from 2003 to 2007) reported 74 marine turtles caught by EU, France and EU, Spain purse seiners over the period 2003–2007³. The most common species reported was olive Ridley, green and hawksbill turtles, and these were mostly caught on log (natural Fish Aggregation Devices – FAD) sets and returned to the sea alive (although there is no systematic information on survivorship after release). Mortality levels of marine turtles due to entanglement in drifting FADs set by the fishery are still unknown and need to be assessed. The EU has indicated that its purse-seine fleet is making progress towards improved FAD designs aimed at reducing the incidence of entanglement of marine turtles, including the use of biodegradable materials. EU, France has indicated that it is already deploying FADs that are likely to reduce the entangled of marine turtles in both the Atlantic and Indian Oceans, while EU, Spain has indicated that it will conduct experiments in the Atlantic Ocean on several FADs designs aimed at reducing the incidence of entanglement of marine turtles, before recommending a final FAD design to replace current FADs. Data collected through observer programs from 1995 to 2011 on purse seine fishing operations suggested that the purse-seine fishery has a low impact on marine turtles with an estimated 240 (SD=157) individuals incidentally captured annually⁴. This study suggested that drifting FADs, considered a critical conservation issue for this fishery, may play a key role in the aggregation of juvenile turtles and could be improved by avoiding entangling devices such as nets. Nevertheless, initial results suggest that this is not the main source of incidental captures of marine turtles in this fishery.

Longline

There is limited information on the interactions of longline fleets in the IOTC with marine turtles and it is not known if this fishing activity represents a serious threat, as is the case in most other regions of the world.

The South African longline fleets have reported that marine turtle bycatch mainly comprises predominantly leatherback turtles, with lesser amounts of loggerhead, hawksbill and green turtles⁵. Estimated average catch rates of marine turtles ranged from 0.005 to 0.3 marine turtles per 1000 hooks and varied by location, season and year. The highest catch rate reported in one trip was 1.7 marine turtles per 1000 hooks in oceanic waters. Over the period 1997 to 2000, the Programme Palangre Réunionnais⁶ examined marine turtle bycatch on 5,885 longline sets in the vicinity of Reunion Island (19-25° S, 48-54° E). The fishery caught 47 leatherback, 30 hawksbill, 16 green and 25 unidentified marine turtles, equating to an average catch rate of less than 0.02 marine turtles per 1000 hooks over the 4 year study period.

The Fishery Survey of India (FSI) carried out a survey covering the whole Indian EEZ using four longline vessels from 2005 to 2009. During this period around 800,000 hooks were deployed in the Arabian Sea, in the Bay of Bengal and in the waters of Andaman and Nicobar. In total 87 marine turtles (79 olive Ridley, 4 green and 2 hawksbill turtles) were caught. Catch rates were: 0.302 marine turtles per 1000 hooks in the Bay of Bengal area, 0.068 marine turtles per 1000 hooks in the Arabian Sea and 0.008 marine turtles per 1000 hooks in the Andaman and Nicobar waters. The highest occurrence of incidental catches in the Bay of Bengal area is probably due to the large abundance of olive Ridley turtles whose main nesting ground in the Indian Ocean is on the east coast of India, in the Orissa region.

Gillnets

Due to the nature of this gear, the incidental catch of marine turtles is thought to be relatively high compared to that of purse-seine and longline gears, however, quantitative data for this gear type are almost non-existent. While the IOTC currently has very little information on interactions between marine turtles and gillnets, the IOSEA database indicates that the coastal mesh net fisheries occur in about 90% of IOSEA Signatory States in the Indian Ocean, and the fishery is considered to have a moderate to relatively high impact on marine turtles in about half of those IOSEA member States. Given the widespread abundance of mesh net fisheries in the Indian Ocean, there is clearly an urgent need for careful, systematic information to be collected and report on this gear type and its impacts on marine turtles.

Other data sources

The IOTC and the Indian Ocean – South-East Asian Marine Turtle Memorandum of Understanding (IOSEA), an agreement under the Convention on Migratory Species, are actively collecting a range of information on fisheries and marine turtle interactions. The IOSEA database covers information from a wider range of fisheries and gears than those held by the IOTC. The IOSEA Online Reporting Facility⁷ compiles information through IOSEA National Reports on

³IOTC-2008-WPEB-08

⁴Bourjea et al. 2014

⁵IOTC-2006-WPBy-15

⁶Poisson F. and Taquet M. (2001) L'espadon: de la recherche à l'exploitation durable. Programme palangre réunionnais, rapport final, 248 p. available in the website www.ifremer.fr/drvreunion

⁷www.ioseaturtles.org/report.php

potential marine turtle fisheries interactions, as well as various mitigation measures put in place by its Signatory States and collaborating organisations. For example, members provide information on fishing effort and perceived impacts of fisheries that may interact with marine turtles, including longlines, purse seines, FADs, and gillnets. While the information is incomplete for some countries and is generally descriptive rather than quantitative, it has begun to provide a general overview of potential fisheries interactions as well as their extent. No information is available for China, Taiwan, China, Japan, Rep. of Korea (among others) which are not yet signatories to IOSEA. Information is also provided on such mitigation measures as appropriate handling techniques, gear modifications, spatial/temporal closures etc. IOSEA is collecting all of the above information with a view to providing a regional assessment of member States' compliance with the FAO Guidelines on reducing fisheries interactions with marine turtles.

It is also useful to compare the impact of large scale fisheries on marine turtles with other fisheries, such as the artisanal fishery, which may have a greater impact on marine turtles than industrial fisheries. A recent study showed for example that the annual turtle catch in the south-western province of Tulear (Madagascar) alone is between 10 000 and 16 000 (Humber et al., 2010). Another recent study estimated that 5 900 turtles were captured annually in Peru only by the national small-scale long-line, bottom set nets fisheries (Alfaro-Shigueto et al., 2011). Lastly, the impacts of fisheries must also be considered in light of other land-based or coastal threats. Nevertheless, it is also clear that, despite strong legislation prohibiting the direct take of turtles throughout, it is still regarded as the most important threat (see review for the IO in Bourjea, 2015).

ASSESSMENT

A number of comprehensive assessments of the status of Indian Ocean marine turtles are available, in addition to the IUCN threat status:

- Hawksbill turtle – Marine Turtle Specialist Group 2008 IUCN Red List status assessment⁸.
- Loggerhead turtle – 2009 status review under the U.S. endangered species act⁹.
- Loggerhead turtle – 2013 Assessment of the conservation status of the loggerhead turtle in the Indian Ocean and South-East Asia. IOSEA Species Assessment: Volume II.
- Leatherback turtle – Assessment of the conservation status of the leatherback turtle in the Indian Ocean and South-East Asia (IOSEA Marine Turtle MoU, 2006)¹⁰.
- Leatherback turtle – 2012 Assessment of the conservation status of the leatherback turtle in the Indian Ocean and South-East Asia – 2012 update. IOSEA Marine Turtle MoU Secretariat report
- Green turtle – Marine Turtle Specialist Group IUCN Red List status assessment expected for 2015 - 2016

LITERATURE CITED

- Abreu-Grobois A, Plotkin P (IUCN SSC Marine Turtle Specialist Group) (2008) *Lepidochelys olivacea*. In: IUCN 2012. IUCN Red List of Threatened Species. Version 2012.2. <www.iucnredlist.org>. Downloaded on 09 November 2012
- Alfaro-Shigueto, J., Mangel, J.C., Bernedo, F., Dutton, P.H., Seminoff, J.A. & Godley, B.J., 2011. Small-scale fisheries of Peru: a major sink for marine turtles in the Pacific. *Journal of Applied Ecology*, doi: 10.1111/j.1365-2664.2011.02040.x
- Bourjea Jerome (2015). Sea turtles; a review of status, distribution and interaction with fisheries in the Southwest Indian Ocean. In Van der Elst RP and Everett BI. 2015. (Eds). *Offshore fisheries of the Southwest Indian Ocean: their status and the impact on vulnerable species*. Oceanographic Research Institute, Special Publication, 10. 448 pp. Chap.9, pp.325-349 (Van der Elst RP and Everett BI)
- Bourjea J, Clermont S, Delgado A, Murua H, Ruiz J, Ciccione S, Chavance P (2014) Marine turtle interactions with purse-seine fishery in the Atlantic and Indian Oceans: Lessons for management. *Biological Conservation* 178: 74-87
- Dalleau M, Ciccione S, Mortimer JA, Garnier J, Benhamou S, Bourjea J (2012) Nesting phenology of Marine Turtles: Insights from a Regional Comparative Analysis on Green Turtle (*Chelonia mydas*). *PLoS ONE* 7(10): e46920. doi:10.1371/journal.pone.0046920
- FAO Species Catalogue (1990) Vol.11: Sea turtles of the world. An annotated and illustrated catalogue of sea turtle species known to date. FAO fisheries synopsis no.125, vol.11. Rome, FAO. 1990. 81p
- Hamann M, Kamrowski RL, Bodine T (2013) Assessment of the conservation status of the loggerhead turtle in the Indian Ocean and South-East Asia. IOSEA Species Assessment: Volume II. 64 p

⁸<http://www.iucnredlist.org/documents/attach/8005.pdf>

⁹<http://www.nmfs.noaa.gov/pr/pdfs/statusreviews/loggerheadturtle2009.pdf>

¹⁰<http://www.ioseaturtles.org/content.php?page=Leatherback%20Assessment>

- Humber, F., Godley, B.J., Ramahery, V. & Broderick, A.C., 2010. Using community members to assess artisanal fisheries: the marine turtle fishery in Madagascar. *Animal conservation* 1 – 11.
- Lewison RL, Freeman SA, Larry B (2004) Quantifying the effects of fisheries on protected species: the impact of pelagic longlines on loggerhead and leatherback sea turtles. *Ecology Letter*, 7(3): 221–231. DOI: 10.1111/j.1461-0248.2004.00573.x
- Limpus CJ (2007) A biological review of Australian marine turtle species. 5. Flatback turtle, *Natator depressus* (Garman). Series: A Biological review of Australian marine turtle species. Vol5. 1-54p
- Marine Turtle Specialist Group (1996) *Caretta caretta*. In: IUCN 2012. IUCN Red List of Threatened Species. Version 2012.2. <www.iucnredlist.org>. Downloaded on 09 November 2012
- Mortimer JA (1984) Marine Turtles in the Republic of the Seychelles: Status and Management. IUCN
- Mortimer JA, Donnelly M (IUCN SSC Marine Turtle Specialist Group) (2008) *Eretmochelys imbricata*. In: IUCN 2012. IUCN Red List of Threatened Species. Version 2012.2. <www.iucnredlist.org>. Downloaded on 09 November 2012
- Mortimer, J.A. & Donnelly, M. 2008. Marine Turtle Specialist Group 2007 IUCN Red List Status Assessment Hawksbill Turtle (*Eretmochelys imbricata*), 121 pages. http://www.iucn-mtsg.org/red_list/ei/index.shtml
- Nel R (2012) Assessment of the conservation status of the leatherback turtle in the Indian Ocean and South-East Asia – 2012 update. IOSEA Marine Turtle MoU Secretariat report, Bangkok, Thailand. 41 p
- Nel R, Punt AE, Hughes GR (2013) Are Coastal Protected Areas Always Effective in Achieving Population Recovery for Nesting Sea Turtles? *PLoS ONE* 8(5): e63525. doi:10.1371/journal.pone.0063525
- Red List Standards & Petitions Subcommittee (1996) *Natator depressus*. In: IUCN 2012. IUCN Red List of Threatened Species. Version 2012.2. <www.iucnredlist.org>. Downloaded on 09 November 2012
- Rees A, Al Saady S, Broderick A, Coyne M, Papatha nasopoulou N, Godley B (2010) Behavioural polymorphism in one of the world's largest populations of loggerhead sea turtles *Caretta caretta*. *Marine Ecology Progress Series* 418:201. doi:10.3354/meps0876
- Sarti Martinez AL (Marine Turtle Specialist Group) (2000) *Dermochelys coriacea*. In: IUCN 2012. IUCN Red List of Threatened Species. Version 2012.2. <www.iucnredlist.org>. Downloaded on 09 November 2012
- Seminoff JA (Southwest Fisheries Science Center, U.S.) (2004) *Chelonia mydas*. In: IUCN 2012. IUCN Red List of Threatened Species. Version 2012.2. <www.iucnredlist.org>. Downloaded on 09 November 2012.

APPENDIX II

OFFICIALLY REPORTED DATA

TABLE 8. Number of turtle interactions by fleet and gear based on observer trip reports submitted to the IOTC Secretariat¹¹ as of 24 November 2016.

| CPC | Gear | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | Coverage level (%) [*] |
|-----------------|------|------|--------|---------|-------|------|-------|---------------------------------|
| Australia | LL | 1 | 0 | 1 | | 1 | | 4.39 |
| China | LL | 1 | | 0 | 0 | 2 | | 0.76 |
| EU(Spain) | PS | | | | | 1 | | 0.54 |
| EU(France)** | LL | 9 | 3 | 10 | 4 | 14 | | 3.03 |
| | PS | | 1+10kg | 2+210kg | 170kg | 40kg | 2 | 15.74 |
| EU(Portugal) | LL | | 3 | 7 | 16 | 0 | 6 | 8.65 |
| Indonesia | LL | | | | | 12 | | 0.02 |
| Japan | LL | 14 | 0 | 2 | | 12 | | 6.05 |
| Korea | LL | 0 | | 0 | 0 | 1 | | 7.24 |
| Korea | PS | | | | | 1 | | 4.35 |
| Sri Lanka | LL | | | | | 0 | 0 | <1.00 |
| | PS | | | | | 0 | | |
| Madagascar | Line | | | 0 | 0 | 0 | | 1.77 |
| Mauritius | PS | | | | | | 3 | 3.67 |
| Mozambique | LL | | | 0 | | | | 0.06 |
| Seychelles | PS | | | | | | 736kg | 9.78 |
| South Africa*** | LL | | 6 | 2 | 5 | 3 | 13 | 3.53 |

^{*}estimated mean annual observer coverage

^{**} A combination of numbers and weights were reported for different trips

^{***}Observer data from South Africa includes both foreign and national flagged vessels

TABLE 9. Number of turtle interactions by fleet and gear based on discard data reported to the IOTC Secretariat¹² as of 24 November 2016.

| CPC | Gear | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 |
|------------------------------|----------|------|------|------|------|------|------|---------|---------|
| Australia | LL | 11 | 30 | 9 | 1 | 1 | | | |
| EU(UK) | LL | | | | | 6 | | | |
| EU(Spain) | PS | | | | | | | | 1 |
| EU(France)** | LL | | | | | 1 | | | 2 |
| | PS | | | | | | 78 | | |
| EU(Portugal) | LL | | | | | 17 | | | |
| Japan | LL | 13 | | | | | | | |
| Korea | LL | | | | | | | 1 | |
| | PS | | | | | | | | 2 |
| Sri Lanka | GILL/RIN | | | | | | 5 | 0 | 70 |
| | LL | | | | | | 17 | 1 | 8 |
| Maldives | LL | | | | | | 93 | 22 | 53 |
| Mozambique | LL | | | | | | | | 3 |
| Taiwan,China | LL | | | | | 14 | 5 | 19 | 1 |
| South Africa (foreign fleet) | LL | 2 | | | | 4 | 6 | 15 (11) | 26 (13) |

^{*}Discard data from South Africa is from South African flagged vessels except for numbers in brackets which are for foreign fleets

¹¹ www.iotc.org/science/regional-observer-scheme-science

¹² www.iotc.org/data/requested-statistics-and-submission-forms

