

DRAFT: EXECUTIVE SUMMARY: PELAGIC THRESHER SHARK

Indian Ocean Tuna Commission
Commission des Thons de l'Océan Indien

**Status of the Indian Ocean pelagic thresher shark (PTH: *Alopias pelagicus*)****TABLE 1.** Pelagic thresher shark: Status pelagic thresher shark (*Alopias pelagicus*) in the Indian Ocean

Area ¹	Indicators		2013 stock status determination
Indian Ocean	Reported catch 2012:	328 t	Uncertain
	Not elsewhere included (nei) sharks:	42,793 t	
Average reported catch 2008–2012:	76 t		
Not elsewhere included (nei) sharks:	48,708 t		
	MSY:	unknown	
	F_{2012}/F_{MSY} :	unknown	
	SB_{2012}/SB_{MSY} :	unknown	
	SB_{2012}/SB_0 :	unknown	

¹Boundaries for the Indian Ocean = IOTC area of competence

Colour key	Stock overfished ($SB_{year}/SB_{MSY} < 1$)	Stock not overfished ($SB_{year}/SB_{MSY} \geq 1$)
Stock subject to overfishing ($F_{year}/F_{MSY} > 1$)		
Stock not subject to overfishing ($F_{year}/F_{MSY} \leq 1$)		
Not assessed/Uncertain		

TABLE 2. Pelagic thresher shark: IUCN threat status of pelagic thresher shark (*Alopias pelagicus*) in the Indian Ocean

Common name	Scientific name	IUCN threat status ¹		
		Global status	WIO	EIO
Pelagic thresher shark	<i>Alopias pelagicus</i>	Vulnerable	–	–

IUCN = International Union for Conservation of Nature; WIO = Western Indian Ocean; EIO = Eastern Indian Ocean

Sources: IUCN 2007, Reardon et al. 2009

NOTE: IOTC Resolution 12/09 *On the conservation of thresher sharks (family Alopiidae) caught in association with fisheries in the IOTC area of competence*, prohibits retention onboard, transshipping, landing, storing, selling or offering for sale any part or whole carcass of thresher sharks of all the species of the family Alopiidae².

INDIAN OCEAN STOCK – MANAGEMENT ADVICE

Stock status. There remains considerable uncertainty in the stock status due to lack of information necessary for assessment or to for the development of other indicators of the stock (Table 1). The current IUCN threat status of ‘Vulnerable’ applies to pelagic thresher shark globally (Table 2). There is a paucity of information available on this species and this situation is not expected to improve in the short to medium term. There is no quantitative stock assessment and limited basic fishery indicators currently available for pelagic thresher shark in the Indian Ocean therefore the stock status is highly uncertain. Pelagic thresher sharks are commonly taken by a range of fisheries in the Indian Ocean. Because of their life history characteristics – they are relatively long lived (+ 20 years), mature at 8-9 years, and have few offspring (2 pups every year), the pelagic thresher shark is vulnerable to overfishing. Therefore stock status remains **uncertain** (Table 1).

Outlook. Current longline fishing effort is directed to other species, however pelagic thresher sharks is a common bycatch these fisheries. Hooking mortality is apparently very high, therefore IOTC regulation 10/12 prohibiting retaining of any part of thresher sharks onboard and promoting life release of thresher shark are apparently ineffective for species conservation. Maintaining or increasing effort will probably result in further declines in biomass,

¹ The process of the threat assessment from IUCN is independent from the IOTC and is presented for information purpose only

² Scientific observers shall be allowed to collect biological samples from thresher sharks that are dead at haulback, provided that the samples are part of the research project approved by the Scientific Committee (or the Working Party on Ecosystems and Bycatch).

productivity and CPUE. However there are few data to estimated CPUE trends, in view of IOTC regulation 10/12 and reluctance of fishing fleet to report information on discards/non-retained catch. The impact of piracy in the western Indian Ocean has resulted in the displacement and subsequent concentration of a substantial portion of longline fishing effort into other areas in the southern and eastern Indian Ocean. It is therefore unlikely that catch and effort on pelagic thresher shark will decline in these areas in the near future, which may result in localised depletion. The following should be noted:

- The available evidence indicates considerable risk to the status of the IO stock at current effort levels.
- Two important sources of data that inform the assessment, total catches and CPUE are highly uncertain and should be investigated further as a priority.
- Noting that current catches (probably largely underestimated) are estimated at an average ~76 t over the last five years ~328 t in 2012, maintaining or increasing effort will probably result in further declines in biomass, productivity and CPUE.
- Mechanisms need to be developed by the Commission to encourage CPCs to comply with their reporting requirement on sharks.

SUPPORTING INFORMATION

(Information collated from reports of the Working Party on Ecosystems and Bycatch and other sources as cited)

CONSERVATION AND MANAGEMENT MEASURES

Pelagic thresher shark in the Indian Ocean are currently subject to a number of Conservation and Management Measures adopted by the Commission:

- Resolution 13/03 *on the recording of catch and effort by fishing vessels in the IOTC area of competence* sets out the minimum logbook requirements for purse seine, longline, gillnet, pole and line, handline and trolling fishing vessels over 24 metres length overall and those under 24 metres if they fish outside the EEZs of their flag States within the IOTC area of competence. As per this Resolution, catch of all sharks must be recorded (retained and discarded).
- Resolution 13/06 *on a scientific and management framework on the conservation of shark species caught in association with IOTC managed fisheries* prohibits, as an interim pilot measure, the retention onboard, transshipment, landing or storing any part or whole carcass of oceanic whitetip sharks (*Carcharhinus longimanus*) (and requests for all other species) by all vessels on the IOTC record of authorised vessels or authorised to fish for tuna or tuna-like species, with the exception of observers who are permitted to collect biological samples (vertebrae, tissues, reproductive tracts, stomachs) from oceanic whitetip sharks that are dead at haulback and artisanal fisheries for the purpose of local consumption, and will conduct a review and an evaluation of the interim measure in 2016.
- Resolution 12/09 *On the conservation of thresher sharks (family Alopiidae) caught in association with fisheries in the IOTC area of competence* prohibits fishing vessels flying the flag of IOTC Members and Cooperating non-Contracting Parties (CPCs) from retaining on board, transshipping, landing, storing, selling or offering for sale any part or whole carcass of thresher sharks of all the species of the family Alopiidae.
- Resolution 11/04 *on a Regional Observer Scheme* requires data on shark interactions to be recorded by observers and reported to the IOTC within 150 days. The Regional Observer Scheme (ROS) started on 1st July 2010.
- Resolution 10/02 *Mandatory statistical requirements for IOTC Members and Cooperating Non-Contracting Parties (CPC's)* indicated that the provisions, applicable to tuna and tuna-like species, are applicable to shark species.
- Resolution 05/05 *Concerning the conservation of sharks caught in association with fisheries managed by IOTC* includes minimum reporting requirements for sharks, calls for full utilisation of sharks and includes a ratio of fin-to-body weight for shark fins retained onboard a vessel.

Extracts from Resolutions 13/03, 13/06, 12/09, 11/04 and 05/05

RESOLUTION 13/03 ON THE RECORDING OF CATCH AND EFFORT BY FISHING VESSELS IN THE IOTC AREA OF COMPETENCE

Para. 1. Each flag CPC shall ensure that all purse seine, longline, gillnet, pole and line, handline and trolling fishing vessels flying its flag and authorized to fish species managed by IOTC be subject to a data recording system.

Para. 8 (start). The flag State and the States which receive this information shall provide all the data for any given year to the IOTC Secretariat by June 30th of the following year on an aggregated basis.

RESOLUTION 13/06 ON A SCIENTIFIC AND MANAGEMENT FRAMEWORK ON THE CONSERVATION OF SHARK SPECIES CAUGHT IN ASSOCIATION WITH IOTC MANAGED FISHERIES

Para. 8. CPCs, especially those targeting sharks, shall submit data for sharks, as required by IOTC data reporting procedures.

RESOLUTION 12/09 ON THE CONSERVATION OF THRESHER SHARKS (FAMILY ALOPIIDAE) CAUGHT IN

ASSOCIATION WITH FISHERIES IN THE IOTC AREA OF COMPETENCE

Para. 2 Fishing Vessels flying the flag of an IOTC Member or Cooperating non-Contracting Party (CPCs) are prohibited from retaining on board, transshipping, landing, storing, selling or offering for sale any part or whole carcass of thresher sharks of all the species of the family Alopiidae, with the exception of paragraph 7.

Para. 3 CPCs shall require vessels flying their flag to promptly release unharmed, to the extent practicable, thresher sharks when brought along side for taking on board the vessel.

Para. 4 CPCs shall encourage their fishers to record and report incidental catches as well as live releases. These data will be then kept at the IOTC Secretariat.

RESOLUTION 11/04 ON A REGIONAL OBSERVER SCHEME

Para. 10. Observers shall:

b) Observe and estimate catches as far as possible with a view to identifying catch composition and monitoring discards, by-catches and size frequency

Resolution 10/02 MANDATORY STATISTICAL REQUIREMENTS FOR IOTC MEMBERS AND COOPERATING NON-CONTRACTING PARTIES (CPC'S)

Para. 3. The provisions, applicable to tuna and tuna-like species, shall also be applicable to the most commonly caught shark species and, where possible, to the less common shark species.

RESOLUTION 05/05 CONCERNING THE CONSERVATION OF SHARKS CAUGHT IN ASSOCIATION WITH FISHERIES MANAGED BY IOTC

Para. 1. CPCs shall annually report data for catches of sharks, in accordance with IOTC data reporting procedures, including available historical data.

Para. 3. CPCs shall take the necessary measures to require that their fishermen fully utilise their entire catches of sharks. Full utilisation is defined as retention by the fishing vessel of all parts of the shark excepting head, guts and skins, to the point of first landing.

FISHERIES INDICATORS***Pelagic thresher shark: General***

Pelagic thresher shark (*Alopias pelagicus*) is a common shark in pelagic coastal and oceanic waters throughout the tropical Indo-Pacific (Fig. 1). This species is commonly confused with common thresher shark (*Alopias vulpinus*), which is mostly temperate species and often recorded under wrong name. Apparently most of tropical records of common thresher sharks in the Indo-Pacific are misidentified pelagic threshers. Due to identification confusions actual distribution and biology of pelagic and common thresher sharks are poorly known. It is probably highly migratory and is epipelagic from the surface to at least 300 m depth (Compagno 2001). It aggregates around seamounts and continental slopes (Compagno 2001). No predation on pelagic thresher sharks has been reported to date; however being smallest species among thresher sharks it may be preyed upon by bigger species such as tiger shark, makos, white sharks, and killer whales. Fishing is a major contributor to adult mortality. This species used its long tail to attack prey (Compagno 2001, Aalbers et al. 2010). Table 3 outlines some of the key life history traits of pelagic thresher shark in the Indian Ocean.




Fig. 1. Pelagic thresher shark: The worldwide distribution of the pelagic thresher shark (source: FAO).

TABLE 3. Pelagic thresher shark: Biology of Indian Ocean pelagic thresher shark (*Alopias pelagicus*).

Parameter	Description
Range and stock structure	In the tropical Indian Ocean, the greatest abundance of pelagic thresher shark occurs at depths of 50 to 300 m, in temperatures ranging from 8 to 25°C. It is considered as highly migratory species however no published information on horizontal movements of pelagic thresher shark is known for the Indian Ocean. Apparently pelagic thresher shark is a solitary fish however it is often aggregated around seamounts or over continental slopes. Area of overlap with IOTC management area = high. No information is available on stock structure.
Longevity	No ageing studies is known for the Indian Ocean, In the Pacific Ocean (China, Taiwan Province) the oldest pelagic thresher

	sharks reported were a 20 year old male (170 cm SL) and a 28 year old female for fish ~ 188 cm SL.
Maturity (50%)	Age: Sexual maturity is attained at 8-9 years (females), 7–8 years (males). Size: Males mature at 140-145 cm standard length (SL) and females at 145–150 cm TL.
Reproduction	Pelagic thresher shark is an ovoviviparous species, without a placental attachment. <ul style="list-style-type: none"> • Fecundity: very low (2) • Generation time: 8–10 years • Gestation period: <12 months • Reproductive cycle: unknown Its potential annual rate of population increase under sustainable fishing is thought to be very low and has been estimated at or 0.033
Size (length and weight)	Maximum size is around 365 cm TL. New-born pups are around 158–190 cm TL. Length–weight relationship for both sexes combined in the Indian Ocean is $TW=0.001*10^{-4}*FL^{2.15243}$

Sources: Lui et al. 1998, Compagno 2001, Reardon et al. 2004, Dulvy et al. 2008

Pelagic thresher shark: Fisheries

Pelagic thresher shark are often targeted by some recreational, semi-industrial and artisanal fisheries and are a bycatch of industrial fisheries (pelagic longline tuna and swordfish fisheries) (Table 4). Typically, the fisheries take pelagic thresher sharks between 120–190 cm FL or 20 to 90 kg (Romanov pers comm). In Australia thresher sharks used to be a target of sport fishermen. Sport fisheries for oceanic sharks are apparently not so common in other Indian Ocean countries.

There is little information on the fisheries prior to the early 1970's. Some countries still fail to collect shark data while others do collect it but fail to report to IOTC. It appears that significant catches of sharks have gone unrecorded in several countries. Furthermore, many existing catch records probably under-represent the actual catches of sharks because they do not account for discards (i.e. do not record catches of sharks for which only the fins are kept or of sharks usually discarded because of their size or condition) or they reflect dressed weights instead of live weights. FAO also compiles landings data on elasmobranchs, but their statistics are limited by the lack of species-specific data and data from the major fleets. Thresher sharks were marketed both locally and in European markets until at least until early 2011 despite IOTC Resolution 12/09. The practice of shark finning is considered to be regularly occurring and on the increase for this species (Clarke et al. 2006, Clarke 2008). The bycatch/release mortality rate is unknown but probably high. In longline fisheries pelagic thresher sharks are often hooked by the tail (Compagno 2001) and die soon afterward. Therefore they are discarded dead if not retained. In most cases discarded sharks are not recorded in fisheries logbooks. Therefore the current IOTC measures (notably Resolution 12/09) appear to have limited conservation effect while contributing to further loss of fisheries data. Other types of conservation efforts such as protected areas should be considered for this species group by the WPEB, taking into account a detailed analysis of catch distribution and 'hotspots' of abundance derived from research data. Extremely common misidentification of this species with common thresher shark aggravate situation with data collection.

TABLE 4. Pelagic thresher shark: Estimated frequency of occurrence and bycatch mortality in the Indian Ocean pelagic fisheries.

Gears	PS	LL		BB/TROL/HAND	GILL	UNCL
		SWO	TUNA			
Frequency	absent	Common		rare	unknown	unknown
Fishing Mortality	no	high	high	unknown	unknown	unknown
Post release mortality	N/A	unknown	unknown	unknown	unknown	unknown

Sources: Boggs 1992, Romanov 2002, 2008

Pelagic thresher shark: Catch trends

The catch estimates for pelagic thresher shark (Table 5) are highly uncertain as is their utility in terms of minimum catch estimates. Five our CPCs have reported detailed data on sharks (i.e. Australia, EU (Spain, Portugal and United Kingdom), I.R. Iran, South Africa, and Sri Lanka) while thirteen CPCs have reported partial data or data aggregated for all species (i.e. Belize, China, Indonesia, Japan, Rep. of Korea, Malaysia, Mozambique, Oman, Philippines, Seychelles, Mauritius, UK-territories, Vanuatu).

TABLE 5. Pelagic thresher shark: Catch estimates for pelagic thresher shark in the Indian Ocean for 2010 to 2012.

Catch		2010	2011	2012
Most recent catch (reported)	pelagic thresher	20 t	17 t	328 t
	nei-sharks	51,581 t	53,658 t	42,793 t

Mean catch (reported) over the last 5 years (2008–2012)	pelagic thresher nei-sharks	76 t 48,708 t
--	--------------------------------	------------------

Note that reported shark catches are incomplete. The catches of sharks are usually not reported and when they are they might not represent the total catches of this species but simply those retained on board. It is also likely that the amounts recorded refer to weights of processed specimens, not to live weights. In 2012, one country reported catches of pelagic thresher sharks in the IOTC region.

A recent project estimated possible thresher shark catches for fleets/countries based on the ratio of shark catch over target species by metier (Murua et al 2013). The estimation was done using target species nominal catch from the IOTC database and assuming that target catches have been accurately declared. The estimated catch from this study highlighted that the possible underestimation of thresher shark in the IOTC database is considerable (i.e. the estimated catch is around 70 times higher than the declared/report and contained in the IOTC database).

Pelagic thresher shark: Nominal and standardised CPUE Trends

Data not available at the IOTC Secretariat. There are no surveys specifically designed to assess shark catch rates in the Indian Ocean. Historical research data shows overall decline both in CPUE and mean weight of thresher sharks (Romanov pers com).

Pelagic thresher shark: Average weight in the catch by fisheries

Data not available.

Pelagic thresher shark: Number of squares fished

Catch and effort data not available.

STOCK ASSESSMENT

No quantitative stock assessment for pelagic thresher shark has been undertaken by the IOTC Working Party on Ecosystems and Bycatch.

LITERATURE CITED

- Aalbers SA, Bernal D, Sepulveda CA (2010) The functional role of the caudal fin in the feeding ecology of the common thresher shark *Alopias vulpinus*. *J Fish Biol* 76, 1863-1868
- Boggs CH (1992) Depth, capture time and hooked longevity of longline-caught pelagic fish: timing bites of fish with chips. *Fish Bull* 90:642-658
- Clarke S (2008) Use of shark fin trade data to estimate historic total shark removals in the Atlantic Ocean. *Aquat Living Res* 21:373-381
- Clarke SC, McAllister MK, Milner-Gulland EJ, Kirkwood GP, Michielsens CGJ, Agnew DJ, Pikitch EK, Nakano H, Shivji MS (2006) Global estimates of shark catches using trade records from commercial markets. *Ecol Lett* 9:1115-1126
- Compagno LJV (2001) Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Volume 2. Bullhead, mackerel and carpet sharks (Heterodontiformes, Lamniformes and Orectolobiformes). *FAO Species Catalogue for Fishery Purposes*. No 1. Vol. 2. FAO, Rome (Italy). 269 p
- Dulvy NK, Baum JK, Clarke S, Compagno LJV, Cortes E, Domingo A, Fordham S, Fowler S, Francis MP, Gibson C, Martinez J, Musick JA, Soldo A, Stevens JD, Valentih S, 2008. You can swim but you can't hide: the global status and conservation of oceanic pelagic sharks and rays. *Aquatic Conservation: Mar FreshwEcosys* 18, 459-482
- IUCN (2007) IUCN Species Survival Commission's Shark Specialist Group. Review of Chondrichthyan Fishes
- Liu K-M, Chen C-T, Liao T-H, Joung S-J (1999) Age, growth, and reproduction of the pelagic thresher shark, *Alopias pelagicus* in the Northwestern Pacific. *Copeia* 1999, 68-74
- Murua H, Santos MN, Chavance P, Amande J, Seret B, Poisson F, Ariz J, Abascal FJ, Bach P, Coelho R, Korta M (2013) EU project for the provision of scientific advice for the purpose of the implementation of the EUPOA sharks: a brief overview of the results for Indian Ocean. IOTC–2013–WPEB09–19
- Reardon M, Márquez F, Trejo T, Clarke SC (2009) *Alopias pelagicus*. In: IUCN 2013. IUCN Red List of Threatened Species. Version 2013.1. <www.iucnredlist.org>. Downloaded on 15 September 2013.
- Romanov EV (2002) Bycatch in the tuna purse-seine fisheries of the western Indian Ocean. *Fish Bull* 100:90-105
- Romanov EV (2008) Bycatch and discards in the Soviet purse seine tuna fisheries on FAD-associated schools in the north equatorial area of the Western Indian Ocean. *Western Indian Ocean J Mar Sci* 7:163-174

Romanov E, Bach P, Romanova N (2008) Preliminary estimates of bycatches in the western equatorial Indian Ocean in the traditional multifilament longline gears (1961-1989) IOTC Working Party on Ecosystems and Bycatch (WPEB) Bangkok, Thailand. 20-22 October, 2008. 18 p