


DRAFT: EXECUTIVE SUMMARY: NARROW-BARRED SPANISH MACKEREL

Indian Ocean Tuna Commission
Commission des Thons de l'Océan Indien



Status of the Indian Ocean narrow-barred Spanish mackerel (COM: *Scomberomorus commerson*) resource

TABLE 1. Narrow-barred Spanish mackerel: Status of narrow-barred Spanish mackerel (*Scomberomorus commerson*) in the Indian Ocean

Area ¹	Indicators		2014 stock status determination
Indian Ocean	Catch ² 2013:	148,695 t	
	Average catch ² 2009–2013:	144,462 t	
	MSY:	137 Kt [93–164 Kt]	
	F _{MSY} :	0.47 (0.41–1.95)	
	B _{MSY} :	229 Kt (132–265K t)	
	F ₂₀₁₂ /F _{MSY} :	0.92 (0.41–1.95)	
B ₂₀₁₂ /B _{MSY} :	1.17 (0.5–1.51)		
B ₂₀₁₂ /B ₀ :	0.59 (0.25–0.75)		

¹Boundaries for the Indian Ocean stock assessment are defined as the IOTC area of competence.

²Nominal catches represent those estimated by the IOTC Secretariat. If these data are not reported by CPCs, the IOTC Secretariat estimates total catch from a range of sources including: partial catch and effort data; data in the FAO FishStat database; catches estimated by the IOTC from data collected through port sampling; data published through web pages or other means; data reported by other parties on the activity of vessels; and data collected through sampling at the landing place or at sea by scientific observers.

Colour key	Stock overfished (SB _{year} /SB _{MSY} < 1)	Stock not overfished (SB _{year} /SB _{MSY} ≥ 1)
Stock subject to overfishing (F _{year} /F _{MSY} > 1)		
Stock not subject to overfishing (F _{year} /F _{MSY} ≤ 1)		
Not assessed/Uncertain		

INDIAN OCEAN STOCK – MANAGEMENT ADVICE

Stock status. Stock Reduction Analysis techniques indicate that the stock is being exploited at a rate that is near F_{MSY} in recent years, and the stock appears to be fully exploited. NWIO (Gulf of Oman Sea countries) indicate that localised depletion may be occurring from an analysis done in 2013, and overfishing is occurring in this area, though the degree of connectivity with other stocks remains unknown. Stock structure issues remain to be clarified with this stock. Based on the weight-of-evidence available, including the two different SRA approaches pursued in 2014, the stock appears to be **not overfished** and **not subject to overfishing** (Table 1, Fig. 2).

Outlook. There remains considerable uncertainty about stock structure and the total catches. The continued increase of annual catches for narrow-barred Spanish mackerel in recent years has further increased the pressure on the Indian Ocean stock as a whole, and the stock is probably near full/optimal utilisation. The apparent fidelity of narrow-barred Spanish mackerel to particular areas/regions is a matter for concern as overfishing in these areas can lead to localised depletion. Research emphasis on improving indicators and exploration of stock structure and stock assessment approaches for data poor fisheries are warranted. There is a high to very high risk of exceeding MSY-based reference points by 2015 if catches are maintained at current (2012) levels (72% risk that SB₂₀₁₅ < SB_{MSY}, and 90% risk that F₂₀₁₅ > F_{MSY}) or a very high risk is catches are increase further (120% of 2012 levels) (90% risk that SB₂₀₁₅ < SB_{MSY}, and 99% risk that F₂₀₁₅ > F_{MSY}) (Table 2).

The following should be noted:

- Maximum Sustainable Yield estimate for the whole Indian Ocean is 137,000 (range 93,000 t–64,000 t).
- Reconstruction of the catch history needs to occur, as do annual catches submitted to the Secretariat.
- Improvement in data collection and reporting is required to assess the stock using more traditional stock assessment techniques.
- Given the rapid increase in narrow-barred Spanish mackerel catch in recent years, some measures need to be taken to slow or reduce catches in the Indian Ocean (Table 2).

- Limit reference points: The Commission has not adopted limit reference points for any of the neritic tunas under its mandate.




Fig. 1. Narrow-barred Spanish mackerel: PFCRA Aggregated Indian Ocean assessment Kobe plot. The Kobe plot presents the trajectories for the range of plausible model options included in the formulation of the final management advice. The trajectory of the geometric mean of the plausible model options is also presented.

TABLE 2. Narrow-barred Spanish mackerel: 2014 PFCRA Aggregated Indian Ocean assessment Kobe II Strategy Matrix. Probability (percentage) of plausible models violating the MSY-based reference points for five constant catch projections (2012 catch level, -10%, -20%, -30% and +20%) projected for 3 and 10 years. Note: from the 2014 stock assessment using catch estimates at that time.

Reference point and projection timeframe	Alternative catch projections (relative to 2012) and weighted probability (%) scenarios that violate reference point				
	70% (100,333 t)	80% (114,666 t)	90% (129,000 t)	100% (143,333 t)	120% (172,000 t)
$B_{2015} < B_{MSY}$	6%	23%	46%	72%	90%
$F_{2015} > F_{MSY}$	0%	10%	54%	90%	99%
$B_{2022} < B_{MSY}$	9%	24%	52%	76%	90%
$F_{2022} > F_{MSY}$	4%	19%	53%	82%	96%

Note: As detailed in Recommendation 14/07, the colour coding used above refers to 25% probability levels associated with the default target reference points of the Commission.

APPENDIX I

SUPPORTING INFORMATION

(Information collated from reports of the Working Party on Neritic Tunas and other sources as cited)

CONSERVATION AND MANAGEMENT MEASURES

Narrow-barred Spanish mackerel (*Scomberomorus commerson*) in the Indian Ocean is currently subject to a number of Conservation and Management Measures adopted by the Commission:

- Resolution 13/03 on the recording of catch and effort by fishing vessels in the IOTC area of competence
- Resolution 14/05 concerning a record of licensed foreign vessels fishing for IOTC species in the IOTC area of competence and access agreement information
- Resolution 12/11 *on the implementation of a limitation of fishing capacity of Contracting Parties and Cooperating Non-Contracting Parties*
- Resolution 10/02 *mandatory statistical requirements for IOTC Members and Cooperating non-Contracting Parties (CPC's)*
- Resolution 10/08 *concerning a record of active vessels fishing for tunas and swordfish in the IOTC area*

FISHERIES INDICATORS

Narrow-barred Spanish mackerel: General

The narrow-barred Spanish mackerel (*Scomberomorus commerson*) is a pelagic, top level predator found throughout tropical marine waters of the Indo-West Pacific. Table 3 outlines some key life history parameters relevant for management.

TABLE 3. Narrow-barred Spanish mackerel. Biology of Indian Ocean narrow-barred Spanish mackerel (*Scomberomorus commerson*)

Parameter	Description
Range and stock structure	A pelagic, top level predator found throughout tropical marine waters of the Indo-West Pacific. Juveniles inhabit shallow inshore areas whereas adults are found in coastal waters out to the continental shelf. Adults are usually found in small schools but often aggregate at particular locations on reefs and shoals to feed and spawn. They appear to undertake lengthy migrations, however, larger individuals may be resident which contributes to a metapopulation structure. Feed primarily on small fishes such as anchovies, clupeids, carangids, also squids and shrimps. Genetic studies carried out on <i>S. commerson</i> from Djibouti, Oman and U.A.E. showed there were small genetic differences among stocks in these three places.
Longevity	~16 years
Maturity (50%)	Age: 1.9 yrs for males and 2.1 yrs for females Size: 72.8 cm for males and 86.3 cm for females.
Spawning season	Females are multiple spawners. Year-round spawning has been observed in east African waters, with peaks during late spring to summer (April-July) and autumn (September-November) coinciding with the two seasonal monsoons which generate high abundances of plankton and small pelagic fish. Spawning in the southern Arabian Gulf occurs in the spring and summer months between April and August.
Size (length and weight)	Maximum: Females and males 240 cm FL; weight 70 kgs.

n.a. = not available. Sources: Grandcourt et al. 2005, Froese & Pauly 2009, Darvishi et al. 2011

Narrow-barred Spanish mackerel – Fisheries and catch trends

Narrow-barred Spanish mackerel¹ is targeted throughout the Indian Ocean by artisanal and recreational fishers. The main method of capture is gillnet, but significant numbers of are also caught trolling (Table 4, Fig. 2).


¹ Hereinafter referred to as Spanish mackerel

TABLE 4. Narrow-barred Spanish mackerel: Best scientific estimates of the catches of narrow-barred Spanish mackerel by type of fishery for the period 1950–2013 (in metric tonnes). (Data as of October 2014)

Fishery	By decade (average)						By year (last ten years)									
	1950s	1960s	1970s	1980s	1990s	2000s	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Purse seine	0	0	285	2,357	4,145	5,611	4,566	5,880	7,632	6,586	6,130	8,459	8,850	8,812	8,758	8,487
Gillnet	8,681	16,863	29,733	51,763	60,006	64,952	61,998	57,819	66,205	71,501	71,263	72,426	74,925	80,050	88,621	82,368
Line	2,581	3,300	7,106	14,467	14,741	19,453	17,398	19,191	19,846	21,293	23,065	25,847	25,550	27,435	31,769	31,941
Other	57	96	468	5,614	9,747	21,345	19,564	20,515	23,905	25,516	22,735	28,170	25,519	27,455	30,970	25,899
Total	11,318	20,260	37,593	74,201	88,639	111,360	103,526	103,406	117,588	124,895	123,192	134,902	134,844	143,753	160,118	148,695

The catch estimates for narrow-barred Spanish mackerel were derived from very small amounts of information and are therefore highly uncertain². The catches provided in Table 4 are based on the information available at the IOTC Secretariat and the following observations on the catches cannot currently be verified. The catches of narrow-barred Spanish mackerel increased from around 50,000 t the late-1970's to over 100,000 t by the late-1990's. The highest catches of narrow-barred Spanish mackerel were recorded in 2012, at over 160,000 t. Narrow-barred Spanish mackerel is caught in both Indian Ocean basins, with approximately equal proportions of catches recorded in the East and West Indian Ocean since the mid-2000s.

In recent years, the countries attributed with the highest catches of narrow-barred Spanish mackerel are Indonesia (29%) and India (23%) and, to a lesser extent, I.R. Iran, Myanmar, the UAE and Pakistan (25%) (Fig. 3).

**Fig. 2.** Narrow-barred Spanish mackerel: Annual catches of narrow-barred Spanish mackerel by gear recorded in the IOTC database (1950–2013). (Data as of October 2014)

² The uncertainty in the catch estimates has been assessed by the Secretariat and is based on the amount of processing required to account for the presence of conflicting catch reports, the level of aggregation of the catches by species and or gear, and the occurrence of non-reporting fisheries for which catches had to be estimated




Fig. 3. Narrow-barred Spanish mackerel: Average catches in the Indian Ocean over the period 2010–12, by country. Countries are ordered from left to right, according to the importance of catches of narrow-barred Spanish mackerel reported. The red line indicates the (cumulative) proportion of catches narrow-barred Spanish mackerel for the countries concerned, over the total combined catches of this species reported from all countries and fisheries. (Data as of October 2014)

Narrow-barred Spanish mackerel – uncertainty of catches

Retained catches are uncertain (Fig. 4), notably for the following fisheries:

- Artisanal fisheries of Indonesia and India: Indonesia and India have only recently reported catches of Spanish mackerel by gear, including catches by gear for the years 2005–08 and 2007–08, respectively. In the past, the IOTC Secretariat used the catches reported in recent years to break the aggregates for previous years, by gear and species. However, in a review conducted by the IOTC Secretariat by an independent consultant in 2012 the catches of narrow-barred Spanish mackerel were reassigned by gear. In recent years, the catches of narrow-barred Spanish mackerel estimated for Indonesia and India component represent around 50% of the total catches of this species in recent years.
- Artisanal fisheries of Madagascar: To date, Madagascar has not reported catches of narrow-barred Spanish mackerel to the IOTC. During 2012 the IOTC Secretariat conducted a review aiming to break the catches recorded in the FAO database as narrow-barred Spanish mackerel by species, on the assumption that all catches of tunas and tuna-like species had been combined under this name (the review used data from various sources including a reconstruction of the total marine fisheries catches of Madagascar (1950–2008), undertaken by the Sea Around Us Project). The new catches estimated are thought to be very uncertain.
- Artisanal fisheries of Somalia: Catch levels are unknown.
- Other artisanal fisheries UAE do not report catches of narrow-barred Spanish mackerel by gear. Although most of the catches are believed to be taken by gillnets, some narrow-barred Spanish mackerel may be also caught by using small surrounding nets, lines or other artisanal gears. In addition, Thailand report catches of narrow-barred Spanish mackerel and Indo-Pacific king mackerel aggregated.
- All fisheries: In some cases the catches of seerfish species are mislabelled, the catches of Indo-Pacific king mackerel and, to a lesser extent, other seerfish species, labelled as narrow-barred Spanish mackerel. Similarly, the catches of wahoo in some longline fisheries are thought to be mislabelled as narrow-barred Spanish mackerel. This mislabelling is thought to have little impact in the case of the narrow-barred Spanish mackerel but may be important for other seerfish species.
- Discard levels are believed to be low although they are unknown for most fisheries.
- Changes to the catch series: There have been no major revisions to the catch series of narrow-barred Spanish mackerel since the WPNT meeting in 2013.




Fig. 4. Narrow-barred Spanish mackerel: Uncertainty of annual catch estimates (1950–2013). Catches are assessed against IOTC reporting standards, where a score of 0 indicates catches that are fully reported according to IOTC standards; catches assigned a score of between 2 – 6 do not report catch data fully by gear and/or species (i.e., partially adjusted by gear and species by the IOTC Secretariat) or any of the other reasons provided in the document; catches with a score of 8 refer to fleets that do not report catch data to the IOTC (estimated by the IOTC Secretariat). (Data as of October 2014)

Narrow-barred Spanish mackerel – Effort trends

Effort trends are unknown for narrow-barred Spanish mackerel in the Indian Ocean.

Narrow-barred Spanish mackerel – Catch–per–unit–effort (CPUE) trends

Catch-and-effort series are available from some fisheries but they are considered highly incomplete (Table 5). In most cases catch-and-effort data are only available for short periods. Reasonably long catch-and-effort data series (extending for more than 10 years) are only available for Sri Lanka gillnets (Fig. 5). The catches and effort recorded are, however, thought to be unrealistic due to the dramatic changes in CPUE recorded in 2003 and 2004.

TABLE 5. Narrow-barred Spanish mackerel: Availability of catches and effort series, by fishery and year (1970–2013). Note that no catches and effort are available at all for 1950–84.

Gear-Fleet	70	72	74	76	78	80	82	84	86	88	90	92	94	96	98	00	02	04	06	08	10	12	
PSS-Indonesia																							
PSS-Malaysia																							
GILL-Indonesia																							
GILL-Sri Lanka																							
GILL-Malaysia																							
GILL-Oman																							
GILL-Pakistan																							
LINE-Australia																							
LINE-Malaysia																							
LINE-Oman																							
LINE-Yemen																							
LINE-South Africa																							
OTHR-Sri Lanka																							
OTHR-Indonesia																							
OTHR-Malaysia																							
OTHR-Oman																							




Fig. 5. Narrow-barred Spanish mackerel: Nominal CPUE series for the gillnet fishery of Sri Lanka derived from the available catches and effort data (1994–2004)

Narrow-barred Spanish mackerel – Fish size or age trends (e.g. by length, weight, sex and/or maturity)

- The size of narrow-barred Spanish mackerel taken by the Indian Ocean fisheries typically ranges between 30 and 140 cm depending on the type of gear used, season and location (Fig. 6). The size of narrow-barred Spanish mackerel taken varies by location with 32–119 cm fish taken in the Eastern Peninsular Malaysia area, 17–139 cm fish taken in the East Malaysia area and 50–90 cm fish taken in the Gulf of Thailand. Similarly, narrow-barred Spanish mackerel caught in the Oman Sea are typically larger than those caught in the Persian Gulf.
- Trends in average weight can only be assessed for Sri Lankan gillnets (from the late-1980s until the early 1990s), and Iranian gillnets from the late 2000s (Fig. 6, Table 6). The length frequency data available from the mid-eighties to the early nineties was obtained with the support of the IPTP (Indo-Pacific Tuna Programme); unfortunately, data collection did not continue after the IPTP activities came to an end.
- Catch-at-Size data are not available for the narrow-barred Spanish mackerel due to the paucity of size data available from most fleets (Table 6) and the uncertain status of the catches for this species (Fig. 4). Length distributions derived from the data available for gillnet fisheries are shown in Fig. 6. No data available for all other fisheries.

TABLE 6. Narrow-barred Spanish mackerel: Availability of length frequency data, by fishery and year (1980–2013)³. Note that no length frequency data are available at all for 1950–84.

Gear-Fleet	80	82	84	86	88	90	92	94	96	98	00	02	04	06	08	10	12	
PSS-Sri Lanka												■	■					
PSS-Thailand													■	■				
GILL-Oman				■	■	■	■	■	■	■						■		
GILL-Pakistan				■	■	■	■	■	■	■	■	■	■	■	■	■	■	
GILL-Sri Lanka				■	■	■	■	■	■	■	■	■	■	■	■	■	■	
GILL-Iran																■	■	
LINE-Iran																■	■	
LINE-Oman																■	■	
LINE-Sri Lanka									■	■	■	■	■	■	■	■	■	
OTHR-Saudi Arabia				■	■	■	■	■	■	■								
OTHR-Sri Lanka													■	■				

Key

- More than 2,400 specimens measured
- Between 1,200 and 2,399 specimens measured
- Less than 1,200 specimens measured

³ Note that the above list is not exhaustive, showing only the fisheries for which size data are available in the IOTC database. Furthermore, when available size data may not be available throughout the year existing only for short periods




Fig. 6. Narrow-barred Spanish mackerel: Left - Narrow-barred Spanish mackerel: Length frequency distributions for gillnet fisheries (total amount of fish measured by 1cm length class) derived from data available at the IOTC Secretariat. Right - number of narrow-barred Spanish mackerel specimens sampled for lengths, by fleet (gillnet only).

STOCK ASSESSMENT

Two assessment approaches were applied to narrow-barred Spanish mackerel in 2014, a Stock Reduction Analysis technique and a Posterior Focused Catch Reduction (PFCRA) method. The trajectories for both approaches were very similar and gave similar outcomes, and for reporting and stock status advice the PFCRA approach is used, as it was statistically robust (Table 7).

Noting that the Commission adopted Resolution 12/01 On the implementation of the precautionary approach, which effectively means that in a situation of increased uncertainty (e.g. data poor situations), a more precautionary approach should be undertaken when developing advice and possible management actions, this approach, combined with the weight-of-evidence available (stock status indicators from data poor assessment approaches, species biology, fishery indicators), were used to determine stock status for narrow-barred Spanish mackerel.

The stock status management advice for narrow-barred Spanish mackerel is based on the catch-based stock reduction method, combined with the known species and fishery attributes for status interpretation purposes. The approach presented is useful to assess stock status in the near term, while more traditional stock assessment approaches in the region are deferred until more data is collected and submitted in accordance with the IOTC data recording and reporting requirements for neritic tunas.

TABLE 7. Narrow-barred Spanish mackerel (*Scomberomorus commerson*) stock status summary

Management Quantity	Aggregate Indian Ocean
2013 catch estimate	148,695 t
Mean catch from 2009–2013	144,462 t
MSY (80% CI)	137 Kt [93–164 Kt]
Data period used in assessment	–
F_{MSY}	0.47 (0.41–1.95)
B_{MSY}	229 Kt (132–265K t)
F_{2012}/F_{MSY} (80% CI)	0.92 (0.41–1.95)
B_{2012}/B_{MSY} (80% CI)	1.17 (0.5–1.51)
SB_{2012}/SB_{MSY}	–
B_{2012}/B_0 (80% CI)	0.59 (0.25–0.75)
SB_{2012}/SB_0	–
$B_{2012}/B_0, F=0$	–
$SB_{2012}/SB_0, F=0$	–

LITERATURE CITED

- Darvishi M, Kaymaram F, Salarpouri A, Behzadi S, Daghooghi B (2011) Population dynamic and biological aspects of *Scomberomorus commerson* in the Persian Gulf and Oman Sea (Iranian coastal). IOTC–2011–WPNT01–23
- Froese R, Pauly DE (2009) FishBase, version 02/2009, FishBase Consortium, www.fishbase.org
- Grandcourt EM, Al Abdessalaam TZ, Francis F, Al Shamsi AT (2005) Preliminary assessment of the biology and fishery for the narrow-barred Spanish mackerel, *Scomberomorus commerson* (Lacépède, 1800), in the southern Arabian Gulf Fish Res 76:277–290
- McPherson GR (1989) North-eastern Australian mackerel (*Scomberomorus*) fishery. In: Chavez EA, Ed). Proceedings of the workshop Australia-Mexico on marine sciences, Quintana Roo, Mexico, 6-17 July 1989. Quintana Roo, Mexico. pp. 341-348