

## Length & weight conversion equations used for IOTC species

Equations used to convert from non-standard measurement to fork length (Table 1) and from fork length to round weight (Table 2)

**Table 1: Regression equations used to convert from non-standard measurements into standard lengths (tunas: tip of the snout to fork length; swordfish: lower-jaw to fork length) per species**

Species: <i>Yellowfin tuna</i>				Standard length: Tip of snout to fork of tail				
Type Measurement	Equation	Parameters	Sample size	Size	Variance	Covariance ab	Mean Residual	Gradient
Weight gilled and gutted <sup>A</sup>	$a*W^b$	a= 44.28699 b= 0.3008591	2,361	Min:14 Max:71	a=0.00752476509 b=2.86244E-07	-4.626246E-05	4.095958	a=3.033852 b=495.6385
Length to the base of the 1 <sup>st</sup> dorsal fin <sup>B</sup>	$a*L^b$	a=2.0759 b=1.1513	7,036	Min: 29 Max: 164				
Species: <i>Bigeye tuna</i>				Standard length: Tip of snout to fork of tail				
Type Measurement	Equation	Parameters	Sample size	Size	Variance	Covariance ab	Mean Residual	Gradient
Weight gilled and gutted <sup>A</sup>	$a*W^b$	a= 42.2186 b= 0.3012349	316	Min:12 Max:107	a=0.0321755341 b=1.299934E-06	-0.0002034041	3.98137	a=3.03806 b=473.1455
Length to the base of the 1 <sup>st</sup> dorsal fin <sup>C</sup>	$\frac{(L+a)^2}{(b)^2}$	a=21.45108 b=5.28756	2,858	Min:13 Max:48				
Species: <i>Albacore</i>				Standard length: Tip of snout to fork of tail				
Type Measurement	Equation	Parameters	Sample size	Size	Variance	Covariance ab	Mean Residual	Gradient
Weight round <sup>D</sup>	$(W/a)^{(1/b)}$	a= 0.000013718 b= 3.0973	n/a	n/a				
Species: <i>Swordfish</i>				Standard length: Tip of lower-jaw to fork of tail				
Type Measurement	Equation	Parameters	Sample size	Size	Variance	Covariance ab	Mean Residual	Gradient
Cleithrum-Fork length <sup>E</sup>	$(L+b)/a$	a= 0.8087 b= 8.6712	n/a	n/a				
Cleithrum-keel length <sup>F</sup>	$(a*L)+b$	a= 1.55108 b= 13.5025	179	Min:88 Max:252				
Eye orbit-fork length <sup>G</sup>	$(a*L)+b$	a= 1.066 b= 10.449	123	Min:48 Max:255				
Pectoral-anal length <sup>J</sup>	$(a*L)+b$	a= 2.5407 b= 25.698	1,806	Min:18 Max:105				
Pectoral-fork length <sup>K</sup>	$(a*L)+b$	a= 1.2398 b= 11.204	55	Min:60 Max:157				
Weight round <sup>L</sup>	$(W/a)^{(1/b)}$	a= 0.000003815 b= 3.188	3,608	Min:89 Max:266				
Weight gilled and gutted <sup>H</sup>	$(W/a)^{(1/b)}$	a= 0.0000043491 b= 3.188	3,608	Min:89 Max:266				
Weight headed <sup>I</sup>	$(W/a)^{(1/b)}$	a= 0.000004592 b= 3.137	n/a	n/a				

**A:** Data from Penang Sampling Programme (1992-93)  
**B:** Data from the Indian Ocean (Marsac, F. et al in IOTC-2006-WPTT-09)  
**C:** Data from the Atlantic Ocean, Champagnat et Pianet (1974) (ibid. B)  
**D:** Inverted length-weight relationship; ICCAT South Atlantic  
**E:** Reference not available (to be provided)  
**F:** Two step conversion as CKL = (0.690253\*EFL) -3.541823 in formula LJFL = 8.00884+(1.07064\*EFL); NOAA Data (Pacific Ocean)  
**G, J, K:** Data from Reunion Island, Indian Ocean Poisson 2001 (in IOTC-2005-WPTT-05)  
**H:** Inverted length-weight equation (ICCAT Mejuto et al 1998 South-East Atlantic Ocean)  
**I:** Inverted length-weight equation; Reference not available (to be provided)  
**L:** Converted to GGT (GGT=RND/1.14 (Mejuto et al. 1998)) and inverted length-weight equation (ICCAT Mejuto et al 1998 South-East Atlantic Ocean)

**Table 1(cont): Regression equations used to convert from non-standard measurements into standard lengths (eye orbit to fork length), per species**

Species: <i>Black marlin</i>				Standard length: Eye orbit to fork of tail				
Type Measurement	Equation	Parameters	Sample size	Size	Variance	Covariance ab	Mean Residual	Gradient
Cleithrum-Keel length	No equation available							
Lower-jaw - fork length <sup>M</sup>	(a*L)+b	a= 0.8972 b= -4.6673	13	Min:119 Max:314				
Weight gilled and gutted <sup>N</sup>	a*L <sup>b</sup>	a= 41.56681 b= 0.309442	24	Min:8.6 Max:279				
Species: <i>Blue marlin</i>				Standard length: Eye orbit to fork of tail				
Type Measurement	Equation	Parameters	Sample size	Size	Variance	Covariance ab	Mean Residual	Gradient
Lower-jaw - fork length <sup>O</sup>	(a*L)+b	a= 0.9039 b= -7.248	26	Min:143 Max:295				
Weight gilled and gutted <sup>P</sup>	a*L <sup>b</sup>	a= 46.0356637 b= 0.283377	154	Min:10 Max:381				
Species: <i>Striped marlin</i>				Standard length: Eye orbit to fork of tail				
Type Measurement	Equation	Parameters	Sample size	Size	Variance	Covariance ab	Mean Residual	Gradient
Lower-jaw - fork length <sup>Q</sup>	(L+b)/a	a= 1.1178 b= -7.769		Min: Max:				
Weight round <sup>R</sup>	a*L <sup>b</sup>	a= 51.3506 b= 0.300417	1427	Min:7 Max:100				
Weight gilled and gutted <sup>S</sup>	a*L <sup>b</sup>	a= 45.443009 b= 0.300417	1427	Min:7 Max:100				
Species: <i>Indo-Pacific sailfish</i>				Standard length: Eye orbit to fork of tail				
Type Measurement	Equation	Parameters	Sample size	Size	Variance	Covariance ab	Mean Residual	Gradient
Cleithrum-Keel length	No equation available							
Lower-jaw - fork length <sup>T</sup>	(L+b)/a	a= 0.8845 b= -3.7025	1166	Min:78 Max:232				
Weight gilled and gutted <sup>U</sup>	a*L <sup>b</sup>	a= 45.5076 b= 0.347166	35	Min:5 Max:38				
<p><b>M:</b> BRS (Ward, pers.com.) Eastern and western Australia (on IOTC-2005-WPTT-05)</p> <p><b>N:</b> PIFSC Administrative report: (Updated Weight-on-Length Relationships for Pelagic Fishes Caught in the Central North Pacific Ocean and Bottom fishes from the Northwestern Hawaiian Islands). With value of a (46.9705) divided by 1.13 to account for conversion of gilled-and-gutted weight into round weight</p> <p><b>O:</b> BRS (Ward, pers.com.) Eastern and western Australia (on IOTC-2005-WPTT-05)</p> <p><b>P:</b> PIFSC Administrative report: (Updated Weight-on-Length Relationships for Pelagic Fishes Caught in the Central North Pacific Ocean and Bottom fishes from the Northwestern Hawaiian Islands) Value of a (52.0203) divided by 1.13 to account for conversion of gilled-and-gutted weight into round weight</p> <p><b>Q:</b> Su, N.J., C.L. Sun, S.Z. Yeh, W.C. Chiang, S.P. Wang, and C.H. Liu (2005) LJFL and EFL relationships for the billfishes caught by the Taiwanese offshore and coastal fisheries; (ISC/05/MAR&amp;SWO-WGs/ 4) inverted equation</p> <p><b>R:</b> PIFSC Administrative report: (Updated Weight-on-Length Relationships for Pelagic Fishes Caught in the Central North Pacific Ocean and Bottom fishes from the Northwestern Hawaiian Islands)</p> <p><b>S:</b> PIFSC Administrative report: (Updated Weight-on-Length Relationships for Pelagic Fishes Caught in the Central North Pacific Ocean and Bottom fishes from the Northwestern Hawaiian Islands) Value of a (51.3506) divided by 1.13 to account for conversion of gilled-and-gutted weight into round weight</p> <p><b>T:</b> Wei-Chuan Chiang et al. , 2004; inverted EFL-FL equation (M+F sexes pooled)</p> <p><b>U:</b> PIFSC Administrative report: (Updated Weight-on-Length Relationships for Pelagic Fishes Caught in the Central North Pacific Ocean and Bottom fishes from the Northwestern Hawaiian Islands) Value of a (51.4235) divided by 1.13 to account for conversion of gilled-and-gutted weight into round weight</p>								

**Table 2: Equations used to convert from standard length into round weight, per species**

Species	Gear Type/s	From type measurement – To type measurement	Equation	Parameters	Sample size	Length
Yellowfin tuna	Purse seine Pole and Line Gillnet	Fork length – Round Weight(kg) <sup>A</sup>	$RND=a*L^b$	a= 0.00001886 b= 3.0195	6,752	Min: 29 Max: 164
	Longline Line Other Gears	Fork length(cm) – Gilled and gutted weight(kg) <sup>B</sup> Gilled and gutted weight(kg) - Round Weight(kg) <sup>C</sup>	$GGT=a*L^b$ $RND=GGT*1.13$	a= 0.0000094007 b= 3.126843987	15,133	Min:72 Max:177
Bigeye tuna	Purse seine Pole and Line Gillnet	Fork length(cm) – Round Weight(kg) <sup>D</sup>	$RND=a*L^b$	a= 0.000027000 b= 2.95100	n/a	n/a
	Longline Line Other Gears	Fork length(cm) – Gilled and gutted weight(kg) <sup>B</sup> Gilled and gutted weight(kg) - Round Weight(kg) <sup>C</sup>	$GGT=a*L^b$ $RND=GGT*1.13$	a= 0.0000159207 b= 3.0415414023	12,047	Min:70 Max:187
Skipjack tuna	All gears	Fork length(cm) – Round Weight(kg) <sup>E</sup>	$RND=a*L^b$	a= 0.0000074800 b= 3.25260	14,140	Min:32 Max:78
Albacore	All gears	Fork length(cm) – Round Weight(kg) <sup>F</sup>	$RND= a*L^b$	a= 0.0000137180 b= 3.0973	n/a	n/a
Swordfish	All gears	Tip of lower-jaw to fork of tail(cm) – Round Weight(kg) <sup>G</sup>	$RND= a*L^b$	a= 0.0000042030 b= 3.21340	2569	Min:80 Max:253
Black marlin	All gears	Eye orbit to fork of tail(cm) – Round Weight(kg) <sup>H</sup>	$RND= a*L^b$	a= 0.0000144217 b= 2.98851	24	Min:95 Max:279
Blue marlin	All gears	Eye orbit to fork of tail(cm) – Round Weight(kg) <sup>H</sup>	$RND= a*L^b$	a= 0.00000272228 b= 3.30967	154	Min:109 Max:269
Striped marlin	All gears	Eye orbit to fork of tail(cm) – Round Weight(kg) <sup>H</sup>	$RND= a*L^b$	a= 0.00000133263 b= 3.41344	17	Min:101 Max:178
Indo-Pac. sailfish	All gears	Eye orbit to fork of tail(cm) – Round Weight(kg) <sup>H</sup>	$RND= a*L^b$	a= 0.0000690103 b= 2.52429	35	Min:86 Max:187

**A: Data from the Indian Ocean (Marsac, F. et al in IOTC-2006-WPTT-09)**

**B: Multilateral catch monitoring Bena (2002-04)**

**C: ICCAT Field Manual (Appendix 4: Population parameters for key ICCAT species. Product Conversion Factors)**

**D: Cort (1986)**

**E: Data from the Atlantic Ocean, Cayré et Laloë (Fonteneau, A. et J. Marcille (eds), 1988: Ressources, pêche et biologie des thonidés tropicaux de l'Atlantique Centre-Est. FAO Doc.Tech.Pêches, (292), page262)**

**F: Data from the South Atlantic (Penney 1994)**

**G: Data from the Atlantic Ocean, Spanish longline fishery (Mejuto et al., 1988, ICCAT)**

**H: PIFSC Administrative report: (Updated Weight-on-Length Relationships for Pelagic Fishes Caught in the Central North Pacific Ocean and Bottom fishes from the Northwestern Hawaiian Islands)**

**Table 3: SHARKS: Equations used to convert from standard length into round weight, per species**

Species	Gear Type/s	From type measurement – To type measurement	Equation	Parameters	Sample size	Length
Blue shark	All gears	Fork length – Round Weight(kg) <sup>A</sup>	$RND=a*L^b$	a= 0.0000031841 b= 3.1313		Min: Max:
	All gears	Precaudal length – Fork Length <sup>C</sup>	$FL=\frac{PCL+a}{b}$	a= 0.3956 b= 0.9075		Min: Max:
	All gears	Total length – Fork length <sup>D</sup>	$FL=a*TL+b$	a= 0.822 b= 1.754	10	Min:217 Max:265
Shortfin Mako Shark	All gears	Fork length – Round Weight <sup>A</sup>	$RND=a*L^b$	a= 0.0000052432 b= 3.1407		Min: Max:
	All gears	Precaudal length – Fork Length <sup>C</sup>	$FL=a*PCL+b$	a= 1.100 b= 0.766	n/a	n/a
	All gears	Total length – Fork length	$FL=a*TL+b$	a= 0.9286 b= -1.7101		Min: Max:
Oceanic whitetip shark	All gears	Fork length – Round Weight <sup>B</sup>	$RND= a*L^b$	a= 0.000018428 b= 2.9245		Min: Max:
	All gears	Total length – Fork length	$FL=a*TL+b$	a= 0.7847 b= 1.211		Min: Max:
Porbeagle	All gears	Fork length – Round Weight <sup>A</sup>	$RND=a*L^b$	a= 0.000014823 b= 2.9641		
	All gears	Precaudal length – Fork Length <sup>C</sup>	$FL=a*PCL+b$			
Silky Shark	All gears	Fork length – Round Weight <sup>A</sup>	$RND=a*L^b$	a= 0.000015406 b= 2.9221		
Bigeye Thresher Shark	All gears	Fork length – Round Weight <sup>A</sup>	$RND=a*L^b$	a= 0.0000091069 b= 3.0802		
Thresher Shark	All gears	Fork length – Round Weight <sup>A</sup>	$RND=a*L^b$	a= 0.00018821 b= 2.5188		

**A:** Data from Western North Atlantic: NOAA Technical Memorandum NMFS-NE-110 (May 1996)

**B:** Data from Indian Ocean: Length-weight relationships, conversion factors and analyses of sex-ratio, by length-range, Observers onboard Spanish Longliners in South Western Indian Ocean during 2005; Ariz J, A Delgado de Molina, M.L Ramos, J.C Santana

**C:** Inverse equation from WCPFC Analysis of North Pacific Shark Data from Japanese Commercial Longline and Research/Training Vessels Record; Shelley Clarke, Kotaro Yokawa, Hiroaki Matsunaga and Hideki Nakano

**D:** Indian Ocean : Maldivian Oceanic Shark Longline Fishery; R.C. Anderson, M.S. Adam & M. Saleem

**E:** Data from the Atlantic Ocean, Cayré et Laloë (Fonteneau, A. et J. Marcille (eds), 1988: Ressources, pêche et biologie des thonidés tropicaux de l'Atlantique Centre-Est. FAO Doc.Tech.Pêches, (292), page262)

**F:** Data from the Indian Ocean, Taiwanese gillnet fishery (Chien-Chung Hsu)

**G:** Data from the Atlantic Ocean, Spanish longline fishery (Mejuto et al., 1988, ICCAT)

**H:** PIFSC Administrative report: (Updated Weight-on-Length Relationships for Pelagic Fishes Caught in the Central North Pacific Ocean and Bottom fishes from the Northwestern Hawaiian Islands)