


## DRAFT RESOURCE STOCK STATUS SUMMARY – BIGEYE TUNA



### Status of the Indian Ocean bigeye tuna (BET: *Thunnus obesus*) resource

**TABLE 1.** Bigeye tuna: Status of bigeye tuna (*Thunnus obesus*) in the Indian Ocean.

Area <sup>1</sup>	Indicators		2019 stock status <sup>3</sup> determination
Indian Ocean <sup>5</sup>	Catch in 2018 <sup>2</sup> :	93,515 t (81,413 t) <sup>4</sup>	<b>38.2%*</b>
	Average catch 2014–2018:	92,140 t (89,720 t) <sup>4</sup>	
MSY (1,000 t) (80% CI):	87 (75-108)		
F <sub>MSY</sub> (80% CI):	0.24 (0.18-0.36)		
SB <sub>MSY</sub> (1,000 t) (80% CI):	503 (370-748)		
F <sub>2018</sub> /F <sub>MSY</sub> (80% CI):	1.20 (0.70-2.05)		
SB <sub>2018</sub> /SB <sub>MSY</sub> (80% CI):	1.22 (0.82-1.81)		
SB <sub>2018</sub> /SB <sub>0</sub> (80% CI):	0.31 (0.21. – 0.34)		

<sup>1</sup> Boundaries for the Indian Ocean stock assessment are defined as the IOTC area of competence.

<sup>2</sup> Proportion of catch estimated or partially estimated by IOTC Secretariat for catches in 2018: 28%

<sup>3</sup> The stock status refers to the most recent years' data used in the last assessment conducted in 2019.

<sup>4</sup> Considering the alternative purse seine log-associated catch composition for the EU fleet in 2018 as per IOTC-2019-WPTT21-R[E].

<sup>5</sup> Results of management quantities presented here are for the revised catches – see footnote 4.

\* Estimated probability that the stock is in the respective quadrant of the Kobe Plot (shown below), derived from the confidence intervals associated with the current stock status. The confidence intervals for SB<sub>2015</sub>/SB<sub>0</sub> were not estimated for the models used.

Colour key	Stock overfished (SB <sub>year</sub> /SB <sub>MSY</sub> < 1)	Stock not overfished (SB <sub>year</sub> /SB <sub>MSY</sub> ≥ 1)
Stock subject to overfishing (F <sub>year</sub> /F <sub>MSY</sub> > 1)	34.6	38.2%
Stock not subject to overfishing (F <sub>year</sub> /F <sub>MSY</sub> ≤ 1)	0%	27.2%
Not assessed/Uncertain		

### INDIAN OCEAN STOCK – MANAGEMENT ADVICE

**Stock status.** In 2019 a new stock assessment was carried out for bigeye tuna in the IOTC area of competence to update the stock status undertaken in 2016. Two models were applied to the bigeye stock (JABBA and Stock Synthesis (SS3)). The stock assessment selected to provide scientific advice was carried out using SS3, a fully integrated model used to provide scientific advice for the three tropical tunas stocks in the Indian Ocean. The reported stock status is based on the SS3 model formulation using a grid of 18 model configurations designed to capture the uncertainty on stock recruitment relationship, the influence of tagging information and selectivity of longline fleets. Due to concerns on the reported catch data for

2018, the stock status is based on SS3 model formulations using the best catch estimate by the Scientific Committee (for details see WPTT report). Spawning stock biomass in 2018 was estimated to be 31% of the unfished levels in 2018 (Table 1) and 122% (82–181%) of the level that can support MSY. The assessment outcome is qualitatively different to the stock assessment conducted in 2016 due to the increase of catch of small size, changes in modelling assumptions about longline selectivity, and the abundance index developed in 2019. Considering the characterized uncertainty, the assessment indicates that  $SB_{2018}$  is above  $SB_{MSY}$  with high probability (65.4%) and that fishing mortality is above  $F_{MSY}$  also with high probability (72.8%). The median value of MSY from the model runs presented with SS3 was 87,000 t with a range between 75,000 and 108,000 t (a median level 16% lower than the estimate in 2016). Catches in 2018 ( $\approx 81,413$  t) remain lower than the estimated median MSY values from the stock assessment conducted in 2019 but within the range of estimated MSY. The average catch over the previous five years (2014–18;  $\approx 89,717$  t) is just above the estimated median MSY and within the range of estimated values. Thus, on the weight-of-evidence available in 2019, the bigeye tuna stock is determined to be **not overfished** but **subject to overfishing** (Table 1).

**Outlook.** Declines in longline effort since 2007, particularly from the Japanese, Taiwanese and Rep. of Korea longline fleets lowered the pressure on the Indian Ocean bigeye tuna stock since 2007. However, recent increase in catch from purse seine fleets have increased this pressure and the stock is estimated to be subject to overfishing. The estimated MSY has declined significantly (16%) from the previous estimate (from 2016) due to the increase of purse seine catch in the overall change in catch composition, changes in modelling assumptions about longline selectivity, and the inclusion of a more pessimistic abundance index in the western tropical region. The Kobe strategy matrix (K2SM) based on the plausible model runs from SS3 in 2019 illustrates the levels of quantified risk associated with varying catch levels over time that could be used to inform future management actions (Table 2). The projections produced to estimate the K2SM (Table 2) are, in the short term, driven by the below average recruitment estimated for the recent years. The SS3 projections from the 2019 assessment show that there is a risk of breaching MSY-based reference points by 2021, and 2028 if catches are maintained at 2018 levels at the current selectivity and therefore size distribution of catch (Table 2). Should the management objective of maintaining biomass at levels higher than  $SB_{MSY}$  with more than 50% probability in 2028 be pursued, the overall catch should be reduced 10% from current levels (73,272 t).

**Management advice.** The stock status determination changed qualitatively in 2019 to **not overfished** but **subject to overfishing**. If catches remain at current levels there is a risk of breaching MSY reference points with 58.9% and 60.8% probability in 2021 and 2028. Reduced catches of at least 10% from current levels will likely reduce the probabilities of breaching reference levels to 49.1% in 2028. Continued monitoring and improvement in data collection, reporting and analyses is required to reduce the uncertainty in assessments (Table 2).

The following key points should also be noted:

- **Maximum Sustainable Yield (MSY):** estimate for the Indian Ocean stock is 87,000 t with a range between 75,000–108,000 t for SS3 (Table 1). The average 2014–2018 catches of  $\approx 89,717$  t, and catches for each year since 2012 are within the range of the estimated MSY level.
- **Interim reference points:** Noting that the Commission in 2015 agreed to Resolution 15/10 *on target and limit reference points and a decision framework*, the following should be noted:
  - **Fishing mortality:** Current fishing mortality is considered to be at 120% of the interim target reference point of  $F_{MSY}$ , and 92% of the interim limit reference point of  $1.3 * F_{MSY}$  (Fig. 2).

- **Biomass:** Current spawning biomass is considered to be at 122% of the interim target reference point of  $SB_{MSY}$  and well above the interim limit reference point of  $0.5 \cdot SB_{MSY}$  (Fig. 2).
- **Main fishing gear** (Average catch 2014–18): Longline  $\approx 42\%$ ; Purse seine  $\approx 31\%$  (FAD associated school (LS)  $\approx 24\%$ ; free swimming school (PS)  $\approx 7\%$ ); All other (artisanal) gears  $\approx 27\%$  (Fig 1).
- **Main fleets** (Average catch 2014–18): Indonesia  $\approx 25\%$ ; Taiwan, China  $\approx 16\%$ ; European Union  $\approx 20\%$  (EU-Spain:  $\approx 15\%$ ; EU-France:  $\approx 5\%$ ); Seychelles  $\approx 13\%$ .





Fig. 1(a-b). Annual catches of bigeye tuna by gear (1950–2018). Data as of October 2019.



**Fig. 2.** . Bigeye tuna: SS3 Aggregated Indian Ocean assessment Kobe plot. The coloured points represent stock status estimates from the 18 model options. The grey dots represent 5000 estimates of 2018 stock status from the multivariate normal approximation from the mean and variance-covariance of the 18 model options. The legend indicates the estimated probability of the stock status being in each of the Kobe quadrant. The white circle (around the purple dot) represents the median stock status in 2018.

**TABLE 2.** Bigeye tuna: Stock Synthesis base case Indian Ocean assessment Kobe II Strategy Matrix. Probability (percentage) of violating the MSY-based target (top) and limit (bottom) reference points for constant catch projections (relative to average catch level from 2018 (81,413 t); -10%, -20%, -30%, -40%) projected for 3 and 10 years.

Reference point and projection timeframe	Alternative catch projections (relative to the catch level from 2018) and weighted probability (%) scenarios that violate reference point				
	60% (48,848t)	70% (56,990t)	80% (65,130t)	90% (73,272t)	100% (81,413t)
$B_{2021} < B_{MSY}$	51.1	53.3	54.2	57.1	58.9
$F_{2021} > F_{MSY}$	7.3	17.8	32	47.9	62.8
$B_{2028} < B_{MSY}$	8	19.5	35.1	49.1	60.8
$F_{2028} > F_{MSY}$	1.1	6.9	19.8	37.7	55.6
Reference point and projection timeframe	Alternative catch projections (relative to the catch level from 2018) and probability (%) of violating MSY-based limit reference points ( $B_{lim} = 0.5 B_{MSY}$ ; $F_{lim} = 1.3 F_{MSY}$ )				
	60%	70%	80%	90%	100%

	(48,848t)	(56,990t)	(65,130t)	(73,272t)	(81,413t)
$B_{2021} < B_{LIM}$	0	0	0	0	0
$F_{2021} > F_{LIM}$	6.0	11.0	17.0	28.0	39.0
$B_{2028} < B_{LIM}$	0.0	0.0	6.0	11.0	22.0
$F_{2028} > F_{LIM}$	0.0	6.0	17.0	22.0	39.0