

Japanese longline CPUE for yellowfin tuna in the Indian Ocean standardized by generalized linear model which includes cluster analysis

T. Matsumoto¹, K. Satoh² and
H. Yokoi²

¹*Fisheries Resources Institute, Japan Fisheries Research and Education Agency, 5-7-1, Orido, Shimizu, Shizuoka, 424-8633, Japan*

²*Fisheries Resources Institute, Japan Fisheries Research and Education Agency, 2-12-4, Fukuura, Kanazawa-ku, Yokohama-shi, 236-8648, Japa*

Abstract

Japanese longline CPUE for yellowfin tuna in the Indian Ocean (area specific) was standardized for 1975-2020 by GLM based using the method in the new CPUE collaborative analysis. Cluster analysis was conducted before standardization, and cluster number was used for main effect as well as year, quarter, vessel ID and five degree latitude/longitude block. Basically, standardized CPUEs showed similar trends among areas. CPUEs continuously decreased from 1970s to around mid 2010s, and those showed slight increasing trend after that. Trend of CPUEs was similar to those in the previous studies.

1. Introduction

Yellowfin tuna is one of main target species for Japanese longline fishery in the Indian Ocean. Its abundance indices are very important for stock assessment or stock indicator of this species. Yellowfin tuna is mainly caught in the tropical and subtropical areas especially in the western Indian Ocean (Matsumoto and Satoh, 2012; Matsumoto 2014). Since 2007, piracy activities off Somalia has increased and spread to whole northwestern Indian Ocean. Japanese longline effort in the Indian Ocean, especially in the northwestern part, has rapidly decreased to avoid the piracy attack. In the IOTC WPTT meeting in 2010, a concern about the effect of the decreased effort on the CPUE trend of the longline fishery was recognized. Okamoto (2011b) estimated the regional effect of the decreased longline effort on the CPUE trend in the Indian Ocean, and suggested that the decreased effort in northwestern Indian Ocean has no more been able to represent the CPUE trend in this region. Therefore, Okamoto (2011a) calculated CPUE trends for both scenarios including and excluding Area 2 (northwestern area) and found that the trends were similar. At 2012-2015 IOTC WPTT meetings, Matsumoto et al. (2012, 2013) and Ochi et al. (2014, 2015) conducted CPUE standardization by using area rate without northwest area because no effort was observed in this area in 2011 due to piracy activities, and the indices were used for stock assessment in 2012 and 2015. Matsumoto et al. (2016) also reported standardization of yellowfin tuna CPUE based on similar methods as those in the previous studies with additionally using the effect of LT1LN1 (1 degree latitude/longitude effect). They found that there was only small difference of CPUE between with LT5LN5 and with LT1LN1. Matsumoto et al. (2016) also relieved the concern that CPUE got higher as the number of hooks between floats (NHF) increases, which did not agree to expected result, by using LT5LN5 instead of subareas for the effect of fishing ground. Matsumoto (2017; 2018; 2019; 2020) reported similar results with updated data. In Matsumoto (2018), vessel effect was used for one of the effects (covariates) in the CPUE standardization using similar approach by Okamoto (2014), and found that it has some effect for CPUE trend.

In 2016, IOTC joint CPUE analysis (CPUE workshop) was conducted and 'joint CPUEs' were created for bigeye and yellowfin tuna, based on Japanese, Taiwanese and Korean longline operational data (Hoyle et al., 2016). These models account for fishing power based on vessel ID where available, and use cluster analysis to incorporate targeting. Joint CPUEs were considered to be more representative of status of the stocks and so were used for base models of stock assessment. At that time fleet-specific CPUE indices were prepared for Japanese longline using the same methods, but were not presented, so it was not possible to compare the joint and Japanese-only longline CPUE indices. In 2017 the joint CPUE analysis workshop was held and CPUE indices for each fleet as well as joint CPUE were created (Hoyle et al., 2017). Japanese longline CPUE for bigeye and yellowfin tuna created at that workshop was reported by Matsumoto et al. (2017). They reported that the trend of both CPUEs was mostly similar to those by traditional method, but there are some differences especially in the early period. Also in 2018 and 2019, joint CPUE analysis workshop was again held

and CPUE indices for each fleet as well as joint CPUE were created (e.g. Matsumoto et al., 2018).

A new collaborative study for developing the abundance index started in 2019 by Japanese, Korean and Taiwanese scientists has been conducted for analyzing CPUE for Indian Ocean yellowfin tuna (Kitakado et al., 2021). In this collaborative study, in addition to the methods similar to those mentioned above, cutting-edge methods such as VAST (advanced spatio-temporal model) will also be developed.

This document reports the standardization of Japanese longline CPUE for yellowfin tuna in the Indian Ocean using the similar methods as those for joint CPUE (Japanese, Korean and Taiwanese longline fishery, conventional method) in the new collaborative study reported by Kitakado et al. (2021). The results may help to compare the results with those by new collaborative study (joint CPUE) and to compare CPUEs among fleets derived using the same or similar methods.

2. Materials and methods

The methods to standardize CPUE are similar to conventional regression analyses in the CPUE collaborative study mentioned above (Kitakado et al., 2021).

Catch and effort data used:

Operational level (set by set) Japanese longline logbook data with vessel ID were used. The data were available for 1975-2020 (data for 2020 were preliminary). The data include the fields year, month and day of operation, location to 1° of latitude and longitude, vessel identifier (call sign and vessel registration number), number of hooks between floats (HBF), number of hooks per set, and catch in number of each species. In the previous collaborative studies, vessel ID was available from 1979, but currently the information for longer period (from 1975) is available.

Each set was allocated to subregion (subarea) (Fig. 1). Area 1 (northwest) was not used because there are not enough catch and effort. These regions are the same as those in the previous studies (e.g. Hoyle et al., 2018, Matsumoto et al., 2018), and also basically the same (except for northern and southern limits for region 1 and 3, respectively) as those for separating fleets or area for SS3 model in the previous stock assessment. Fig. 2 shows annual change in the proportion of fishing effort (number of operation) in each area. Fig. 3 shows species composition of the catch by longline fishery in each area.

Cluster analysis

We clustered the data using the approach described by Kitakado et al. (2021), which used Ward's minimum variance and the complete linkage methods. Species composition in number of the catch was aggregated for 10-days period (1st-10th, 11th-20th, and 21st~ for each month), and was used for cluster analysis. In the previous analyses (e.g. Hoyle et al., 2018), the data was aggregated for 1 month period, but shorter period was used in this study for better reflecting targeting. Catch for southern bluefin tuna (SBT), albacore (ALB), bigeye tuna (BET), yellowfin tuna (YFT), swordfish (SWO), sharks (SKX) and other fish (OTH) were used for species composition. Data were also clustered using the kmeans method, which minimises the sum of squares from points to the cluster centres.

GLM (Generalized Linear Model):

After cluster analysis, cluster numbers were assigned to catch and effort data aggregated by year, month, vessel ID and 1 degree latitude/longitude blocks. This data set was used for CPUE standardization.

GLM (generalized linear models) that assumed a lognormal distribution was conducted. In this approach the response variable $\log(\text{CPUE}+k)$ was used, and a normal distribution was assumed. The constant k , added to allow for modelling sets with zero catches of the species of interest, was 10% of the mean CPUE for all sets. CPUE was defined at the set level as catch in number divided by 1000 hooks. The following model was used:

$$\ln(\text{CPUEs}+k) \sim \text{year} + q + \text{vessel} + \text{latlon5} + \text{cluster} + \text{yr} * q + \epsilon$$

where *year*: effect of year, *q*: effect of quarter; *vessel*: effect of vessel ID; *latlon5*: effect of five degree latitude and longitude; *cluster*: effect of cluster; *year*q*: interaction between year and quarter; ϵ : error term, *k*: 10% of the mean CPUE

All the covariates were incorporated as fixed effect. As for diagnostics of CPUE standardization, residual distributions, Q-Q plots and influence plots were produced.

3. Results and discussion

Species compositions were plotted by cluster for each region (**Fig. 4**) and each region and year (**Fig. 5**). Dominant species differed depending on clusters, but there was at least one cluster in each region in which yellowfin tuna was dominant. Number of clusters were 4 or 5 for each region.

ANOVA indicates that as for all the regions all the effects were effective at 1% significance level (Table 1). Fig. 6 shows comparison of yellowfin CPUE by area, and Fig. 7 shows comparison of yellowfin CPUE in each area with nominal CPUE and standardized CPUE in the previous studies (Matsumoto et al. 2019, Matsumoto 2020). The trend of CPUE is similar among areas. CPUEs continuously decreased from 1970s to around mid 2010s, and those showed slight increasing trend after that. Very high jump of CPUE in 2020 was observed in area 2, which may be due to small number of operations (Fig. 2). The trend of CPUE in this study is similar to those in the previous studies, and there are some small scale differences.

Fig. 8 shows distribution of standardized residuals and QQ plots. It seems that the distributions are not largely skewed. Fig. 9 shows influence plots. In many cases there is historical change of the effect. Difference of historical change of the effect by area is also observed. For example, *latlon5* effect is increasing in region 3 and 5, although there is no clear trend in region 4.

4. References

- Hoyle, S., Chang, Y., Kim, D. N., Lee, S., Matsumoto, T., Satoh, K. and Yeh, Y. (2016) Collaborative study of tropical tuna CPUE from multiple Indian Ocean longline fleets in 2016. IOTC-2016-WPTT18-14.
- Hoyle S., Assan C., Chang, S. T., Fu, D., Govinden R., Kim D, N, Lee S. I., Lucas J., Matsumoto T. Satoh K., Yeh Y. M., and Kitakado T. (2017) Collaborative study of tropical tuna CPUE from multiple Indian Ocean longline fleets in 2017. IOTC-2017-WPTT19-32. 52p.
- Kitakado T., K. Satoh, SI Lee, NJ Su, T Matsumoto, H Yokoi, K Okamoto, MK Lee, JH Lim, Y Kwon, SP Wang, WP Tsai, ST Chang, and FC Chang. (2021) Report of trilateral collaborative study among Japan, Korea and Taiwan for producing joint abundance indices for the yellowfin tunas in the Indian Ocean using longline fisheries data up to 2019. IOTC-2021-WPTT23DP-14.
- Matsumoto, T. (2014): Review of Japanese fisheries and tropical tuna catch in the Indian Ocean. IOTC 2014/WPTT16/10. 28pp.
- Matsumoto, T. (2017) Japanese longline CPUE for yellowfin tuna in the Indian Ocean standardized by generalized linear model. IOTC-2017-WPTT19-48. pp 17.
- Matsumoto, T. (2018) Japanese longline CPUE for yellowfin tuna in the Indian Ocean standardized by generalized linear model. IOTC-2018-WPTT20-38. pp 19.
- Matsumoto, T. (2019) Japanese longline CPUE for yellowfin tuna in the Indian Ocean standardized by generalized linear model. IOTC-2019-WPTT21-46. pp 18.
- Matsumoto, T. (2020) Japanese longline CPUE for yellowfin tuna in the Indian Ocean standardized by generalized linear model. IOTC-2020 -WPTT22(DP) -INF04.
- Matsumoto, T. Okamoto, H. and Kitakado, T. (2012): Japanese longline CPUE for yellowfin tuna in the Indian Ocean

- up to 2011 standardized by general linear model. IOTC 2012/WPTT14/35. 34pp.
- Matsumoto, T. and Satoh, K. (2012): Review of Japanese fisheries and tropical tuna catch in the Indian Ocean. IOTC 2012/WPTT14/17. 28pp.
- Matsumoto, T., Okamoto, H. and Kitakado, T. (2013): Japanese longline CPUE for yellowfin tuna in the Indian Ocean up to 2012 standardized by generalized linear model. IOTC-2013-WPTT15-37, p. 43.
- Matsumoto, T., Nishida, H., Satoh, K and Kitakado, T. (2016): Japanese longline CPUE for yellowfin tuna in the Indian Ocean standardized by generalized linear model. IOTC-2016-WPTT18-25, p. 22.
- Matsumoto, T., Satoh, K. and Hoyle, S. (2017) Standardization of bigeye and yellowfin tuna CPUE by Japanese longline in the Indian Ocean which includes cluster analysis. IOTC-2017-WPTT19-29. pp 26.
- Matsumoto, T., Satoh, K. and Hoyle, S. (2018) Standardization of bigeye and yellowfin tuna CPUE by Japanese longline in the Indian Ocean which includes cluster analysis. IOTC-2018-WPTT20-37. pp 36.
- Ochi, D., Matsumoto, T., Okamoto, H. and Kitakado, T. (2014): Japanese longline CPUE for yellowfin tuna in the Indian Ocean up to 2013 standardized by generalized linear model. IOTC-2014-WPTT16-47, 37pp.
- Ochi, D., Matsumoto, T., Okamoto, H., T. Nishida and Kitakado, T. (2015): Update of standardized Japanese longline CPUE for yellowfin tuna in the Indian Ocean and consideration of standardization methods. IOTC-2014/WPTT16/26, 53pp.
- Okamoto, H. (2011a): Japanese longline CPUE for yellowfin tuna in the Indian Ocean up to 2010 standardized by general linear model. IOTC 2011/WPTT13/34. 45 pp.
- Okamoto, H. (2011b): Preliminary analysis of the effect of the Piracy activity in the northwestern Indian Ocean on the CPUE trend of bigeye and yellowfin. IOTC 2011/ WPTT13/44. 9pp.
- Okamoto, H. (2014): CPUE of bigeye and yellowfin tuna caught by Japanese longliner in the Indian Ocean standardized by GLM considering several aspects of area, catchability and data resolution. IOTC 2014/WPTT16/31. 23 pp.

Table 1. ANOVA table of GLM for year based area specific CPUE standardization.

R2

```

Response: log(CPUE + const)
      LR Chisq Df Pr(>Chisq)
Year    2797.1  44 < 2.2e-16 ***
Q        534.2   3 < 2.2e-16 ***
LatLon   3720.3  29 < 2.2e-16 ***
Cluster  27517.5  3 < 2.2e-16 ***
Vessel   10357.3 697 < 2.2e-16 ***
Year:Q    4441.0 117 < 2.2e-16 ***
---
```

R3

```

Response: log(CPUE + const)
      LR Chisq Df Pr(>Chisq)
Year    2239  45 < 2.2e-16 ***
Q        272   3 < 2.2e-16 ***
LatLon   9983  28 < 2.2e-16 ***
Cluster  69772  4 < 2.2e-16 ***
Vessel   12193 818 < 2.2e-16 ***
Year:Q    4077 134 < 2.2e-16 ***
---
```

R4

```

Response: log(CPUE + const)
      LR Chisq Df Pr(>Chisq)
Year    2571.4  45 < 2.2e-16 ***
Q        817.7   3 < 2.2e-16 ***
LatLon  15782.7  66 < 2.2e-16 ***
Cluster  25353.4  3 < 2.2e-16 ***
Vessel   17699.5 964 < 2.2e-16 ***
Year:Q    3009.6 135 < 2.2e-16 ***
---
```

R5

```

Response: log(CPUE + const)
      LR Chisq Df Pr(>Chisq)
Year    2858.1  45 < 2.2e-16 ***
Q        564.6   3 < 2.2e-16 ***
LatLon   1036.4  32 < 2.2e-16 ***
Cluster  28306.3  3 < 2.2e-16 ***
Vessel   9831.3 807 < 2.2e-16 ***
Year:Q    2152.2 132 < 2.2e-16 ***
---
```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1




Fig. 1. Definition of areas used in this study.




Fig. 2. Historical change in the proportion of fishing effort (number of hooks) in each area (Fig. 1).





Fig. 3. Species composition of catch in number in the Indian Ocean by the Japanese longline fishery in each area shown in Fig. 1.

R2



R3





Fig. 4. Beanplots for yellowfin region showing species composition by cluster for albacore (ALB), bigeye tuna (BET), yellowfin tuna (YFT), swordfish (SWO), southern bluefin tuna (SBT), sharks (SKX) and other fish (OTH). The horizontal bars indicate the medians.

R4



R5






Fig. 4. Beanplots for yellowfin region showing species composition by cluster for albacore (ALB), bigeye tuna (BET), yellowfin tuna (YFT), swordfish (SWO), southern bluefin tuna (SBT), sharks (SKX) and other fish (OTH). The horizontal bars indicate the medians. (continued)


R2



R3



R4



R5




Fig. 5. Annual change in species composition for albacore (ALB), bigeye tuna (BET), yellowfin tuna (YFT), swordfish (SWO), bluefin tuna (BFT), southern bluefin tuna (SBT), sharks (SKX) and other fish (OTH) by cluster and area.




Fig. 6. Standardized year based CPUE in number for each area.








Fig. 7. Standardized year based CPUE in number for each area with comparison of nominal CPUE and CPUE in the previous studies (Matsumoto et al., 2019 and Matsumoto, 2020).



R2



R3



R4



R5





Fig. 8. Standardized residuals of year based CPUE standardization for each of four areas expressed as histograms and QQ plots.

R2




Fig. 9. Influence plot for CPUE standardization for yellowfin.

R3




Fig. 9. Influence plot for CPUE standardization for yellowfin. (continued)

R4




Fig. 9. Influence plot for CPUE standardization for yellowfin. (continued)

R5




Fig. 9. Influence plot for CPUE standardization for yellowfin. (continued)