




Contents lists available at ScienceDirect

Marine Policy

journal homepage: www.elsevier.com/locate/marpol

Developing a science-based framework for the management of drifting Fish Aggregating Devices

Manuela Capello^{a,*}, Gorka Merino^b, Mariana Tolotti^a, Hilario Murua^c, Laurent Dagorn^a

^a MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, Sète, France

^b AZTI Marine Research, Basque Research and Technology Alliance (BRTA), Herrera Kaia Portualdea z/g, 20110 Pasaia, Gipuzkoa, Spain

^c ISSF, International Seafood Sustainability Foundation, Pittsburgh, PA, USA

ARTICLE INFO

Keywords:

Tropical tuna
Purse-seine fisheries
Indian ocean
Scientific advice
Indicators
Operating models

ABSTRACT

Fish Aggregating Devices (FADs) are man-made floating objects deployed by fishers to aggregate tuna and facilitate their catch. Currently, more than half of the global tropical tuna purse-seine catches occur at FADs. The fast development of the purse-seine fisheries operating on drifting FADs (DFADs) has raised concerns regarding their impacts on tuna populations, on non-target species like sharks, as well as on the pelagic and coastal habitats. Consequently, the management of DFAD fisheries is a priority for all tuna regional fisheries management organizations. Due to the little availability of science-based advice to support management decisions, resolutions on DFADs have been mainly based on precautionary principles. In this study we propose a science-based framework for the management of DFADs, relying on indicators and operating models. A set of models and indicators that help evaluate the ecological impacts of DFADs is presented, considering the case study of DFAD fisheries management in the Indian Ocean. The objective of this framework is to assess and predict the effects of DFADs on coastal and pelagic ecosystems, in order to support and/or evaluate past, present and future management actions.

1. Introduction

Tuna fisheries are among the world's most important fisheries in terms of global yields and economic value [1]. In 2020, the catch of major commercial tunas reached 4.9 million tonnes, of which around 95% correspond to tropical tuna catches [2], with skipjack tuna (*Katsuwonus pelamis*) exceeding 2.8 millions tonnes in 2020 and being the third top marine species in terms of total yield, only after the Peruvian anchoveta and Alaska pollock [1]. Tropical tuna species manifest an associative behavior with floating objects (FOBs), forming large multi-species aggregations around them [3]. Both artisanal and industrial fisheries worldwide take advantage of this associative behavior by deploying man-made floating objects, called Fish Aggregating Devices (FADs). Anchored FAD arrays (AFADs) are generally exploited by the artisanal/semi-industrial fisheries of coastal countries (e.g. trollers, handliners, small purse-seiners and pole-and-line vessels) [4–6], whereas drifting FADs (DFADs) are deployed offshore by the industrial purse seine fleets in all oceans [7]. In recent years the catches of tropical tuna at DFADs have peaked to more than 50% of the total tropical tuna catch [8,9,2,7]. The increasing DFAD-based fishing efficiency that

characterizes the industrial purse-seine tuna fisheries is linked to the development of novel technologies that equip DFADs, such as instrumented buoys with GPS beacons and, more recently, echosounders. These satellite-linked devices facilitate their location and provide remote estimates of the abundance of fish underneath the DFAD [10–14]. The use of DFADs has direct effects on:

- i. **Target species**, i.e. skipjack, yellowfin (*Thunnus albacares*) and bigeye tuna (*T. obesus*): by increasing their catchability and by shifting the purse-seine fishing effort on juveniles (for yellowfin and bigeye tuna) [8,9,15,16];
- ii. **Non-target (bycatch) species**: by increasing their catchability, with major concerns for some Endangered, Threatened and Protected (ETP) species, such as the silky shark (*Carcharhinus falciformis*) and the oceanic whitetip shark (*C. longimanus*) [17] but also through ghost fishing of DFADs when their design causes entanglements of sharks [18];
- iii. **Habitats**: the increased number of floating objects at sea leads to modifications of surface marine habitats, with unknown consequences on the fitness and physiological condition of tuna and

* Correspondence to: MARBEC, Avenue Jean Monnet, CS 30171, Sète 34203, France.

E-mail address: manuela.capello@ird.fr (M. Capello).

<https://doi.org/10.1016/j.marpol.2023.105657>

Received 3 January 2023; Received in revised form 18 April 2023; Accepted 18 April 2023

Available online 16 May 2023

0308-597X/© 2023 Elsevier Ltd. All rights reserved.

associated species [19,20], DFAD stranding on sensitive ecosystems (e.g. coral reefs) [21,22], and increased marine pollution when built up using non-biodegradable materials [23].

Accordingly, the use of DFADs has raised concerns about their sustainability [8,9,15,16,24] and has led all tuna regional fisheries management organizations (RFMOs) to establish limits to the number of instrumented buoys attached to DFADs (e.g.: Indian Ocean: IOTC Res.23/02 [25]; Atlantic ocean: ICCAT Rec 22-01 [26]; Eastern Pacific Ocean: IATTC C-21-04 [27]; Western and Central Pacific Ocean: WCPFC CMM 2021-01 [28]). Other management measures have also been adopted for purse seiners by tuna RFMOs to reduce the impact of DFAD fishing, such as time-area closures (e.g., ICCAT Rec 21-01; WCPFC CMM 2021-01) and discard bans (e.g., IOTC Res.19/05). Furthermore, specific regulations have imposed the use of fully non-entangling DFADs made without netting material (IOTC Res. 23/02 [25] or non-entanglement risk DFADs (ICCAT Rec.19-02 [29], IATTC Res. C-19-01 [30], and WCPFC CMM 2021-01 [28]) to reduce shark mortality and avoid the entanglement of other marine megafauna (e.g. sea turtles). In parallel to these DFAD management measures, a series of mitigation measures have been voluntarily adopted by certain purse-seine fleets to reduce the impacts of the use of DFADs, including changes in the design of DFADs to reduce shark entanglements [31], the transition towards biodegradable materials [31,32], the development of DFAD recovery programs [33] and the implementation of best practices for releasing sharks and other bycatch species [34-36].

Options to reduce the impacts of DFAD fishing are regularly discussed within dedicated RFMOs technical FAD working groups, Scientific Committees and RFMOs Commission meetings, where ultimately management measures are adopted [37]. However, due to the lack of quantitative science-based advice for DFADs management and empirical evidences, it is difficult to ascertain the effectiveness of those adopted management measures. Similarly, the mitigation measures adopted by the fishing industries to reduce the impacts of DFADs are often questioned, because either the measures are not considered efficient enough or due to their lack of enforcement [38-40]. More globally, the participation of increasingly diverse stakeholders in RFMO technical scientific working groups, where scientific advice for fisheries management is developed, often leads to deviate discussions from a scientific perspective towards an advocacy and political angle. Therefore, it is paramount to clarify the role of the scientific advice in the development of DFAD

management measures.

This study presents a framework for a science-based management of DFAD fisheries, relying on operating models and indicators to support the development and implementation of management measures within tuna RFMOs. The main focus of this study are DFADs, although the same framework can also be applied to the management of AFAD fisheries.

2. Definition of a science-based DFAD management framework

The conceptual framework of a science-based DFAD management scheme proposed in this study consists in a feedback-loop process, going back and forth from stakeholders to policymakers (Fig. 1). Similar to the process used in the Management Strategy Evaluation (MSE), the global aim of this scheme is to identify a management strategy that will allow achieving the adopted management objectives [41]. The role of scientists lies at the heart of this loop and aims to provide ecological knowledge and advice to: (i) support the formulation of novel management and mitigation measures, depending on the management objectives (ii) provide feedback information on the effectiveness of past management measures that may cause the management decisions/objectives to be revised.

This conceptual framework is headed by the definition of *management objectives* and Target Reference Points (TRPs) (Fig. 1). Defining and prioritizing clear management objectives is an essential step for fisheries management [42]. Currently, management objectives driving the decision-making process, such as controlling the fishing capacity or reducing ecosystem impacts of DFADs, have either not been set, or not been prioritized [39]. The same holds for the definition of specific TRPs, i.e., target indicators' levels related to DFAD fisheries to be achieved through management measures. The definition of management objectives and TRPs should be agreed at Commission level taking into account different considerations (e.g., socioeconomic impacts, sustainability, etc.) raised by all stakeholders. For example, a clear management objective that has been recently identified by tuna RFMOs is reducing the risk of entanglement of sharks at DFADs, for which specific management measures have already been adopted. However, the effectiveness of shark entanglement mitigation measures at DFADs is still largely debated when fully non entangling DFADs are not used. In this respect, a TRP related to shark entanglement risks, such as reducing them of a given percentage with respect to a reference year, has not been defined yet.




Fig. 1. Schematic view of a science-based approach for FAD management relying on indicators and operating models.

A clear definition of objectives and TRPs can guide the provision of scientific advice, the science underpinning management decisions and the adoption of management measures. Once management objectives and TRPs are agreed, scientists can develop a set of indicators and performance metrics to quantitatively evaluate the management options against the agreed management objectives and TRPs. For example, if the objective is to reduce shark entanglement, it is necessary to monitor individual shark entanglement rates and estimate which levels would be acceptable from a management perspective (see [18] and Section 2.1 for further details). Performance metrics would consist in analyzing how far are these indicators from TRPs. Furthermore, other indicators than those specifically addressing the management objectives themselves can be produced by scientists, in order to evaluate additional impacts of management decisions. Indeed, management measures aimed at reducing fishing mortality on one stock can induce a change in fishing strategies and produce unintended impacts on target species, non-target species and habitats. For example, a recent study [43] demonstrated that the rebuilding plan for yellowfin tuna adopted by the Indian Ocean Tuna Commission (IOTC Res. 16/01) was followed by an increase on the number of DFAD sets and fishing effort, resulting in higher by-catch of silky sharks (see more details in Section 2.1). In summary, within the proposed science-based FAD management framework, the direct and indirect impacts of management decisions would be assessed using multiple indicators.

Indicators alone lack predictive capabilities to test candidate management options. Therefore, another important building block of the proposed science-based framework, requires the development of *operating models* (OM). Numerical models constitute an essential, complementary tool to support decision-making. In fisheries science, OMs simulate the past and future dynamics of the fish stocks and the fisheries to evaluate the consequences of different management procedures (MP, also referred to as harvest strategies) on exploited fish populations. The use of OMs is widespread in fisheries management within MSE frameworks [44,41,45], where OMs can be used to test different FAD management strategies and the resultant performance metrics for each management option will estimate the probability of achieving management objectives and TRPs. To date, most applications of MSE have focused on evaluating MPs that define input and output controls (e.g., effort and catch limits). As a consequence, within tuna RFMOs, the OMs considered in the MSE frameworks developed so far account for the population dynamics of tuna species and their associated fisheries [41]. From a more general perspective, the main scope (and challenge) of an OM is to describe the main population and fisheries processes that are relevant to fisheries management. In the case of DFAD fisheries management, OMs certainly need to account for the *stocks dynamics* of target species, which allow understanding at which level the catches at DFADs, are sustainable (subsection 2.2.1). However, other dimensions need to be considered too. Namely, OMs accounting for the *fish and fleets behavior* can predict the trends of DFAD catches of tuna and non-target species as well as their sustainability according to different management measures and DFAD densities (subsection 2.2.2). Second, accounting for the spatio-temporal dynamics of the DFADs themselves using OM that simulate *FOB drifts* is key to predict DFAD stranding and loss events, as well as changes in surface habitats (number of FOBs) according to different management strategies (subsection 2.2.3). Within such OM-based approach, the uncertainty related to the biology and behavior of target and non-target species can be characterized, providing a quantitative framework for the application of the precautionary approach [46].

In summary, indicators and performance metrics will allow evaluating how well the different management options tested using the OMs perform, to achieve the agreed management objectives and TRPs. The development of OMs, as well as testing/evaluating candidate/past MPs through indicators and performance metrics, is carried out by scientists in technical scientific working groups. The performance of those candidate management options is then presented and discussed in joint

dialogue meetings among scientists, stakeholders and managers. Ultimately, depending on the management objectives, managers will adopt a set of “best” FAD **management measures**, which are tailored to achieve the agreed objectives, during tuna RFMO Commission meetings. These management measures can also be complemented with **technical mitigation measures** developed by the fishery itself on a voluntary basis [31,34-36,32]. The key aspect of a science-based management relies on the fact that the selection of management/mitigation measures, as well as the evaluation of their effectiveness and global impacts on marine ecosystems, can draw on indicators and OMs’ outputs provided by scientists separating clearly the role of science and managers.

In the following sections specific indicators and OMs that can help evaluating the ecological impacts of DFADs are presented. The effect of past management decisions (e.g., total limits on the number of operational buoys, but also management measures that are not strictly related to DFADs, like the introduction of quotas) is discussed, by evaluating the trends of the indicators that can be built from the available information, considering the Indian Ocean as a case study.

2.1. Indicators

In order to quantitatively evaluate the ecological impacts of DFADs and support management decisions, a number of indicators can be set up considering (i) target species, (ii) non-target species and (iii) coastal and pelagic habitats (Table 1). In the following sections, for each set of indicators, an example describing the data requirements/availability and a rationale of their relevance for DFAD management is provided.

2.1.1. Target tuna species

Tuna catches (by species) harvested at FOBs (i.e., any natural and artificial floating objects) by purse seiners can be obtained from logbook data and are generally available within all tuna RFMOs for stock assessments and other scientific purposes. Generally, the logbook catches of tuna by species are corrected considering port sampling data [47], in order to account for possible bias related to the difficulty to identify species for logbook declarations, and/or observer sampling [48]. This correction is conducted separately for FOB-associated schools and free-swimming schools (not associated), because their corresponding species and size composition differ. The timeline of total catches of FOB-associated tuna can provide useful insights on the effects of past DFAD management measures on the fishery. Of course, discussing catch trends on the light of past management measures alone should be taken with precaution, because the tuna catches depend both on the tuna abundance, their catchability and the fishing effort as well as catch/effort of other gears. However, catch trends still remain a straightforward indicator that can inform on the evolution of the fishery through time.

For example, in the case of the Indian Ocean, the timeline of total catches of FOB-associated tuna obtained from the main purse-seine fleets that exploit DFADs (Fig. 2) demonstrates increasing trends for skipjack tuna until 2018, with a clear decrease in 2019 and 2020. The catches of yellowfin and bigeye tuna followed similar trajectories but increased to a minor extent in the period prior to 2018. Despite the difficulty to disentangle the reasons of the observed trends in relation to the above-mentioned factors (changes in tuna abundance, catchability and fishing effort), the first DFAD management plans adopted in the IOTC (Res. 15/08; Res. 17/08) which limited the number of instrumented buoys that could be used by the purse seine fleets, did not seem to alter the global increasing trends of FOB-associated catches. The recent DFAD management plan (Res. 19/02), that entered into force in January 2020, occurred after a significant decrease in tuna catches, which could be explained by the entry into force of the new rebuilding plan for yellowfin tuna adopted in October 2018 (Res.18/01).

A second indicator related to target tuna species concerns their physiological conditions. The so-called “ecological trap hypothesis” considers that FADs can alter tuna movements and retain them in less

Table 1
Possible ecological indicators to monitor the impacts of DFADs on target tuna species, non-target species and coastal and pelagic habitats.

Category	Ecological Indicators	Data source	
Target species	Species-specific catches at DFADs	Logbook data Observers' data	
	Species-specific physiological condition	Scientific cruises	
Non-target species	Species-specific catches at DFADs	Logbook data Observers' data	
	Species-specific physiological condition	Scientific cruises	
	Entanglement rates at DFADs	Scientific cruises	
Habitats	Number of DFAD stranding events	Buoys data Logbook data	
	Amount of plastics stranded/sank	Buoys data Logbook data Observers' data In-situ observations	
		Total number of DFADs relative to the number of natural floating objects	Observers' data




Fig. 2. Timeline of purse seine catches of FOB-associated tuna in the Indian ocean (YFT: yellowfin tuna; BET: bigeye tuna; SKJ: Skipjack tuna; TOTAL: sum of the catches of the three species). Only the tuna catches for the main purse-seine fleets exploiting DFADs have been considered: EU-Spain, EU-France, Seychelles, Mauritius and Korea. Resolutions related to DFAD management plans are indicated with black vertical dotted lines (AB= limit for the number of active buoys per vessel; BY= limit in the total number of buoys purchased yearly per vessel). Other relevant resolutions affecting the DFAD fisheries are indicated in red. All resolutions are indicated considering their date of entry into force in the abscissa.

Data Source: Indian Ocean Tuna Commission (<https://www.iotc.org/WPTT/23AS/Data/05-CESurface>).

productive regions, with potential negative consequences for their fitness [20]. In order to evaluate to which extent the physiological condition of tuna has been affected by the increase on the number of DFADs deployments, it is necessary to elucidate the link between the

number of FADs and tuna biological condition (e.g. development of gonads, thorax girth, size etc.). So far, only ad-hoc studies, focusing on restricted zones and time periods, have tried to address this issue in different oceans [49-53]. Regular data collection should be conducted to

provide temporal and spatial trends of tuna biological and physiological indicators, considering the evolution of the number of DFADs as a co-variable. The role of DFAD density on tuna condition relative to other environmental variables (e.g., sea-surface temperature and chlorophyll) could also be evaluated, since these variables can also change through time and affect tuna physiology [54].

2.1.2. Non-target species

Indicators on non-target species should primarily focus on ETPs of major concern such as the silky shark, which constitute the main shark bycatch species caught at DFADs and is listed as Vulnerable by the International Union of Conservation of Nature (IUCN), and the oceanic whitetip shark, considered as Critically Endangered according to the IUCN [55-58,17,59]. A straightforward way to evaluate the effectiveness of any management measures that aim to reduce shark bycatch consists in developing a timeline of the overall number of sharks caught at DFADs. The information on the catch per set of sharks originates from observers' data. Scientific observers on-board purse seiners report the number of sharks caught for each observed set. The data is gathered by tuna RFMOs and can be made available to scientists, in aggregated forms. Extrapolation factors are generally applied in the case of partial observer's coverage, considering the shark catch per set of the observed DFAD sets and the overall number of purse seine sets operated on DFADs in the same period/region. Because sharks are often released by purse-seiners after the catch, trends on the overall number of sharks caught at DFADs should be further refined to account for actual mortality. This could be done by considering at-vessel and post-release mortality rates from dedicated studies that also consider release practices. A few independent studies estimated mortality rates for silky sharks caught by purse seine vessels in the Pacific, Atlantic and Indian oceans [60,61,35,62]. These studies estimate total mortality rates between 60% and 80%, even when individuals are released following good practices. This is mainly due to the high at-vessel mortality rate, as most silky sharks are already dead by the time they reach the deck. The best way to incorporate the results of the different studies into total mortality estimates needs to be discussed, since they vary depending on the implementation of good release practices, that could be vessel or fleet specific. Because these practices can also evolve with time, such dedicated post-release survivorship studies should be periodically updated.

In the Indian ocean, catches of silky sharks (Fig. 3 and Supplementary Material S1) attained a maximum in 2016 and remained relatively

stable in the subsequent years. However, this indicator's trend should be considered with care. Indeed, information on the total number of DFAD sets was not readily available from IOTC Form 3FA and one issue with using national reports (see details in Supplementary Material S1) is that the information on the number DFAD sets is not georeferenced. Therefore, total silky shark catches can only be derived globally and not by area as is in Tolotti et al. [43]. Furthermore, further improvements of this indicator should include at-vessel and post-release mortality rate.

In addition to the indicator shown in Fig. 3, a dedicated indicator providing the temporal evolution of shark entanglement risks should be produced to evaluate this additional source of mortality. A first study [18], quantified the extent of shark entanglement within the underwater structure of DFADs in the Indian Ocean. Since this study unveiled this issue, mitigation measures such as changes in the design of DFADs were voluntarily adopted by the purse seine fleets to reduce the risk of entanglement [63]. Resolutions for the adoption of fully non-entangling DFADs without netting [25] and with low entanglement risk have also been adopted across RFMOs [29],[30],[28] with the objective of reducing the risk of shark entanglement. The assessment of shark entanglement at DFADs relies on pop-up satellite electronic tags and scientific diving data that inform respectively on the vertical behavior of silky sharks (by which entanglement events can be detected) and on the number of sharks observed entangled in the DFADs' underwater structure [18,61,35,36]. Observers can report entanglements when vessels lift DFADs out of the water, but decomposed carcasses may detach from the DFAD before emerging from the water. To the purpose of monitoring the temporal evolution of shark entanglement risks and the effectiveness of management measures, regular scientific cruises should be conducted. Electronic tagging of sharks could also be performed by trained observers on board purse-seiners.

Finally, similar to tuna, the physiology of other associated species like sharks can also be affected by increasing DFAD densities. Recent electronic tagging studies conducted on both tuna and non-tuna species found around DFADs demonstrated that silky sharks spend significant amounts of time associated with FADs [64] with a similar associative dynamics as tunas (i.e., comparable residence times) [65]. Therefore, if the "ecological trap" scenario is proven, it could equally be applied for these species. Future data collection and research effort should be dedicated to evaluating shark physiological condition and their trends relative to changes in the DFAD density. No data is currently available to build this indicator in the Indian Ocean.




Fig. 3. Estimated purse seine catches of silky sharks at DFADs (number of individuals) by the main fleets operating in the western Indian Ocean (Spain, France and Seychelles). Resolutions related to FAD management plans are indicated with black vertical dotted lines (AB= limit for the number of active buoys per vessel; BY= limit in the total number of buoys purchased yearly per vessel). Other relevant resolutions affecting the DFAD fisheries are indicated in red. All resolutions are indicated considering their date of entry into force in abscissa.

2.1.3. Habitats

Indicators on the impacts of DFADs on coastal and pelagic habitats aim at assessing how management actions affect the number of DFAD stranding events, the amount of marine litter, pollution and habitat damages caused by DFADs, as well as changes in the number of floating objects found at the sea. The main data sources to build such indicators consist in GPS data transmitted by the satellite-linked buoys which equip all DFADs [66,21,67,22] as well as observers' and logbook data [19]. Furthermore, in-situ observations can provide useful information on materials and size of the DFADs, as well as their environmental impacts (e.g. coral reef damage) [68].

Stranding events correspond to DFADs stranded in coastal environments. Their impacts are particularly critical for coral reef areas, since the DFAD structure can damage these sensitive habitats [40]. The percentage of DFAD stranding events can be assessed from satellite-linked buoys data [66];[21,22]. However, in order to evaluate the ecological impacts of DFADs on marine habitats, assessing the trends in the *total number of DFAD stranding events* is necessary. Namely, a stable percentage of DFAD stranding events can still imply higher/lower numbers of stranded DFADs (and therefore higher/lower ecological impacts) if the number of DFAD deployments increases/decreases through time. To this purpose, disposing of reliable data on the number of DFAD deployments is key. In the Indian ocean, using satellite-linked buoys data provided by the French purse seine fleet, Imzilen et al. [21] highlighted a steady increase in the percentage of DFAD stranding events (from 3.5% in 2008 to nearly 20% in 2013), followed by a stabilization (between 15% and 20% in the period 2013–2017). On the other hand, so far no study has been conducted for other purse seine fleets operating in the Indian Ocean, to evaluate whether similar trends have occurred over the same period. Furthermore, the IOTC Secretariat recommends caution when analyzing DFAD deployment data provided through Form 3FA [69]. Given these uncertainties and data limitations on the total number of DFADs deployed available at the IOTC level, to date, a reliable timeline of the number of DFAD stranding events cannot be built for the Indian ocean.

DFAD stranding events are not the only possible fate of DFADs. Indeed, DFADs can also sink in the open ocean. Secondly, when a DFAD drifts out the fishing zones, its echosounder buoy can be deactivated by the vessels. The deactivation of a buoy implies that its position and echosounder data is not received any more by its end users. Consequently, the DFADs whose buoys are deactivated are considered lost or abandoned by the vessel. This practice can be encouraged by resolutions setting limits on the number of operational buoys (i.e., buoys actively transmitting information to their end users): in order to comply with the authorized limits, purse-seiners can deactivate buoys that depart from their fishing grounds. Despite this practice can be restricted by the limits in the annual purchase of instrumented buoys and the total number of buoys in stock per vessel (e.g., Resolution 23/02 in the case of the Indian Ocean), the actual effect of past and current management measures on this practice remain to be assessed. Finally, the retrieval of the echosounder buoy by another fishing vessel, a common practice in all oceans, may be another issue of DFAD loss if it is not replaced by another buoy. More generally, this practice makes the DFAD traceability more complex, since a single DFAD can be associated to more than one instrumented buoy during its lifetime. Indicators accounting for the number of DFAD stranding/sinking events should include these additional and unknown sources of DFAD losses. Because DFADs also include non-biodegradable materials for the raft frames, floats, and subsurface structure, all these DFAD-loss events can be a source of marine pollution and should be monitored [70]. In this respect, the amount of plastics lost into the oceans should be quantified through a specific indicator accounting for the DFAD design (weight and composition of their constituent materials) [71]. This indicator would also allow quantifying the effectiveness of management and mitigation measures promoting the shift towards biodegradable materials for some of the DFAD components [23,70], as well as designing adequate DFAD recovery programs [67].

Finally, changes in the surface habitats induced by the deployment of DFADs can be monitored considering the temporal evolution of the number of DFADs relative to the number of natural floating objects (NLOGs), which constitute a natural component of the pelagic habitat [8,9,19]. In the western Indian Ocean, such indicator can be built from observers' data, which report the position and type of floating objects encountered by the fishing vessels. A recent study suggests that the ratios between the number of DFADs and the number of NLOGs have increased in recent years (2014–2018) relative to 2007–2008. The entire western Indian Ocean has been affected, with DFADs representing more than 85% of the overall FOBs for the period 2014–2018, NLOGs less than 10%, and other floating objects originating from human-induced pollution corresponding to 5% (e.g., land-based pollution, as well as pollution resulting from lost fishing equipment and gears other than FADs) [19].

2.2. Operating models

An ensemble of operating models can be developed to support management decisions aiming at mitigating the ecological impacts of DFADs (Table 2).

2.2.1. Stocks dynamics OM

The issue of the increased catch of juveniles of yellowfin and bigeye tuna on floating objects and the relative reduction of sets on free-swimming schools targeting adult individuals is a major concern of the DFAD fisheries [15,16,72]. Understanding the consequences of this shift in the exploitation of the juvenile component of yellowfin and bigeye stocks on the yield-per-recruits is key to ensure the sustainability of the DFAD fisheries. However, considering the DFAD catches alone may not be sufficient to draw conclusions: it is necessary to account for the whole stocks and fisheries dynamics, including the catches of juveniles by other gears and adults from other fisheries (e.g., longline fisheries). To this purpose, OMs similar to those currently used in stock assessment and in the MSE framework [45], should be used to quantify the impacts of increased fishing mortalities of juvenile yellowfin and bigeye tunas on the yield-per-recruit and stock biomass in conjunction with the impact on fishing in the adult component. These models can be run considering the main sources of uncertainty on tropical tuna stocks' dynamics. Simulations can be conducted considering different fishing-mortalities-at-age to identify the best total-allowable-catch-at-age options for the stock rebuilding plans. However, this implies catch allocation rules between gears which are beyond the scientific discussion and involves managers' decisions. Fishery impact analysis [73] can be done by analyzing how the population spawning potential has been impacted, historically and at present, by major fishery types over years. Fishery impact analysis is done by estimating the spawning biomass dynamics over time that would have occurred in the absence of historical fishing. The reduction in spawning biomass potential induced by a particular fishing gear is estimated, so

Table 2

List of operating models focusing on target tuna species, non-target species and coastal and pelagic habitats.

Category	Operating model	Output/Prediction
Target species	Stocks dynamics	Optimal catch and catch-at-age options for maintaining the stock on the target reference points and rebuilding tuna stocks Fishery impact plots per fishing gear Catch trends for variable DFAD densities
Target/ non-target species	Fish and fleets behavior	Catch trends for variable DFAD densities
Habitats	FOB drifts	Number of DFAD stranding and sinking events Changes in the density of floating objects (DFADs and NLOG inside and outside the fishing grounds)

that the relative fishery impact on Spawning Stock Biomass (SSB) by major gear type can be compared (see figure 46 in [73]). As such, fishery impact plots could inform managers on the relative contribution of each gear to the stock status. In the Indian ocean, this topic is particularly important in relation to the rebuilding plans of overfished stocks, such as the yellowfin tuna (IOTC Res. 16/01; Res. 18/01; Res. 19/01; Res. 21/01).

2.2.2. Fish and fleets behavior OM

Another ensemble of OMs is necessary to assess how increasing DFAD densities affect the catches of tuna and associated species, including ETP species such as silky sharks. Previous studies already demonstrated that increasing DFAD densities do not necessarily imply larger associated populations [74] or a higher number of DFAD sets [75], revealing that the relationship between number of DFADs and tuna catches can be non-linear and/or non-monotonic. In order to provide scientific advice on DFAD MPs, operating models that can predict tuna and ETP species catch trends for variable DFAD densities should be developed. These models should account for both the associative dynamics of tuna/non-target species at DFADs and the purse seine fishing practices at DFADs (number of DFAD sets). Indeed, for a given DFAD density, the amount of catches of each species depends on both their associative dynamics (i.e. the proportion of time spent associated with DFADs and, hence, the proportion of the population which is vulnerable to the fishery [76]) and fishers' behavior (which affects the number of DFAD sets) [75]. Building and conditioning such models requires combining information from multiple data sources: logbook/observers' data (which inform on catches per set and number of DFAD sets), echosounder buoys data (which inform on the presence/absence of tuna at DFADs as well as on DFAD densities), and on electronic tagging data (which inform on the amount of time that tagged individuals spent at and away from DFADs), see [Supplementary Material S2 \(Tables S1 and S2\)](#). Recently, in the Indian ocean, progress in the developments of such OM have been made [77]. The model of Dupaux et al. accounts for tuna associative behavior obtained from electronic tagging data, allowing to assess changes in the time spent by tuna between two associations as a function of the DFAD density. Combined with information on DFAD densities obtained from buoys data provided by the IOTC secretariat (Form 3BU), this model has been used to quantify for the first time the proportion of FOB-associated tuna population within the fishing grounds exploited by the purse-seine fisheries in the Western Indian Ocean [77]. Further developments of this model, coupling tuna behavior with models of fleets behavior accounting for the number of DFAD sets as a function of the DFAD densities, will allow predicting the evolution of FOB-associated tuna catches, and fishing mortalities, for different DFAD densities. Uncertainties related to the behavioral processes setting the aggregation dynamics (e.g., the role of schooling behavior [78]) could also be accounted for in this OM.

2.2.3. FOB drifts OM

Models of FOB drifts, capable of predicting the trajectories of FOBs from their release location and the local surface currents can be used to assess the risks of DFAD stranding [66,79] and changes in the surface habitats [19]. These models, building on previous results showing that FOBs drift similarly to oceanographic drifters [80] simulate FOB trajectories using Lagrangian simulations [81]. Model inputs include (i) FOBs numbers and initial positions (ii) ocean surface currents (iii) the average lifetime of FOBs. To account for uncertainties in the modeled FOB trajectories, a random walk component of particles motion can be considered within the Lagrangian model [79]. However, the models could be further validated when using observed FOB trajectories that are being submitted to various tuna RFMOs (e.g., IATTC C-21-04, IOTC Res. 23/02). FOB lifetimes also constitute a source of uncertainty and FOB drift OMs run considering different lifetimes allow to assess the sensitivity of results [19].

3. Discussion

This study proposes a science-based framework, based on an ensemble of OMs and indicators, to support the development of FAD-fisheries management plans. We show ways to evaluate the effectiveness of DFAD management measures by providing specific examples of indicators related to the ecological impacts of DFADs and their trends, considering the Indian Ocean as a case study.

Some of the indicators proposed in this study are available in the literature and have been presented to tuna RFMOs [8,9,38,19,82,21,75,83]. However, scientific studies generally provide only a snapshot over a given time window where the studies were conducted or consist of one-time studies [18]. Here, a continuous monitoring of such indicators is proposed, in order to provide scientific advice and adapt management if necessary. Depending on the management objectives, some indicators could be spatialized and considered on a quarterly basis to allow for spatio-temporal management decisions. Derived indicators, that are drawn from the former, can also be built (for example, the ratio between the DFAD catches of target and non-target species), depending on the management objectives.

Other indicators which include data from other fishing gears and techniques should complement those currently proposed: indicators showing the target and non-target relative catch of DFADs compared to other gears, or, for the purse seine fisheries only, indicators comparing the catches conducted at DFADs and free-swimming schools. Global catches of non-target species and discard rates could also be considered. Furthermore, data collection of in-situ observations of stranded DFADs [68] will allow building new indicators on environmental impacts of DFADs (e.g., coral reef damaged) on coastal habitats. Economic indicators would also be key to weight the interests of all parties and evaluate the impacts of management decisions beyond the ecological aspects. For example, the implications of total retention of non-target species in coastal countries, where long-distance fisheries ports are based, could be measured by monitoring fish prices and sales in local markets, or indicators related to the creation of employment due to canneries. Moreover, indicators accounting for cultural, political and social implications of DFAD use could be defined. Finally, indicators monitoring the purse seine carbon emissions evolves with time and are affected by management decisions could also be used [84] and compared with other types of gears.

Due to the complexity of the relationship between number of DFADs, tuna abundance, tuna associative dynamics, the fishing strategies adopted by purse seiners, and the catches of target and non-target species, it is difficult to predict the effects of management measures from indicators alone. For this reason, the use of OMs should complement the proposed indicators. In the current framework, an ensemble of operating models is proposed, each aiming to provide science-based advice on a specific ecological impact of DFADs on target species, non-target species and habitats. So far, OMs developed within the MSE approach have been used to support management decisions for single species stocks management [85]. Remarkably, although the approach discussed in this paper follows the same spirit as the MSE, it substantially differs regarding its targets and objectives, since it is devoted to the management of a fishing tool rather than a fish stock. In conjunction with the management of other fishing tools/gears, this approach will contribute to the sustainability of tuna fisheries and, more globally, the good health of marine ecosystems.

The scientific advice that can be produced through the proposed indicators and operating models is within the scope of the ecosystem-based for fisheries management (EBFM). Several studies advocated the adoption of an integrated EBFM in RFMOs [86-92]. However, so far very little applications of an EBFM can be found within tuna RFMOs, where management decisions are most often taken considering single species management approaches. In the case of DFAD fisheries, similar to other fishing practices and gears, whose impacts involve not only target tuna species, but also non-target species, coastal and pelagic habitats,

adopting an integrated EBFM and overcoming the limits of single-species management approaches is an essential step. In this respect, this study offers new pathways towards the implementation of an EBFM. Similarly, the use of OMs allows accounting for uncertainties on biological and behavioral processes, which is a pre-requisite of the Precautionary Approach framework [46].

Both indicators and OMs rely on catch-dependent and independent data provided by fisheries monitoring programs, scientific surveys and stakeholders such as fishers. The synergy between data collection programs and the development of models and indicators is a key aspect of the proposed framework. In this respect the number of deployments and density of DFADs are undoubtedly a key input for building and interpreting the proposed indicators for monitoring the ecological impacts of DFADs. Until recently, instrumented buoys data (buoy daily position data and/or echosounder raw acoustic biomass data) were provided to national scientists through specific agreements which concerned only national fleets. These data have certainly proven to be useful for scientists [10,11,13,93], but their partial coverage has so far hindered accurate estimations of total DFAD densities. Nowadays, specific resolutions have been adopted to request all contracting parties using DFADs to provide buoys position data to tuna RFMOs (e.g. IOTC Res 23/02) and buoy daily position data and biomass data (IATTC CMM 21–04). These new datasets allow estimating the density of operational buoys (i.e., the buoys which are transmitting their data remotely), which can be used as a proxy of the DFAD density. However, as mentioned before, buoys can currently be deactivated or retrieved while at sea. As a result, the density of operational buoys can only be considered as a lower bound for the actual number of DFADs floating in the ocean. More importantly, depending on the buoy deactivation practices (which may change with time depending on the limits imposed on the maximum number of operational buoys allowed by tuna RFMOs) the gap between the number of operational buoys and DFADs could vary, further biasing the estimates of the total number of DFADs at sea estimated based on operational buoy numbers. To cope with this issue, buoys data should be complemented with additional information allowing a higher traceability of the fate of DFADs. First, the information on the number of deactivated buoys should be made available at the same scale as the operational buoys' position data. Second, the number of DFAD deployments and retrievals should accurately be provided to tuna RFMOs and scientists. Moreover, DFAD should be associated with unique identifiers and their encounters (since the time of deployment) should be recorded within dedicated DFAD logbooks.

Since January 2020, in the Indian ocean, due to the entry into force of IOTC resolution 19/02, the buoys GPS positions are provided to the IOTC Secretariat at a daily scale by the member countries, to support the monitoring of compliance with the limitation established by the same resolution. The IOTC secretariat can currently deliver this data in an aggregated form, at a 1°/monthly scale. This novel data availability constitutes a big step forward, because the information on the overall density of echosounder buoys (and not only the buoys of a single purse-seine fleets delivered through specific agreements with national scientists) is key to evaluate DFAD densities and their ecological impacts. However, the unavailability of fine-scale buoy data (i.e., daily GPS positions of the buoys) still limits its exploitation for scientific purposes. Indeed, the operational information exploited by skippers has much higher resolutions with data being transmitted every few hours (e.g., 6–12 h in default mode, depending on the buoy model). Similarly, both tuna and shark species respond to the local number of FADs at time scales of the order of few days [65]. In order to build and condition operating models accounting for the behavior and catchability trends of tuna and associated species for variable FAD densities, making these data available to scientists at a finer scale will be necessary.

Finally, the data related to FOB-associated catches of tuna species provided by tuna RFMOs generally include the aggregated tuna catches conducted both on NLOGs, DFADs or other floating objects. Management decisions promoted by tuna RFMOs can set limits in the number of

DFADs but cannot apply to other types of floating objects such as the NLOGs. However, the latter can also show variable densities through time, due to natural and anthropogenic factors, such as extreme weather events or climate change. Reporting separately the tuna catches conducted on NLOGs and DFADs is key to discern the specific effects of management decisions devoted to set DFADs limits from other sources of variability in the total number of FOBs that are independent from management.

4. Conclusion

This study presents a new science-based framework to evaluate options for the management of DFADs built upon operating models and indicators. The operating models can support scientists and decision-makers alongside a set of indicators obtained from both fisheries-dependent and independent FAD-related data, towards a science-based approach for the management of FAD fisheries. The operating models and indicators proposed in this study can also be considered as part of adaptive management strategies. In this respect models and indicators can be used to build performance measures and to understand the effectiveness of past, present and future management actions. The same approach can be applied to other fisheries, including the science-based management of AFAD fisheries by coastal states. The use of DFADs is currently the source of conflicting debates. Recent discussions on FADs in tuna RFMOs have gone through at the point of turning scientific and technical groups into political debates. As a result, scientific evidence and advice have lost its importance to provide management advice and mainly the political position of interest groups has prevailed in the discussions and scientific groups. With the approach suggested here, science can be put back at the forefront of the subject to provide the DFAD fishery management advice.

CRedit authorship contribution statement

Manuela Capello: Conceptualization, Writing – original draft. **Mariana Tolotti:** Silky shark Data curation and analysis, paper Writing – review & editing. **Hilario Murua:** Paper Writing – review & editing. **Gorka Merino:** Paper Writing – review & editing. **Laurent Dagorn:** paper Writing – review & editing.

Data Availability

Data will be made available on request.

Acknowledgements

This study was supported by the MANFAD project funded by *France Filière Pêche*.

Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at [doi:10.1016/j.marpol.2023.105657](https://doi.org/10.1016/j.marpol.2023.105657).

References

- [1] FAO, The State of World Fisheries and Aquaculture 2022. Towards Blue Transformation, FAO, Rome, 2022, <https://doi.org/10.4060/cc0461en>.
- [2] ISSF. (2022). Status of the World Fisheries for Tuna: March 2022. ISSF Technical Report 2022–04. International Seafood Sustainability Foundation, Washington, D. C., USA., *March*.
- [3] P. Freon, L. Dagorn, Review of fish associative behaviour: toward a generalisation of the meeting point hypothesis, *Rev. Fish. Biol. Fish.* 10 (2000) 183–207, <https://doi.org/10.1023/A:1016666108540>.
- [4] A.R. Jauharee, M. Capello, M. Simier, F. Forget, M.S. Adam, L. Dagorn, Tuna behaviour at anchored FADs inferred from Local Ecological Knowledge (LEK) of pole-and-line tuna fishers in the Maldives, *PLoS ONE* 16 (2021) 1–19, <https://doi.org/10.1371/journal.pone.0254617>.

- [5] E.D. Macusi, N.A.S. Abreo, R.P. Babaran, Local ecological knowledge (LEK) on fish behavior around anchored FADs: the case of tuna purse seine and ringnet fishers from Southern Philippines, *Front. Mar. Sci.* 4 (JUN) (2017) 1–13, <https://doi.org/10.3389/fmars.2017.00188>.
- [6] H. Sadusky, P. Chaibongsai, D.J. Die, M. Shivilani, Management of moored fish aggregation devices (FADs) in the Caribbean, *Collect. Vol. Sci. Pap. ICCAT* 74 (5) (2018) 2230–2242.
- [7] Scott J. and Lopez, J. (2014). The use of FADs in tuna fisheries. Available at: [http://www.europarl.europa.eu/regdata/etudes/Note/Join/2014/514002/IPOL-PECH_NT\(2014\)514002_EN.Pdf](http://www.europarl.europa.eu/regdata/etudes/Note/Join/2014/514002/IPOL-PECH_NT(2014)514002_EN.Pdf) (Last Accessed: October 6, 2020).
- [8] L. Dagorn, N. Bez, T. Fauvel, E. Walker, How much do fish aggregating devices (FADs) modify the floating object environment in the ocean, *Fish. Oceanogr.* 22 (3) (2013) 147–153, <https://doi.org/10.1111/fog.12014>.
- [9] L. Dagorn, K.N. Holland, V. Restrepo, G. Moreno, Is it good or bad to fish with FADs? What are the real impacts of the use of drifting FADs on pelagic marine ecosystems? *Fish. Res.* 14 (3) (2013) 391–415, <https://doi.org/10.1111/j.1467-2979.2012.00478.x>.
- [10] Y. Baidai, L. Dagorn, M.J. Amande, D. Gaertner, M. Capello, Machine learning for characterizing tropical tuna aggregations under Drifting Fish Aggregating Devices (DFADs) from commercial echosounder buoys data, *Fish. Res.* (2020) 229, <https://doi.org/10.1016/j.fishres.2020.105613>.
- [11] Y. Baidai, L. Dagorn, M.J. Amande, D. Gaertner, M. Capello, Tuna aggregation dynamics at drifting fish aggregating devices: a view through the eyes of commercial echosounder buoys, *ICES J. Mar. Sci.* 77 (2020) 2960–2970, <https://doi.org/10.1093/icesjms/fsaa178>.
- [12] J. Lopez, G. Moreno, I. Sancristobal, J. Murua, Evolution and current state of the technology of echo-sounder buoys used by Spanish tropical tuna purse seiners in the Atlantic, Indian and Pacific Oceans, *Fish. Res.* 155 (2014) 127–137, <https://doi.org/10.1016/j.fishres.2014.02.033>.
- [13] G. Moreno, L. Dagorn, M. Capello, J. Lopez, J. Filmlalter, F. Forget, I. Sancristobal, K. Holland, Fish aggregating devices (FADs) as scientific platforms, *Fish. Res.* (2016) 178, <https://doi.org/10.1016/j.fishres.2015.09.021>.
- [14] G. Wain, L. Guéry, D.M. Kaplan, D. Gaertner, Quantifying the increase in fishing efficiency due to the use of drifting FADs equipped with echosounders in tropical tuna purse seine fisheries, *ICES J. Mar. Sci.* 78 (2021) 235–245, <https://doi.org/10.1093/icesjms/fsaa216>.
- [15] A. Fonteneau, E. Chassot, N. Bodin, Global spatio-temporal patterns in tropical tuna purse seine fisheries on drifting fish aggregating devices (DFADs): taking a historical perspective to inform current challenges, *Aquat. Living Resour.* 48 (2013) 37–48, <https://doi.org/10.1051/alr/2013046>.
- [16] S.P. Griffiths, S.J. Nicol, V. Allain, S.D. Hoyle, T.A. Lawson, Just a FAD? Ecosystem impacts of tuna purse - seine fishing associated with fish aggregating devices in the western Pacific Warm Pool Province, *Fish. Oceanogr.* 28 (2019) 94–112, <https://doi.org/10.1111/fog.12389>.
- [17] M. Travassos Tolotti, J.D. Filmlalter, P. Bach, P. Travassos, B. Seret, L. Dagorn, Banning is not enough: the complexities of oceanic shark management by tuna regional fisheries management organizations, *Glob. Ecol. Conserv.* 4 (2015) 1–7, <https://doi.org/10.1016/j.gecco.2015.05.003>.
- [18] J.D. Filmlalter, M. Capello, J.-L. Deneubourg, P.D. Cowley, L. Dagorn, Looking behind the curtain: quantifying massive shark mortality in fish aggregating devices, *Front. Ecol. Environ.* 11 (6) (2013) 291–296, <https://doi.org/10.1890/130045>.
- [19] A. Dupaix, M. Capello, C. Lett, M. Andreollo, N. Barrier, G. Viennois, L. Dagorn, Surface habitat modification through industrial tuna fishery practices, *ICES J. Mar. Sci.* 78 (9) (2021) 3075–3088, <https://doi.org/10.1093/icesjms/fsab175>.
- [20] F. Marsac, A. Fonteneau, F. Ménard, Drifting FADs used in tuna fisheries: an ecological trap? *Proc. 1st Symp. Tuna Fish. FADs 1999 (2000)* 537–552 (Martinique, October).
- [21] T. Imzilen, C. Lett, E. Chassot, D.M. Kaplan, Spatial management can significantly reduce dFAD beachings in Indian and Atlantic Ocean tropical tuna purse seine fisheries, *Biol. Conserv.* 254 (November 2020) (2021), 108939, <https://doi.org/10.1016/j.biocon.2020.108939>.
- [22] A. Maufroy, E. Chassot, R. Joo, D.M. Kaplan, Large-scale examination of spatio-temporal patterns of drifting fish aggregating devices (dFADs) from tropical tuna fisheries of the Indian and Atlantic Oceans, *PLoS ONE* 10 (5) (2015), e0128023, <https://doi.org/10.1371/journal.pone.0128023>.
- [23] H. Murua, I. Zudaire, M. Tolotti, J. Murua, M. Capello, M. Grande, I. Arregui, J. Uranga, J. Maria, P. Sabarros, J. Ruiz, Y. Baidai, M. Lourdes, J.C. B. F. Abascal, H. Arrizabalaga, G. Moreno, L. Dagorn, Lessons learnt from the first large-scale biodegradable FAD research experiment to mitigate drifting FADs impacts on the ecosystem, *Mar. Policy* 148 (2023), 105394, <https://doi.org/10.1016/j.marpol.2022.105394>.
- [24] B. Leroy, J. Phillips, S. Nicol, G. Pilling, S. Harley, E. Al, A critique of the ecosystem impacts of drifting and anchored FADs used by purse-seine tuna fisheries in the Western and Central Pacific Ocean, *Aquat. Living Resour.* 61 (2013) 49–61, <https://doi.org/10.1051/alr/2012033>.
- [25] IOTC. (2023). RESOLUTION 23/02 ON MANAGEMENT OF DRIFTING G FISH AGGREGATING DEVICES (DFADS) IN THE IOTC AREA OF COMPETENCE.
- [26] ICCAT. (2022). RECOMMENDATION BY ICCAT REPLACING RECOMMENDATION 21–01 ON A MULTI-ANNUAL CONSERVATION AND MANAGEMENT PROGRAMME FOR TROPICAL TUNAS. 1–21.
- [27] IATTC. (2021). RESOLUTION C-21–04 CONSERVATION MEASURES FOR TROPICAL TUNAS IN THE EASTERN PACIFIC OCEAN DURING 2022–2024.
- [28] WCPFC. (2021). CONSERVATION AND MANAGEMENT MEASURE FOR BIGEYE, YELLOWFIN AND SKIPJACK TUNA IN THE WESTERN AND CENTRAL PACIFIC OCEAN. Conservation and Management Measure 2021–01. December, 1–15.
- [29] ICCAT. (2020). COMPENDIUM MANAGEMENT RECOMMENDATIONS AND RESOLUTIONS ADOPTED BY ICCAT FOR THE CONSERVATION OF ATLANTIC TUNAS AND TUNA-LIKE SPECIES.
- [30] IATTC. (2019). Inter-American Tropical Tuna Commission Resolution C-19–01 Amendment of Resolution C-18–05 on the Collection and Analyses of Data on Fish-Aggregating Devices. July, 22–26.
- [31] ISSF. (2019). Non-entangling & biodegradable FADs guide: best practices for fishers, RFMOs, governments & vessel owners.
- [32] Zudaire, I., Moreno, G., Murua, J., Murua, H., Tolotti, M.T., Roman, M., Hall, M., Lopez, J., & Grande, M. (2021). Biodegradable DFADs: Current status and prospects. *IOTC-2021-WGFAD02-09*.
- [33] Zudaire, I., Santiago, J., Grande, M., Murua, H., Adam, P., Moniz, I., & Herrera, M. (2018). FAD watch: a collaborative initiative to minimize the impact of FADs in coastal ecosystems. *14th IOTC Working Party on Ecosystems and Bycatch*. IOTC-2018-WPEB14–12.
- [34] Murua, J., Ferrarios, J.M., M., G., Moreno, G., M., Onandia, I., Ruiz, J., Zudaire, I., Santiago, J., Murua, H., & Restrepo, V. (2021). Developing solutions to increase survival rates of vulnerable bycatch species in tuna purse seiner FAD fisheries. 2nd AD-HOC WORKING GROUP ON FADS MEETING, IOTC-2021–.
- [35] F. Poisson, B. Séret, A. Vernet, M. Goujon, L. Dagorn, Collaborative research: Development of a manual on elasmobranch handling and release best practices in tropical tuna purse-seine fisheries, *Mar. Policy* 44 (2014) 312–320, <https://doi.org/10.1016/j.marpol.2013.09.025>.
- [36] F. Poisson, J.D. Filmlalter, A. Vernet, L. Dagorn, Mortality rate of silky sharks (*Carcharhinus falciformis*) caught in the tropical tuna purse seine fishery in the Indian Ocean, *Can. J. Fish. Aquat. Sci.* 71 (6) (2014) 795–798.
- [37] L. Song, H. Shen, An integrated scheme for the management of drifting fish aggregating devices in tuna purse seine fisheries, *Fish. Manag. Ecol.* 30 (2022) 56, <https://doi.org/10.1111/fme.12600>.
- [38] T.K. Davies, C.C. Mees, E.J. Milner-Gulland, The past, present and future use of drifting fish aggregating devices (FADs) in the Indian Ocean, *Mar. Policy* 45 (4) (2014) 163–170, <https://doi.org/10.1016/j.marpol.2013.12.014>.
- [39] Gershman, D., Galland, G.R., & Holmes, G. (2019). Toward true FAD deployment limits in the t-RFMOs. *J-T-RFMO FAD WG 2019, Gershman.S.04*.
- [40] G. Gomez, S. Farquhar, H. Bell, E. Laschever, S. Hall, The IUU nature of FADs: implications for tuna management and markets, *Coast. Manag.* 48 (6) (2020) 534–558, <https://doi.org/10.1080/08920753.2020.1845585>.
- [41] A. Punt, D.S. Butterworth, C.L. Moor, De, M. Haddon, Management strategy evaluation: best practices, *Fish. Res.* 17 (2016) 303–334, <https://doi.org/10.1111/faf.12104>.
- [42] S. Su, Y. Tang, J. Chen, B. Chang, Y. Chen, A comprehensive framework for operating science-based fisheries management: a checklist for using the best available science, *Fish. Res.* 22 (4) (2021) 798–811, <https://doi.org/10.1111/faf.12551>.
- [43] M. Tolotti, P. Guillotreau, F. Forget, M. Capello, L. Dagorn, Unintended effects of single - species fisheries management, *Environ. Dev. Sustain.* (2022), 0123456789, <https://doi.org/10.1007/s10668-022-02432-1>.
- [44] G. Merino, H. Arrizabalaga, I. Arregui, J. Santiago, H. Murua, Adaptation of North Atlantic albacore fishery to climate change: yet another potential benefit of harvest control rules, *Front. Mar. Sci.* 6 (2019) 620, <https://doi.org/10.3389/fmars.2019.00620>.
- [45] R. Sharma, P. Levontin, T. Kitakado, L. Kell, I. Mosqueira, A. Kimoto, R. Scott, C. M. Paul, Y. Ye, J. Kleiber, J. Lindsay, W. Shana, A. Magnusson, Operating model design in tuna regional fishery management organizations: current practice, issues and implications, May, 940–961, *Fish. Res.* 21 (2020) 940, <https://doi.org/10.1111/faf.12480>.
- [46] S.M. Garcia, The precautionary principle: its implications in capture fisheries management, *Ocean Coast. Manag.* 22 (1994) 99–125, [https://doi.org/10.1016/0964-5691\(94\)90014-0](https://doi.org/10.1016/0964-5691(94)90014-0).
- [47] Duparc, A., Cauquil, P., Depetris, M., Dewals, P., Floch, L., Gaertner, D., Hervé, A., Lebranchu, J., Marsac, F., & Bach, P. (2018). Assessment of accuracy in processing purse seine tropical tuna catches with the T3 methodology using French fleet data. *IOTC-2018-WPTT20-16*.
- [48] Peatman, T., Williams, P., & Nicol, S. (2022). *Project 60: Progress towards achieving SC17 recommendations*. WCPFC-SC18–2022/ST-IP-03.
- [49] J. Hallier, D. Gaertner, Drifting fish aggregation devices could act as an ecological trap for tropical tuna species, *Mar. Ecol. Prog. Ser.* 353 (2008) 255–264, <https://doi.org/10.3354/meps07180>.
- [50] S. Jaquemet, M. Potier, F. Ménard, Do drifting and anchored Fish Aggregating Devices (FADs) similarly influence tuna feeding habits? A case study from the western Indian Ocean, *Fish. Res.* 107 (1–3) (2011) 283–290, <https://doi.org/10.1016/j.fishres.2010.11.011>.
- [51] M. Robert, L. Dagorn, N. Bodin, F. Pernet, E.J. Arsenault-Pernet, J.L. Deneubourg, Comparison of condition factors of skipjack tuna (*Katsuwonus pelamis*) associated or not with floating objects in an area known to be naturally enriched with logs, *Can. J. Fish. Aquat. Sci.* 71 (3) (2014) 472–478, <https://doi.org/10.1139/cjfas-2013-0389>.
- [52] I. Zudaire, H. Murua, M. Grande, N. Goñi, M. Potier, F. Ménard, E. Chassot, N. Bodin, Variations in the diet and stable isotope ratios during the ovarian development of female yellowfin tuna (*Thunnus albacares*) in the Western Indian Ocean, *Mar. Biol.* 162 (12) (2015) 2363–2377, <https://doi.org/10.1007/s00227-015-2763-0>.
- [53] I. Zudaire, H. Murua, M. Grande, F. Pernet, N. Bodin, Accumulation and mobilization of lipids in relation to reproduction of yellowfin tuna (*Thunnus albacares*) in the Western Indian Ocean, *Fish. Res.* 160 (2014) 50–59, <https://doi.org/10.1016/j.fishres.2013.12.010>.

- [54] S. Dueri, L. Bopp, O. Maury, Projecting the impacts of climate change on skipjack tuna abundance and spatial distribution, *Glob. Change Biol.* 20 (3) (2014) 742–753, <https://doi.org/10.1111/gcb.12460>.
- [55] M.J. Amandé, J. Ariz, E. Chassot, A.D. De Molina, D. Gaertner, H. Murua, R. Pianet, J. Ruiz, P. Chavance, Bycatch of the European purse seine tuna fishery in the Atlantic Ocean for the 2003–2007 period, *Aquat. Living Resour.* 23 (4) (2010) 353–362, <https://doi.org/10.1051/alr/2011003>.
- [56] E.L. Gilman, Bycatch governance and best practice mitigation technology in global tuna fisheries, *Mar. Policy* 35 (5) (2011) 590–609, <https://doi.org/10.1016/j.marpol.2011.01.021>.
- [57] IUCN. (2022). The IUCN Red List of Threatened Species.
- [58] N. Lezama-Ochoa, H. Murua, J. Ruiz, P. Chavance, A. Delgado de Molina, A. Caballero, I. Sancristobal, Biodiversity and environmental characteristics of the bycatch assemblages from the tropical tuna purse seine fisheries in the eastern Atlantic Ocean, *Mar. Ecol.* 39 (3) (2018), <https://doi.org/10.1111/maec.12504>.
- [59] E. Torres-Irineo, M.J. Amandé, D. Gaertner, A.D. de Molina, H. Murua, P. Chavance, J. Ariz, J. Ruiz, N. Lezama-Ochoa, Bycatch species composition over time by tuna purse-seine fishery in the eastern tropical Atlantic Ocean, *Biodivers. Conserv.* 23 (5) (2014) 1157–1173, <https://doi.org/10.1007/s10531-014-0655-0>.
- [60] C. Eddy, R. Brill, D. Bernal, Rates of at-vessel mortality and post-release survival of pelagic sharks captured with tuna purse seines around drifting fish aggregating devices (FADs) in the equatorial eastern Pacific Ocean, *Fish. Res.* 174 (2016) 109–117, <https://doi.org/10.1016/j.fishres.2015.09.008>.
- [61] M.R. Hutchinson, D.G. Itano, J.A. Muir, K.N. Holland, Post-release survival of juvenile silky sharks captured in a tropical tuna purse seine fishery, *Mar. Ecol. Prog. Ser.* 521 (2015) 143–154, <https://doi.org/10.3354/meps11073>.
- [62] Onandia I, Grande M, Galaz JM, et al (2021) New assessment on accidentally captured silky shark post-release survival in the Indian Ocean tuna purse seine fishery. In: IOTC- 17th Working Party on Ecosystems & Bycatch (Data Prep). IOTC-2021-WPEB17(DP)-13 Rev1.
- [63] Murua, J., Moreno, G., Hall, M., Dagorn, L., Itano, D., & Restrepo, V. (2017). Towards global non-entangling fish aggregating device (FAD) use in tropical tuna purse seine fisheries through a participatory approach. *ISSF Technical Report 2017-07*, December, 25.
- [64] L. Bonnin, C. Lett, L. Dagorn, J.D. Filmlalter, F. Forget, P. Verley, M. Capello, Can drifting objects drive the movements of a vulnerable pelagic shark? *Aquat. Conserv.: Mar. Freshw. Ecosyst.* 31 (2020) 74–82, <https://doi.org/10.1002/aqc.3420>.
- [65] M.T. Tolotti, F. Forget, M. Capello, J.D. Filmlalter, M. Hutchinson, D. Itano, K. Holland, L. Dagorn, Association dynamics of tuna and purse seine bycatch species with drifting fish aggregating devices (FADs) in the tropical eastern Atlantic Ocean, *Fish. Res.* (2020) 226, <https://doi.org/10.1016/j.fishres.2020.105521>.
- [66] L. Escalle, J.S. Phillips, M. Brownjohn, S. Brouwer, A. Gupta, Sen, E. Sebille, Van, J. Hampton, G. Pilling, Environmental versus operational drivers of drifting FAD beaching in the Western and Central Pacific Ocean, *Sci. Rep.* 9 (2019) 14005, <https://doi.org/10.1038/s41598-019-50364-0>.
- [67] T. Imzilen, C. Lett, E. Chassot, A. Maufroy, M. Goujon, D.M. Kaplan, Recovery at sea of abandoned, lost or discarded drifting fish aggregating devices, *Nat. Sustain.* 5 (7) (2022) 593–602, <https://doi.org/10.1038/s41893-022-00883-y>.
- [68] Escalle, L. et al. (2022). Preliminary analyses of the regional database of stranded drifting FADs in the Pacific Ocean. *WCPFC Sci. Comm. WCPFC-SC18-2022/EB-IP-03*.
- [69] IOTC. (2021). REVIEW OF DATA ON DRIFTING FISH AGGREGATING DEVICES, IOTC-2021-WGFAD02-03 Rev1. 2nd IOTC FAD Working Group, 1–51.
- [70] G. Moreno, J. Salvador, I. Zudaire, J. Murua, J. Lluís, J. Uranga, H. Murua, M. Grande, J. Santiago, V. Restrepo, The jelly-FAD: a paradigm shift in the design of biodegradable fish aggregating devices, *Mar. Policy* 147 (2023), 105352, <https://doi.org/10.1016/j.marpol.2022.105352>.
- [71] I. Zudaire, et al., Biodegradable drifting fish aggregating devices: Current status and future prospects, *Marine Policy* 153 (2023), 105659, <https://doi.org/10.1016/j.marpol.2023.105659>. ISSN 0308-597X.
- [72] F. Ménard, A. Fonteneau, D. Gaertner, V. Nordstrom, B. Stéquent, E. Marchal, Exploitation of small tunas by a purse-seine fishery with fish aggregating devices and their feeding ecology in an eastern tropical Atlantic ecosystem, *ICES J. Mar. Sci.* 57 (3) (2000) 525–530, <https://doi.org/10.1006/jmsc.2000.0717>.
- [73] Ducharme-Barth, N., Vincent, M., Hampton, J., Hamer, P., Williams, P., & Pilling, G. (2020). *Stock assessment of bigeye tuna in the western and central Pacific Ocean WCPFC-SC16-2020/SA-WP-03. August*. <https://doi.org/10.13140/RG.2.2.14696.90885>.
- [74] G. Sempo, L. Dagorn, M. Robert, J.-L. Deneubourg, Impact of increasing deployment of artificial floating objects on the spatial distribution of social fish species, *J. Appl. Ecol.* 50 (2013) 1081, <https://doi.org/10.1111/1365-2664.12140>.
- [75] C.E. Lennert-cody, G. Moreno, V. Restrepo, M.H. Roma, M.N. Maunder, Recent purse-seine FAD fishing strategies in the eastern Pacific Ocean: what is the appropriate number of FADs at sea ? *ICES J. Mar. Sci.* 75 (2018) 1748–1757, <https://doi.org/10.1093/icesjms/fsy046>.
- [76] M. Capello, J.L. Deneubourg, M. Robert, K.N. Holland, K.M. Schaefer, L. Dagorn, Population assessment of tropical tuna based on their associative behavior around floating objects, *Sci. Reports* 6 (2016) 36415, <https://doi.org/10.1038/srep36415>.
- [77] Dupaix, A., Dagorn, L., Deneubourg, J.-L., & Capello, M. (2022). Impact of DFAD density on tuna associative behavior and catchability in the Indian Ocean. *IOTC-2022-WGFAD03-06*.
- [78] M. Capello, J. Rault, J. Deneubourg, L. Dagorn, Schooling in habitats with aggregative sites: the case of tropical tuna and floating objects, *J. Theor. Biol.* 547 (2022), 111163, <https://doi.org/10.1016/j.jtbi.2022.111163>.
- [79] D.J. Curnick, D.A. Feary, G.H. Cavalcante, Risks to large marine protected areas posed by drifting fish aggregation devices, *Conserv. Biol.* 35 (4) (2020) 1222–1232, <https://doi.org/10.1111/cobi.13684>.
- [80] T. Imzilen, E. Chassot, J. Barde, H. Demarcq, A. Maufroy, L. Roa-pascuali, J. Ternon, Fish aggregating devices drift like oceanographic drifters in the near-surface currents of the Atlantic and Indian Oceans, *Prog. Oceanogr.* 171 (May 2018) (2019) 108–127, <https://doi.org/10.1016/j.poccean.2018.11.007>.
- [81] C. Lett, P. Verley, C. Mullon, C. Parada, A Lagrangian tool for modelling ichthyoplankton dynamics, *Environ. Model. Softw.* 23 (2008) 1210–1214, <https://doi.org/10.1016/j.envsoft.2008.02.005>.
- [82] A. Fonteneau, J. Ariz, D. Gaertner, V. Nordstrom, P. Pallares, Observed changes in the species composition of tuna schools in the Gulf of Guinea between 1981 and 1999, in relation with the fish aggregating, *Aquat. Living Resour.* 13 (2000) 253–257, [https://doi.org/10.1016/S0990-7440\(00\)01054-8](https://doi.org/10.1016/S0990-7440(00)01054-8).
- [83] A. Maufroy, D.M. Kaplan, N. Bez, A.D. Molina, De, H. Murua, L. Floch, E. Chassot, Massive increase in the use of drifting Fish Aggregating Devices (dFADs) by tropical tuna purse seine fisheries in the Atlantic and Indian oceans, *ICES J. Mar. Sci.* 74 (2017) 215–225, <https://doi.org/10.1093/icesjms/fsw175>.
- [84] E. Chassot, S. Antoine, P. Guillotreau, J. Lucas, C. Assan, M. Marguerite, N. Bodin, Fuel consumption and air emissions in one of the world's largest commercial fisheries, *Environ. Pollut.* 273 (2021), 116454, <https://doi.org/10.1016/j.envpol.2021.116454>.
- [85] G. Merino, H. Murua, J. Santiago, H. Arrizabalaga, V. Restrepo, Characterization, communication, and management of uncertainty in tuna fisheries, *Sustainability* 12 (19) (2020) 8245, <https://doi.org/10.3390/su12198245>.
- [86] F.-X. Bard, J.-M. Stretta, M. Slepoukha, Les épaves artificielles comme auxiliaires de la pêche thonière en océan Atlantique: quel avenir? *Pêche Marit.* 1291 (1985) 655–659. (<http://www.documentation.ird.fr/fdi/notice.php?niv=fdi:010022579>).
- [87] S.C. Clarke, M.K. McAllister, E.J. Milner-Gulland, G.P. Kirkwood, C.G. J. Michielsens, D.J. Agnew, E.K. Pikitch, H. Nakano, M.S. Shivji, Global estimates of shark catches using trade records from commercial markets, *Ecol. Lett.* 9 (10) (2006) 1115–1126, <https://doi.org/10.1111/j.1461-0248.2006.00968.x>.
- [88] V. Huckstorff, W.-C. Lewin, T. Mehner, C. Wolter, Performance level and efficiency of two differing predator-avoidance strategies depend on nutritional state of the prey fish, *Behav. Ecol. Sociobiol.* 63 (12) (2009) 1735–1742, <https://doi.org/10.1007/s00265-009-0790-x>.
- [89] M.C. Melnychuk, E. Peterson, M. Elliott, R. Hilborn, Fisheries management impacts on target species status, *Proc. Natl. Acad. Sci. USA* 114 (1) (2017) 178–183, <https://doi.org/10.1073/pnas.1609915114>.
- [90] W.S. Patrick, J.S. Link, Myths that continue to impede progress in ecosystem-based fisheries management, *Fisheries* 40 (4) (2015) 155–160, <https://doi.org/10.1080/03632415.2015.1024308>.
- [91] E.K. Pikitch, C. Santora, E.A. Babcock, A. Bakun, R. Bonfil, D.O. Conover, P. Dayton, P. Doukakis, D. Fluharty, B. Heneman, E.D. Houde, J. Link, P. A. Livingston, M. Mangel, M.K. McAllister, J. Pope, K.J. Sainsbury, Ecosystem-based fishery management, *Science* 305 (5682) (2004) 346–347, <https://doi.org/10.1126/science.1098222>.
- [92] T.J. Pitcher, D. Kalikoski, K. Short, D. Varkey, G. Pramod, An evaluation of progress in implementing ecosystem-based management of fisheries in 33 countries, *Mar. Policy* 33 (2) (2009) 223–232, <https://doi.org/10.1016/j.marpol.2008.06.002>.
- [93] J. Santiago, J. Lopez, G. Moreno, H. Murua, I. Quincoces, M. Soto, Towards a tropical tuna buoy-derived abundance index (TT-BAI), *Collect. Vol. Sci. Pap. ICCAT* 72 (3) (2016) 714–724.