

## **Draft executive summary of the status of the blue shark resource**

Draft dated 11 July 2007

### **BIOLOGY**

The blue shark (*Prionace glauca*) is common in pelagic oceanic waters throughout the tropical and temperate oceans worldwide. It has one of the widest ranges of all the shark species. It may also be found close inshore and in estuaries. Blue shark is most common in relatively cool waters (7 to 16°C) often close to the surface. In the tropical Indian Ocean, the greatest abundance of blue sharks occurs at depths of 80 to 220 m, in temperatures ranging from 12 to 25°C. The distribution and movements of blue shark are strongly influenced by seasonal variations in water temperature, reproductive condition, and availability of prey.

The blue shark is often found in large single sex schools containing individuals of similar size. Adult blue sharks have no known predators; however, subadults and juveniles are eaten by both shortfin makos and white sharks as well as by sea lions. Fishing is likely to be a major contributor to adult mortality.

In the Atlantic Ocean, the oldest blue sharks reported were a 16 year old male and a 13 year old female. Longevity is estimated to be between 20-26 years of age and maximum size is around 3.8 m FL. Size increases when latitude decreases.

Sexual maturity is attained at 5 years of age in both sexes. Blue shark is a viviparous species, with a yolk-sac placenta. Once the eggs have been fertilised there is a gestation period of between 9 and 12 months. Litter size is quite variable, ranging from four to 135 pups and may be dependent on the size of the female. The average litter size observed from the Indian Ocean is 38. New-born pups are around 40 to 51 cm in length. Generation time is about eight years. In Indian Ocean, between latitude 2 °N and 6 °S, pregnant females are present for most of the year.

### **FISHERIES**

Blue sharks are often targeted by some semi-industrial, artisanal and recreational fisheries and are a bycatch of industrial fisheries (pelagic longline tuna and swordfish fisheries and purse seine fishery). The blue shark appears to have a similar distribution to swordfish. Typically, the fisheries take blue sharks between 1.8-2.4 m fork length or 30 to 52 kg. Males are slightly smaller than the females. In other Oceans, angling clubs are known for organising sharks fishing competitions where blue sharks and mako sharks are targeted. Sport fisheries for sharks are apparently not so common in the Indian Ocean.

There is little information on the fisheries prior to the early 1970's, and some countries continue not to collect shark data while others do collect it but do not report it to IOTC. It appears that significant catches of sharks have gone unrecorded in several countries. Furthermore, many catch records probably under-represent the actual catches of sharks because they do not account for discards (i.e. do not record catches of sharks for which only the fins are kept or of sharks usually discarded because of their size or condition) or they reflect dressed weights instead of live weights.

In 2005, seven countries reported catches of blue sharks in the IOTC region. These are not given in this summary because their representativeness is highly uncertain. Apparently, as other shark stocks have declined less blue sharks are being discarded.

FAO also compiles landings data on elasmobranchs, but the statistics are limited by the lack of species-specific data and data from the major fleets.

## **AVAILABILITY OF INFORMATION FOR STOCK ASSESSMENT**

There is little information on blue shark biology and no information is available on stock structure.

Possible fishery indicators:

1. **Trends in catches:** The catch estimates for blue shark are highly uncertain as is their utility in terms of minimum catch estimates.
2. **Nominal CPUE Trends:** Data not available. There are no surveys specifically designed to assess shark catch rates in the Indian Ocean. Trends in localised areas might be possible in the future (for example, from the Kenyan recreational fishery).
3. **Average weight in the catch by fisheries:** data not available
4. **Number of squares fished:** CE data not available

## **STOCK ASSESSMENT**

No quantitative stock assessment has been undertaken by the IOTC Working Party on Ecosystems and Bycatch.

## **MANAGEMENT ADVICE**

There is a paucity of information available on this species and this situation is not expected to improve in the short to medium term. There is no quantitative stock assessment or basic fishery indicators currently available for blue shark in the Indian Ocean therefore the stock status is highly uncertain.

Blue sharks are commonly taken by a range of fisheries in the Indian Ocean and in some areas they are fished in their nursery grounds. Because of their life history characteristics – they are relatively long lived (16-20 years), mature at 4-6 years, and have relatively few offspring (25-50 pups every two years), the blue shark is vulnerable to overfishing.