

DRAFT executive summary of the status of the scalloped hammerhead shark resource

Draft dated 11 July 2007

BIOLOGY

The scalloped hammerhead shark (*Sphyrna lewini*) is widely distributed and common in warm temperate and tropical waters down to 275 m. It is also found in estuarine and inshore waters.

In some areas, the scalloped hammerhead shark forms large resident populations. In other areas, large schools of small-sized sharks are known to migrate pole wards seasonally.

Scalloped hammerhead sharks feeds on pelagic fishes, other sharks and rays, squids, lobsters, shrimps and crabs.

The maximum age for Atlantic Ocean scalloped hammerheads is estimated to be over 30 years with the largest individuals reaching over 2.4 m.

Males in the Indian Ocean mature at around 1.4-1.65 m TL. Females mature at about 2.0 m TL. The scalloped hammerhead shark is viviparous with a yolk sac-placenta. The young are around 38-45 cm TL at birth, and litters consist of 15-31 pups. The reproductive cycle is annual and the gestation period is 9-10 months. The nursery areas are in shallow coastal waters.

FISHERIES

Scalloped hammerhead sharks are often targeted by some semi-industrial, artisanal and recreational fisheries and are a bycatch of industrial fisheries (pelagic longline tuna and swordfish fisheries and purse seine fishery).

There is little information on the fisheries prior to the early 1970's, and some countries continue not to collect shark data while others do collect it but do not report it to IOTC. It appears that significant catches of sharks have gone unrecorded in several countries. Furthermore, many catch records probably under-represent the actual catches of sharks because they do not account for discards (i.e. do not record catches of sharks for which only the fins are kept or of sharks usually discarded because of their size or condition) or they reflect dressed weights instead of live weights.

Catches of scalloped hammerhead sharks in the IOTC region are not given in this summary because their representativeness is highly uncertain.

FAO also compiles landings data on elasmobranchs, but the statistics are limited by the lack of species-specific data and data from the major fleets.

AVAILABILITY OF INFORMATION FOR STOCK ASSESSMENT

There is little information available on scalloped hammerhead shark biology and no information is available on stock structure.

Possible fishery indicators:

1. **Trends in catches:** The catch estimates scalloped hammerhead are highly uncertain as is their utility in terms of minimum catch estimates.
2. **Nominal CPUE Trends:** data not available
3. **Average weight in the catch by fisheries:** data not available
4. **Number of squares fished:** CE data not available

STOCK ASSESSMENT

No quantitative stock assessment has been undertaken by the IOTC Working Party on Ecosystems and Bycatch.

MANAGEMENT ADVICE

There is a paucity of information available on this species and this situation is not expected to improve in the short to medium term. There is no quantitative stock assessment or basic fishery indicators currently available for scalloped hammerhead shark in the Indian Ocean therefore the stock status is highly uncertain.

Scalloped hammerhead sharks are commonly taken by a range of fisheries in the Indian Ocean. They are extremely vulnerable to gillnet fisheries. Furthermore, pups occupy shallow coastal nursery grounds, often heavily exploited by inshore fisheries. Because of their life history characteristics – they are relatively long lived (over 30 years), and have relatively few offspring (<31 pups each year), the scalloped hammerhead shark is vulnerable to overfishing.