

EXECUTIVE SUMMARY: STATUS OF THE INDIAN OCEAN YELLOWFIN TUNA (*THUNNUS ALBACARES*) RESOURCE

TABLE 1. Status of yellowfin tuna (*Thunnus albacares*) in the Indian Ocean.

Area ¹	Indicators – 2011 assessment		2011 stock status determination
			2009 ²
Indian Ocean	Catch 2010 (1000 t): Average catch 2006–2010 (1000 t): MSY: F ₂₀₀₉ /F _{MSY} : SB ₂₀₀₉ /SB _{MSY} : SB ₂₀₀₉ /SB ₀ :	299.1 326.6 357 (290–435) 0.84 (0.63–1.10) 1.61 (1.47–1.78) 0.35 (0.31–0.38)	

¹Boundaries for the Indian Ocean stock assessment are defined as the IOTC area of competence.

²The stock status refers to the most recent years' data used for the assessment.

Colour key	Stock overfished (SB _{year} /SB _{MSY} < 1)	Stock not overfished (SB _{year} /SB _{MSY} ≥ 1)
Stock subject to overfishing (F _{year} /F _{MSY} > 1)		
Stock not subject to overfishing (F _{year} /F _{MSY} ≤ 1)		

INDIAN OCEAN STOCK – MANAGEMENT ADVICE

The WPTT **RECOMMENDED** the following management advice for yellowfin tuna in the Indian Ocean, for the consideration of the Scientific Committee.

Stock status. The stock assessment model used in 2011 suggests that the stock is currently not overfished ($B_{2009} > B_{MSY}$) and overfishing is not occurring ($F_{2009} < F_{MSY}$) (Table 1 and Fig. 1). Spawning stock biomass in 2009 was estimated to be 35% (31–38%) (from Table 1) of the unfished levels. However, estimates of total and spawning stock biomass show a marked decrease over the last decade, accelerated in recent years by the high catches of 2003–2006. Recent reductions in effort and, hence, catches has halted the decline.

The main mechanism that appears to be behind the very high catches in the 2003–2006 period is an increase in catchability by surface and longline fleets due to a high level of concentration across a reduced area and depth range. This was likely linked to the oceanographic conditions at the time generating high concentrations of suitable prey items that yellowfin tuna exploited. A possible increase in recruitment in previous years, and thus in abundance, cannot be completely ruled out, but no signal of it is apparent in either data or model results. This means that those catches probably resulted in considerable stock depletion.

Outlook. The decrease in longline and purse seiner effort in recent years has substantially lowered the pressure on the Indian Ocean stock as a whole, indicating that current fishing mortality has not exceeded the MSY-related levels in recent years. If the security situation in the western Indian Ocean were to improve, a rapid reversal in fleet activity in this region may lead to an increase in effort which the stock might not be able to sustain, as catches would then be likely to exceed MSY levels. Catches in 2010 (299,074 t) are within the lower range of MSY values. The current assessment indicates that catches of about the 2010 level are sustainable, at least in the short term. However, the stock is unlikely to support higher yields based on the estimated levels of recruitment from over the last 15 years.

In 2011, the WPTT undertook projections of yellowfin tuna stock status under a range of management scenarios for the first time, following the recommendation of both the Kobe process and the Commission, to harmonise technical advice to managers across RFMOs by producing Kobe II management strategy matrices. The purpose of the table is to quantify the future outcomes from a range of management options (Table 2). The table describes the presently estimated probability of the population being outside biological reference points at some point in the future, where “outside” was assigned the default definitions of $F > F_{MSY}$ or $B < B_{MSY}$. The timeframes represent 3 and 10 year projections (from the last data in the model), which corresponds to predictions for 2013 and 2020. The management options represent three different levels of constant catch projection: catches 20% less than 2010, equal to 2010 and 20% greater than 2010.

The projections were carried out using 12 different scenarios based on similar scenarios used in the assessment for the combination of those different MFCL runs: LL selectivity flat top vs. dome shape; steepness values of 0.7, 0.8 and 0.9; and computing the recruitment as an average of the whole time series vs. 15 recent years (12 scenarios). The probabilities in the matrices were computed as the percentage of the 12 scenarios being $B > B_{MSY}$ and $F < F_{MSY}$ in each year. In that sense, there are not producing the uncertainty related to any specific scenario but the uncertainty associated to different scenarios.

There was considerable discussion on the ability of the WPTT to carry out the projections with MFCL for yellowfin tuna. For example, it was not clear how the projection redistributed the recruitment among regions as recent distribution of recruitment differs from historic; which was assumed in the projections. The WPTT agreed that the true uncertainty is unknown and that the current characterization is not complete; however, the WPTT feels that the projections may provide a relative ranking of different scenarios outcomes. The WPTT recognised at this time that the matrices do not represent the full range of uncertainty from the assessments. Therefore, the inclusion of the K2SM at this time is primarily intended to familiarise the Commission with the format and method of presenting management advice.

The WPTT **RECOMMENDED** that the Scientific Committee consider the following:

- The Maximum Sustainable Yield estimate for the whole Indian Ocean is 357,000 t with a range between 290,000–435,000 t (Table 1), and annual catches of yellowfin tuna should not exceed the lower range of MSY (300,000 t) in order to ensure that stock biomass levels could sustain catches at the MSY level in the long term.
- Recent recruitment is estimated to be considerably lower than the whole time series average. If recruitment continues to be lower than average, catches below MSY would be needed to maintain stock levels.




Fig. 1. MULTIFAN-CL Indian Ocean yellowfin tuna stock assessment Kobe plot. Blue circles indicate the trajectory of the point estimates for the B ratio and F ratio for each year 1972–2009. The equal weighted mean trajectory of the scenarios investigated in the assessment. The range is given by the different scenarios investigated.

TABLE 2. MULTIFAN-CL Indian Ocean yellowfin tuna stock assessment Kobe II Strategy Matrix. Percentage probability of violating the MSY-based reference points for five constant catch projections (2010 catch level, $\pm 20\%$ and $\pm 40\%$) projected for 3 and 10 years. In the projection, however, 12 scenarios were investigated: the six scenarios investigated above as well as the same scenarios but with a lower mean recruitment assumed for the projected period.

Reference point and projection timeframe	Alternative catch projections (relative to 2010) and probability (%) of violating reference point				
	60%	80%	100%	120%	140%
	(165,600 t)	(220,800 t)	(276,000 t)	(331,200 t)	(386,400 t)
$B_{2013} < B_{MSY}$	<1	<1	<1	<1	<1
$F_{2013} > F_{MSY}$	<1	<1	58.3	83.3	100
$B_{2020} < B_{MSY}$	<1	<1	8.3	41.7	91.7
$F_{2020} > F_{MSY}$	<1	41.7	83.3	100	100

SUPPORTING INFORMATION

(Information collated from reports of the Working Party on Tropical Tunas and other sources as cited)

CONSERVATION AND MANAGEMENT MEASURES

Yellowfin tuna (*Thunnus albacares*) in the Indian Ocean is currently subject to a number of conservation and management measures adopted by the Commission:

- Resolution 08/04 concerning the recording of catch by longline fishing vessels in the IOTC area.
- Resolution 09/02 On the implementation of a limitation of fishing capacity of contracting parties and cooperating non-contracting parties.
- Resolution 10/02 mandatory statistical requirements for IOTC Members and Cooperating non-Contracting Parties (CPC's).
- Resolution 10/03 concerning the recording of catch by fishing vessels in the IOTC area.
- Resolution 10/07 concerning a record of licensed foreign vessels fishing for tunas and swordfish in the IOTC area.
- Resolution 10/08 concerning a record of active vessels fishing for tunas and swordfish in the IOTC area.
- Recommendation 10/13 On the implementation of a ban on discards of skipjack tuna, yellowfin tuna, bigeye tuna, and non targeted species caught by purse seiners.
- Recommendation 11/06 Concerning the Recording of Catch by Fishing Vessels in the IOTC Area of Competence.

FISHERIES INDICATORS

General

Yellowfin tuna (*Thunnus albacares*) is a cosmopolitan species distributed mainly in the tropical and subtropical oceanic waters of the three major oceans, where it forms large schools. Table 3 outlines some of the key life history traits of yellowfin tuna relevant for management.

TABLE 3. Biology of Indian Ocean yellowfin tuna (*Thunnus albacares*)

Parameter	Description
Range and stock structure	A cosmopolitan species distributed mainly in the tropical and subtropical oceanic waters of the three major oceans, where it forms large schools. Feeding behaviour has been extensively studied and it is largely opportunistic, with a variety of prey species being consumed, including large concentrations of crustaceans that have occurred recently in the tropical areas and small mesopelagic fishes which are abundant in the Arabian Sea. It has also been observed that large individuals can feed on very small prey, thus increasing the availability of food for this species. Archival tagging of yellowfin tuna has shown that this species can dive very deep (over 1000 m) probably to feed on meso-pelagic prey. Longline catch data indicates that yellowfin tuna are distributed throughout the entire tropical Indian Ocean. The tag recoveries of the RTTP-IO provide evidence of large movements of yellowfin tuna, thus supporting the assumption of a single stock for the Indian Ocean. The average distance travelled by yellowfin between being tagging and recovered is 710 nautical miles, and showing increasing distances as a function of time at sea.
Longevity	9 years
Maturity (50%)	Age: females and males 3–5 years. Size: females and males 100 cm.
Spawning season	Spawning occurs mainly from December to March in the equatorial area (0-10°S), with the main spawning grounds west of 75°E. Secondary spawning grounds exist off Sri Lanka and the Mozambique Channel and in the eastern Indian Ocean off Australia.
Size (length and weight)	Maximum length: 240 cm FL; Maximum weight: 200 kg. Newly recruited fish are primarily caught by the purse seine fishery on floating objects. Males are predominant in the catches of larger fish at sizes than 140 cm (this is also the case in other oceans). The sizes exploited in the Indian Ocean range from 30 cm to 180 cm fork length. Smaller fish (juveniles) form mixed schools with skipjack tuna and juvenile bigeye tuna and are mainly limited to surface tropical waters, while larger fish are found in surface and sub-surface waters. Intermediate age yellowfin tuna are seldom taken in the industrial fisheries, but are abundant in some artisanal fisheries, mainly in the Arabian Sea.

SOURCES: Froese & Pauly (2009)

Catch trends

Contrary to the situation in other oceans, the artisanal fishery component of yellowfin tuna catches in the Indian Ocean is substantial, taking approximately 20–25% of the total catch landed. Catches of yellowfin tuna remained more or less stable between the mid-1950s and the early-1980s, ranging between 30,000 and 70,000 t, owing to the activities of longliners and, to a lesser extent, gillnetters (Fig. 2).

Catches of yellowfin tuna increased rapidly with the arrival of the purse seine fleets in the early 1980s (Figs. 2 and 3), along with increased activity by longline vessels, with more than 400,000 t landed in 1993. Purse seiners typically take fish ranging from 40–140 cm fork length and smaller fish are more common in the catches taken north of the equator.

The purse seine fishery is characterized by the use of two different fishing modes: a fishery on drifting objects (FADs), which catches large numbers of small yellowfin in association with skipjack tuna and juvenile bigeye tuna, and a fishery on free swimming schools, which catches larger yellowfin tuna on multi-specific or mono-specific sets. Between 1995 and 2003, the FAD component of the purse seine fishery represented 48–66% of the sets undertaken (60–80% of the positive sets) and took 36–63% of the yellowfin tuna catch by weight (59–76% of the total catch). The proportion of yellowfin tuna caught (in weight) on free-schools during 2003–2006 (64%) was much higher than in previous (49% for 1999–2002) or following years (55% for 2007–2009).

The longline fishery primarily catches large fish, from 80–160 cm fork length, although smaller fish in the size range 60–100 cm have been taken and reported by longliners from Taiwan,China since 1989 in the Arabian Sea. The longline fishery targets several tuna species in different parts of the Indian Ocean, with yellowfin tuna and bigeye tuna being the main target species in tropical waters. The longline fishery can be subdivided into a deep-freezing longline component (large scale deep-freezing longliners operating on the high seas from Japan, Rep. of Korea and Taiwan,China) and a fresh-tuna longline component (small to medium scale fresh tuna longliners from Indonesia and Taiwan,China). As was the case with purse seine fisheries, since 2005 longline catches have decreased substantially with current catches estimated to be at around 41,000 t, representing a more than three-fold decrease over the catches in 2005 (Fig. 2).

Total yellowfin tuna catches dropped markedly from the peak catches taken in 2006, with the lowest catches recorded since the early 1990's reported in 2009, at around 275,955 t. Preliminary catch levels in 2010 are estimated to be around 299,074 t (Tables 4, 5).

The recent drop in catches of yellowfin tuna could be related, at least in part, to the expansion of piracy in the western tropical Indian Ocean, which has led to a marked drop in the levels of longline effort in the core fishing area of the species (Figs. 4a, b) as well as to the decline in the number of purse seiners in the Indian Ocean (~30% reduction).

Catches by other gears, i.e. pole-and-line, gillnet, troll, hand line and other minor gears, have increased steadily since the 1980s (Fig. 2). In recent years the total artisanal yellowfin tuna catch has been between 140,000–160,000 t, with the catch by gillnets (the dominant artisanal gear) at around 80,000 t.

Most yellowfin tuna are caught in the Indian Ocean, north of 12°S, and in the north of the Mozambique Channel (Figs. 4a, b). In recent years the catches of yellowfin tuna in the western Indian Ocean have dropped considerably, especially in areas off Somalia, Kenya and Tanzania and in particular between 2008 and 2010. The drop in catches is the consequence of a generalised drop in fishing effort due to the effect of piracy in the western Indian Ocean region.




Fig. 2. Annual catches of yellowfin tuna by gear recorded in the IOTC Database (1961–2010) (Data as of September 2011).




Fig. 3. Annual catches of yellowfin tuna by fleet recorded in the IOTC Database (1961–2010) (Data as of September 2011).




Fig. 4a–b. Time-area catches (total combined in tonnes) of yellowfin tuna estimated for 2009 and 2010 by type of gear: Longline (LL), Purse seine free-schools (FS), Purse seine associated-schools (LS), pole-and-line (BB), and other fleets (OT), including drifting gillnets, and various coastal fisheries (Data as of September 2011).

TABLE 4. Best scientific estimates of the catches of yellowfin tuna (*Thunnus albacares*) by gear and main fleets [or type of fishery] by decade (1950–2000) and year (2001–2010), in tonnes. Data as of October 2011. Catches by decade represent the average annual catch, noting that some gears were not used for all years (refer to Fig. 2).

Fishery	By decade (average)						By year (last ten years)									
	1950s	1960s	1970s	1980s	1990s	2000s	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
FS	0	0	18	32,590	64,942	89,761	78,969	77,059	137,492	168,799	124,024	85,021	53,529	74,990	36,263	31,951
LS	0	0	17	18,090	56,304	61,909	50,997	61,933	86,585	59,597	69,873	74,454	43,843	41,453	51,565	72,199
LL	21,990	41,256	29,512	33,889	66,689	57,668	43,932	53,132	55,741	86,415	116,847	69,831	54,414	29,128	21,242	17,130
LF	0	0	615	4,286	47,570	32,827	39,323	34,429	31,292	31,125	33,991	30,475	28,752	30,424	23,157	24,089
BB	1,754	1,452	4,380	6,621	11,765	17,162	14,233	19,393	19,451	16,177	16,607	18,644	18,133	18,351	18,463	12,755
GI	2,604	7,569	12,861	15,261	50,192	76,053	60,748	62,982	83,283	99,254	76,660	86,286	66,693	80,086	82,695	101,418
HD	679	1,175	2,615	6,990	20,002	31,762	29,790	34,093	31,105	40,820	38,993	31,789	30,274	28,895	23,952	20,472
TR	832	1,514	3,502	7,193	16,825	19,479	19,453	18,288	17,270	25,798	19,136	19,160	19,061	19,770	17,682	18,177
OT	118	130	497	1,275	1,344	1,107	543	463	1,396	1,734	1,123	1,436	1,290	1,567	936	883
Total	27,978	53,096	54,017	126,193	335,634	387,728	337,988	361,772	463,615	529,719	497,254	417,096	315,989	324,664	275,955	299,074

Fisheries: Purse seine free-school (FS); Purse seine associated school (LS); Deep-freezing longline (LL); Fresh-tuna longline (LF); Pole-and-Line (BB); Gillnet (GI); Hand line (HD); Trolling (TR); Other gears nei (OT).

TABLE 5. Best scientific estimates of the catches of yellowfin tuna (*Thunnus albacares*) in the Western and Eastern Indian Ocean areas for the period 1950–2010 (in metric tons). Data as of October 2011.


Area*	By decade (average)						By year (last ten years)									
	1950s	1960s	1970s	1980s	1990s	2000s	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
R1	2,164	5,430	9,376	18,462	73,169	83,578	65,544	73,160	82,854	119,183	129,226	92,860	74,179	72,600	62,861	65,123
R2	11,899	23,101	20,921	72,400	143,122	183,679	156,045	164,369	265,456	278,103	248,113	204,035	126,450	135,499	100,973	111,041
R3	919	7,857	4,483	9,646	28,681	33,100	32,009	34,377	31,004	36,490	33,887	33,480	35,123	30,867	28,990	27,545
R4	918	1,799	1,370	1,075	3,314	2,122	3,376	3,328	2,387	3,802	2,904	1,363	540	507	427	498
R5	12,079	14,909	17,869	24,611	87,347	85,250	81,014	86,538	81,914	92,141	83,124	85,358	79,697	85,191	82,704	94,867
Total	27,978	53,096	54,017	126,193	335,634	387,728	337,988	361,772	463,615	529,719	497,254	417,096	315,989	324,664	275,955	299,074

*See Fig. 9 for a description of the areas

Uncertainty of catches

Retained catches are generally well known for the major fleets (Fig. 5); but are less certain for:

- Many coastal fisheries, notably those from Indonesia, Sri Lanka, Yemen, Madagascar and Comoros.
- The gillnet fishery of Pakistan.
- Non-reporting industrial purse seiners and longliners (NEI), and commercial longliners from India.



- the catch series for yellowfin tuna has not been significantly revised since the WPTT12 in 2010, although there has been some revision to the time series of catch from the fisheries of India leading to changes in catches by gear.
- levels of discards are believed to be low although they are unknown for most industrial fisheries, excluding industrial purse seiners flagged in EU countries for the period 2003–2007.
- catch-and-effort series are available from the major industrial and artisanal fisheries. However, these data are not available for some important artisanal fisheries or they are considered to be of poor quality for the following reasons:
 - no data are available for the fresh-tuna longline fishery of Indonesia, over the entire time series, and very little data available for the fresh-tuna longline fishery of Taiwan,China.
 - no data are available for the gillnet fisheries of Pakistan.
 - although Iran has provided catch and effort data, it is not reported as per the IOTC standards.
 - the poor quality effort data for the significant gillnet/longline fishery of Sri Lanka.
 - no data are available from important coastal fisheries using hand and/or troll lines, in particular Yemen, Indonesia, Madagascar and Comoros.

Effort trends

Total effort from longline vessels flagged to Japan, Taiwan,China and EU,Spain by five degree square grid from 2007 to 2010 are provided in Fig. 6, and total effort from purse seine vessels flagged to the EU and Seychelles (operating under flags of EU countries, Seychelles and other flags), and others, by five degree square grid and main fleets, for the years 2007 to 2010 are provided in Fig. 7. The total number of fishing trips by vessels flagged to the Maldives by 5 degree square grid, type of boat and gear, for the years 2009 and 2010 are provided in Fig. 8.




Fig. 6. Number of hooks set (millions) from longline vessels by five degree square grid and main fleets, for the years 2009 (left) and 2010 (right) (Data as of August 2011).

- LLJP (light green): deep-freezing longliners from Japan
- LLTW (dark green): deep-freezing longliners from Taiwan,China
- SWLL (turquoise): swordfish longliners (Australia, EU, Mauritius, Seychelles and other fleets)
- FTLL (red) : fresh-tuna longliners (China, Taiwan,China and other fleets)
- OTLL (blue): Longliners from other fleets (includes Belize, China, Philippines, Seychelles, South Africa, Rep. of Korea and various other fleets)




Fig. 7. Number of hours of fishing (Fhours) from purse seine vessels by 5 degree square grid and main fleets, for the years 2009 (left) and 2010 (right) (Data as of August 2011).

PS-EU (red): Industrial purse seiners monitored by the EU and Seychelles (operating under flags of EU countries, Seychelles and other flags)
 PS-OTHER (green): Industrial purse seiners from other fleets (includes Japan, Mauritius and purse seiners of Soviet origin) (excludes effort data for purse seiners of Iran and Thailand)




Fig. 8. Number of fishing trips by vessels flagged to the Maldives by 5 degree square grid, type of boat and gear, for the years 2009 (left) and 2010 (right) (Data as of August 2011).

BBN (blue): Baitboat non-mechanized; BBM (Green): Baitboat mechanized; BB (Red): Baitboat unspecified; UN (Purple): Unclassified gears
 Note that the above maps were derived using the available catch-and-effort data in the IOTC database, which is limited to the number of baitboat calls (trips) by atoll by month for Maldivian baitboats for the period concerned. Note that some trips may be fully devoted to handling, trolling, or other activities (data by gear type are not available since 2002). No data are available for the pole-and-line fisheries of India (Lakshadweep) and Indonesia.

Standardised catch-per-unit-effort (CPUE) trends

For the longline fisheries (LL fisheries in regions 1–5; Fig. 9), CPUE indices were derived using generalized linear models (GLM) from the Japanese longline fleet (LL regions 2–5) and for the Taiwanese longline fleet (LL region 1) to be used in the stock assessment. Standardised longline CPUE indices for the Taiwanese fleet were available for 1979–2008. The GLM analysis used to standardise the Japanese longline CPUE indices was refined for the 2011 assessment to include a spatial (latitude*longitude) variable. The resulting CPUE indices were generally comparable to the indices derived from the previous model and were adopted as the principal CPUE indices for the 2011 assessment (Fig. 10). There is considerable uncertainty associated with the Japanese CPUE indices for region 2 in the most recent year (2010) and no CPUE indices are available for region 1 for 2009–10.




Fig. 9. Spatial stratification of the Indian Ocean for the MFCL assessment model.




Fig. 10. Annualised GLM standardised catch-per-unit-effort (CPUE) for the principal longline fisheries (longline region 1: Taiwan, China and longline regions 2–5: Japan) and the whole Indian Ocean (IO), scaled by the respective region scalars.

Fish size or age trends (e.g. by length, weight, sex and/or maturity)

- trends in average weight (Fig. 11) can be assessed for several industrial fisheries but they are very incomplete or of poor quality for some fisheries, namely hand lines (Yemen, Comoros, Madagascar), troll lines (Indonesia) and many gillnet fisheries (see paper IOTC–2011–WPTT13–08).




Fig. 11. Changes in average weight (kg) of yellowfin tuna from 1950 to 2010 – all fisheries combined (Data as of September 2011).

- catch-at-Size and Age tables are available although the estimates are more uncertain in some years and some fisheries due to:
 - size data not being available from important fisheries, notably Yemen, Pakistan, Sri Lanka and Indonesia (lines and gillnets) and Comoros and Madagascar (lines).
 - the paucity of size data available from industrial longliners from the late-1960s up to the mid-1980s.
 - the paucity of catch by area data available for some industrial fleets (NEI, Iran, India, Indonesia, Malaysia).

Tagging data

A total of 63,310 yellowfin tuna were tagged during the Indian Ocean Tuna Tagging Programme (IOTTP) which represented 31.4% of the total number of fish tagged. Most of the yellowfin tuna tagged (86.4%) were tagged during the main Regional Tuna Tagging Project-Indian Ocean (RTTP-IO) and were primarily released off the coasts of the Seychelles, in the Mozambique Channel, along the coast of Oman and off the coast of Tanzania (Fig. 12) between May 2005 and September 2007. The remaining were tagged during small-scale projects around the Maldives, India and the southwest and eastern Indian Ocean by institutions with the support of IOTC. To date 10,560 (16.7%) tagged fish have been recovered and reported to the IOTC Secretariat.




Fig. 12. Densities of releases (in red) and recoveries (in blue) of yellowfin tuna. Data as of September 2011.

STOCK ASSESSMENT

A single quantitative modelling method (MULTIFAN-CL) was applied to the yellowfin tuna assessment in 2011, using data from 1972–2010. The following is worth noting with respect to the modelling approach used:

- The main features of the model in the 2010 assessment included a fixed growth curve (with variance) with an inflection, an age-specific natural mortality rate profile (M), the modelling of 24 fisheries including the separation of two purse seine fisheries into three time blocks, using a cubic spline method to estimate longline selectivities in the place of a logistic curve, the down-weighting of length frequency data in the fitting, separation of the analysis into five regions of the Indian Ocean and the specification of four steepness parameters for the stock recruitment relationship ($h=0.6, 0.7, 0.8$ and 0.9).
- In addition to another year of data, the 2011 assessment included several changes to the previous assessment: the longline CPUE indices were modified (Japanese updated with latest year which included information about latitude and longitude in the standardisation process for Regions 2–5 was supplied and the Taiwan,China index was revised for region 1); major historical catch revisions for fisheries in Region 5, splitting the longline fleet in Region 5 into distant water and fresh tuna logline fleets leaving 25 total fleets in the model; and the range of steepness evaluated was expanded to $h=0.55-0.95$.

While the biomass trends were very similar between the 2010 and 2011 assessments, the estimates of stock productivity and thus, the status, differed. There were several reasons for this: there was poor convergence in the 2010 assessment, thus the fits were suboptimal and alternative solutions were near optimal. Refitting the 2010 assessment is now more optimistic. Also, fitting the 2010 model to 2011 data was more optimistic. Thus, revisiting of key parameters and the inclusion of the latest year of data in the 2011 assessment appeared to be important. These issues are difficult to explore in the MFCL framework. The WPTT reviewed several alternative model structures and parameter formulations for the model that were presented in the assessment. These included: the new longline model structure for Region 5; alternative Japanese CPUE indices; a single region model where all 5 Regions were collapsed into one; a Region 2 model estimated separately from other Regions; the 5 values of steepness and alternative tag mixing periods (1–4 quarters). Additionally, an attempt was made to estimate age-specific M's. In regards to the latter, this parameter was not well estimated and the WPTT adopted the low M profile as the most appropriate way to proceed.

The problems identified in the catch data from some fisheries, and especially on the length frequencies in the catches of various fleets, a very important source of information for stock assessments. Length frequency data is almost unavailable for some fleets, while in other cases sample sizes are too low to reliably document changes in abundance and selectivity by age. Moreover, in general, catch data from some coastal fisheries is considered as poor.

The available tagging data has provided the WPTT with relevant information on various biological parameters, such as natural mortality and growth. Further use of these data should better support the analyses conducted by the WPTT.

In the previous assessment purse seine selectivity in the period 2003–2007 was separated into three blocks of time surrounding 2005 to accommodate the unusually large catches in the middle of that time period. This was continued in the current assessment. However, the WPTT questioned whether this was the most appropriate way to do this. An alternative was suggested in which the time blocks of PS fleet were removed and the same selectivity was applied throughout the period. This was explored in new model runs. Results were not demonstrably different.

Longline selectivity will be revisited in 2012 as it was suggested that this selectivity might still be best described by a logistic (flat-topped) model instead of a cubic spline approach, whereby the resulting selectivity was dome-shaped. This option reinvigorated a long standing debate that has yet to be resolved. A run whereby logistic selectivities were imposed was evaluated.

Generally, the runs with alternative parameter and model structures did not suggest large differences in the approach and resulted in qualitatively predictable outcomes. The WPTT felt that the alternative outcomes were an expression of uncertainties in the model, data and assessment. Therefore, the WPTT focused on following basic alternatives for characterizing the uncertainty: logistic versus cubic spline longline selectivity; using the low M profile; alternative steepness of the stock-recruitment relationship of 0.7, 0.8 and 0.9, and estimation of MSY based reference points using the average recruitment for the whole time series. It was determined that with current knowledge outcomes using these alternatives are equally likely and a combined evaluated was generated based upon this.

The final range of model options adopted by the WPTT included the 2 alternative parametrization of longline selectivity (cubic spline and logistic) and three steepness options (0.7, 0.8 and 0.9). For the cubic spline model option, there is a strong temporal trend in recruitment and recent recruitments (average of the last 15 years) is estimated to be lower (80%) than the long term recruitment level. On that basis, it was agreed to also derived alternative MSY estimates based on the recent levels of recruitment for comparative purposes. Key assessment results for the MFCL stock assessment are shown in Tables 1, 2 and 6; Fig. 1.

Table 6. Key management quantities from the MFCL assessment, for the agreed scenarios of yellowfin tuna in the Indian Ocean. Values represent an equal weighting mean of the scenarios investigated. The range is described by the range values between those scenarios.

Management Quantity	Indian Ocean
2010 catch estimate (1000 t)	299.1
Mean catch from 2006–2010 (1000 t)	326.6
MSY (1000 t)	357 (290–435)
Data period used in assessment	1972–2010
F_{2009}/F_{MSY}	0.84 (0.63–1.10)
B_{2009}/B_{MSY}	1.46 (1.35–1.59)
SB_{2009}/SB_{MSY}	1.61 (1.47–1.78)
B_{2009}/B_0	0.49
SB_{2009}/SB_0	0.35 (0.31–0.38)
$B_{2009}/B_{0, F=0}$	0.58
$SB_{2009}/SB_{0, F=0}$	–

LITERATURE CITED

Froese R, & Pauly DE 2009. *FishBase*, version 02/2009, FishBase Consortium, <www.fishbase.org>.