

**EXECUTIVE SUMMARY: STATUS OF THE INDIAN OCEAN NARROW-BARRED SPANISH MACKEREL
(*SCOMBEROMORUS COMMERSON*) RESOURCE**

TABLE 1. Status of narrow-barred Spanish mackerel (*Scomberomorus commerson*) in the Indian Ocean.

Area ¹	Indicators – 2011 assessment		2011 stock status determination
			2010 ²
Indian Ocean	Catch ³ 2010: 124,107 t Average catch ³ 2006–2010: 116,444 t MSY: unknown F ₂₀₁₀ /F _{MSY} : unknown SB ₂₀₁₀ /SB _{MSY} : unknown SB ₂₀₁₀ /SB ₀ : unknown		UNCERTAIN

¹Boundaries for the Indian Ocean stock assessment are defined as the IOTC area of competence.

²The stock status refers to the most recent years' data used for the assessment.

³Nominal catches represent those estimated by the IOTC Secretariat. If these data are not reported by CPCs, the IOTC Secretariat estimates total catch from a range of sources including: partial catch and effort data; data in the FAO FishStat database; catches estimated by the IOTC from data collected through port sampling; data published through web pages or other means; data reported by other parties on the activity of vessels; and data collected through sampling at the landing place or at sea by scientific observers.

Colour key	Stock overfished (SB _{year} /SB _{MSY} < 1)	Stock not overfished (SB _{year} /SB _{MSY} ≥ 1)
Stock subject to overfishing (F _{year} /F _{MSY} > 1)		
Stock not subject to overfishing (F _{year} /F _{MSY} ≤ 1)		

INDIAN OCEAN STOCK – MANAGEMENT ADVICE

The WPNT **RECOMMENDED** the following management advice for narrow-barred Spanish mackerel in the Indian Ocean, for the consideration of the Scientific Committee, noting that there remains considerable uncertainty about stock structure and about the total catches.

Stock status. No quantitative stock assessment is currently available for narrow-barred Spanish mackerel in the Indian Ocean, and due to a lack of fishery data for several gears, only preliminary stock indicators can be used. Therefore stock status remains *uncertain* (Table 1). However, aspects of the fisheries for this species combined with the lack of data on which to base a more formal assessment are a cause for considerable concern. Although indicators from the Gulf and Oman Sea suggest that overfishing is occurring in this area, the degree of connectivity with other regions remains unknown.

Outlook. The continued increase of annual catches for narrow-barred Spanish mackerel in recent years has further increased the pressure on the Indian Ocean stock as a whole, however there is not sufficient information to evaluate the effect this will have on the resource. The apparent fidelity of narrow-barred Spanish mackerel to particular areas/regions is a matter for concern as overfishing in these areas can lead to localised depletion. Research emphasis on improving indicators and exploration of stock structure and stock assessment approaches for data poor fisheries are warranted.

The WPNT **RECOMMENDED** that the Scientific Committee consider the following:

- the Maximum Sustainable Yield estimate for the whole Indian Ocean is unknown.
- annual catches urgently need to be reviewed.
- improvement in data collection and reporting is required to assess the stock.

SUPPORTING INFORMATION

(Information collated from reports of the Working Party on Neritic Tunas and other sources as cited)

CONSERVATION AND MANAGEMENT MEASURES

Narrow-barred Spanish mackerel (*Scomberomorus commerson*) in the Indian Ocean is currently subject to a number of conservation and management measures adopted by the Commission, although none are species specific:

- Resolution 08/04 concerning the recording of catch by longline fishing vessels in the IOTC area.
- Resolution 09/02 On the implementation of a limitation of fishing capacity of contracting parties and cooperating non-contracting parties.
- Resolution 10/02 mandatory statistical requirements for IOTC Members and Cooperating non-Contracting Parties (CPC's).

- Resolution 10/03 concerning the recording of catch by fishing vessels in the IOTC area.
- Resolution 10/08 concerning a record of active vessels fishing for tunas and swordfish in the IOTC area.
- Recommendation 11/06 Concerning the Recording of Catch by Fishing Vessels in the IOTC Area of Competence.

FISHERIES INDICATORS

General

The narrow-barred Spanish mackerel (*Scomberomorus commerson*) is a pelagic, top level predator found throughout tropical marine waters of the Indo-West Pacific. Table 2 outlines some key life history parameters relevant for management.

TABLE 2. Biology of Indian Ocean narrow-barred Spanish mackerel (*Scomberomorus commerson*).

Parameter	Description
Range and stock structure	A pelagic, top level predator found throughout tropical marine waters of the Indo-West Pacific. Juveniles inhabit shallow inshore areas whereas adults are found in coastal waters out to the continental shelf. Adults are usually found in small schools but often aggregate at particular locations on reefs and shoals to feed and spawn. Appear to undertake lengthy migrations. Feed primarily on small fishes such as anchovies, clupeids, carangids, also squids and shrimps. Genetic studies carried out on <i>S. commerson</i> from Djibouti, Oman and U.A.E. showed there were small genetic differences among stocks in these three places.
Longevity	~16 years
Maturity (50%)	Age: n.a.; females n.a. males n.a. Size: females ~81 cm FL and males ~52 cm FL.
Spawning season	Females are multiple spawners. Year-round spawning has been observed in east African waters, with peaks during late spring to summer (April-July) and autumn (September-November) coinciding with the two seasonal monsoons which generate high abundances of plankton and small pelagic fish.
Size (length and weight)	Maximum: Females and males 240 cm FL; weight 70 kgs.

n.a. = not available. SOURCES: Grandcourt et al. (2005); Froese & Pauly (2009); Darvishi et al. (2011)

Narrow-barred Spanish mackerel – Catch trends

Narrow-barred Spanish mackerel is targeted throughout the Indian Ocean by artisanal and recreational fishers. The main method of capture is gillnet, but significant numbers of are also caught trolling (Fig. 1).

The catch estimates for narrow-barred Spanish mackerel were derived from very small amounts of information and are therefore highly uncertain. The catches of narrow-barred Spanish mackerel increased from around 50,000 t the mid-1970's to over 100,000 t by the mid-1990's. The highest catches of Spanish mackerel were recorded in 2010, amounting to 124,107 t. In recent years, catches have been increasing, with average annual catches for 2006–2010 estimated to be around 116,444 t (Table 3). Narrow-barred Spanish mackerel is caught in both Indian Ocean basins, with higher catches recorded in the West.

In recent years, the countries attributed with the highest catches of Spanish mackerel are India (29%) and Indonesia (23%) and, to a lesser extent, Iran, Pakistan, and Madagascar (20%) (Fig. 2).





TABLE 3. Best scientific estimates of the catches of narrow-barred Spanish mackerel by type of fishery for the period 1950–2010 (in metric tonnes). Data as of October 2011.

Fishery	By decade (average)						By year (last ten years)									
	1950s	1960s	1970s	1980s	1990s	2000s	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Purse seine	0	0	237	1,141	2,571	1,782	1,404	1,928	2,325	1,590	2,116	3,926	1,877	1,951	1,920	2,874
Gillnet	7,164	15,184	26,883	54,952	71,418	78,404	78,408	73,231	76,410	73,571	64,618	74,173	77,371	84,124	84,225	89,352
Line	2,330	3,350	6,529	13,733	14,964	16,823	16,773	15,420	17,023	15,214	16,145	17,137	15,811	17,394	18,099	18,045
Other	1,368	2,012	4,255	6,635	10,616	13,932	13,264	15,354	14,566	12,996	13,537	16,239	15,547	14,793	13,527	13,836
Total	10,862	20,546	37,904	76,462	99,570	110,941	109,849	105,933	110,324	103,370	96,416	111,475	110,605	118,262	117,770	124,107

Narrow-barred Spanish mackerel – uncertainty of catches

Retained catches are uncertain (Fig. 3), notably for the following fisheries:

- Artisanal fisheries of India and Indonesia: India and Indonesia have only recently reported catches of narrow-barred Spanish mackerel by gear, including catches by gear for the years 2005–2008 and 2007–2008, respectively. In both cases, the IOTC Secretariat used the catches reported by gear to break previous catches of this species by gear. The catches of narrow-barred Spanish mackerel estimated for this component represent more than 52% of the total catches of this species in recent years.
- Artisanal fisheries of Madagascar: Madagascar has never reported catches of narrow-barred Spanish mackerel to the IOTC Secretariat. During 2010 the IOTC Secretariat conducted a review aiming to break the catches recorded in the FAO database as narrow-barred Spanish mackerel by species, on the assumption that all catches of neritic tunas had been combined under this name. The new catches estimated are thought to be very uncertain.
- Artisanal fisheries of Mozambique, Myanmar and Somalia: None of these countries have ever reported catches to the IOTC Secretariat. Catch levels are unknown.
- Other artisanal fisheries: Oman and the United Arab Emirates do not report catches of narrow-barred Spanish mackerel by gear. Although most of the catches are believed to be taken by gillnets, some fish may be also caught by using small surrounding nets, lines or other artisanal gears. Thailand and Malaysia report catches of narrow-barred Spanish mackerel and Indo-Pacific king mackerel aggregated.
- All fisheries: In some cases the catches of seerfish species are mislabelled, the catches of Indo-Pacific king mackerel and, to a lesser extent, other seerfish species, labelled as narrow-barred Spanish mackerel. Similarly, the catches of wahoo in some longline fisheries are thought to be mislabelled as narrow-barred Spanish mackerel. This mislabelling is thought to have little impact in the case of the narrow-barred Spanish mackerel but may be important for other seerfish species.
- Discard levels are believed to be low although they are unknown for most fisheries.
- Changes to the catch series: The catch series of narrow-barred Spanish mackerel has changed since those estimated in 2010, following reviews of catches for the coastal fisheries in Indonesia and India, involving marked changes in catches by species. Overall, the new catches estimated represent the 98% of those recorded in the past.




Fig. 3. Narrow-barred Spanish mackerel: Uncertainty of annual catch estimates (1960–2010) (Data as of November 2011).


Catches below the zero-line (Type B) refer to fleets that do not report catch data to the IOTC (estimated by the IOTC Secretariat), do not report catch data by gear and/or species (broken by gear and species by the IOTC Secretariat) or any of the other reasons provided in the document. Catches over the zero-line (Type A) refer to fleets for which no major inconsistencies have been found to exist. Light bars represent data for artisanal fleets and dark bars represent data for industrial fleets.

Narrow-barred Spanish mackerel – Effort trends

Effort trends are unknown for narrow-barred Spanish mackerel in the Indian Ocean.

Narrow-barred Spanish mackerel – Catch-per-unit-effort (CPUE) trends

Standardised CPUE series have not yet been developed. Nominal CPUE series are however available from some fisheries but they are considered highly incomplete. In most cases catch-and-effort data are only available for short periods. Reasonably long catch-and-effort data series (extending for more than 10 years) are only available for Sri Lanka gillnets (Fig. 4). The catches and effort recorded are, however, thought to be unrealistic due to the dramatic changes in CPUE recorded in 2003 and 2004.



Narrow-barred Spanish mackerel – Fish size or age trends (e.g. by length, weight, sex and/or maturity)

- The size of narrow-barred Spanish mackerel taken by the Indian Ocean fisheries typically ranges between 30–140 cm depending on the type of gear used, season and location. The size of narrow-barred Spanish mackerel taken varies by location with 32–119 cm fish taken in the Eastern Peninsular Malaysia area, 17–39 cm fish taken in the East Malaysia area and 50–90 cm fish taken in the Gulf of Thailand. Similarly, Spanish mackerel caught in the Oman Sea are typically larger than those caught in the Persian Gulf.
- Trends in average weight can only be assessed for Sri Lankan gillnets but the amount of specimens measured has been very low in recent years. The length frequency data available from the mid-eighties to the early nineties was obtained with the support of the IPTP (Indo-Pacific Tuna Programme). Unfortunately, data collection did not continue after the IPTP activities came to an end.
- Catch-at-Size(Age) tables are not available for narrow-barred Spanish mackerel due to the paucity of size data available from most fleets and the uncertain status of the catches for this species.
- Sex ratio data have not been provided to the Secretariat by CPCs.

STOCK ASSESSMENT

No quantitative stock assessment for narrow-barred Spanish mackerel in the Indian Ocean is known to exist and no such assessment has been undertaken by the IOTC Working Party on Neritic Tunas. However, a preliminary estimation of stock indicators was attempted on the catch and effort datasets from the Sri Lankan gillnet fishery (described above). However, there is considerable uncertainty about the degree to which this and other indicators represent abundance as factors such as changes in targeting practices, discarding practices, fishing grounds and management practices are likely to interact in the depicted trends. Further work must be undertaken to derive additional stock indicators for this species, because in the absence of a quantitative stock assessment, such indicators represent the only means to monitor the status of the stock and assess the impacts of fishing.

TABLE 4. Narrow-barred Spanish mackerel (*Scomberomorus commerson*) stock status summary.

Management Quantity	Aggregate Indian Ocean
2010 catch estimate (1000 t)	124.1
Mean catch from 2006–2010 (1000 t)	116.4
MSY (1000 t) (80% CI)	unknown
Data period used in assessment	–

F_{2010}/F_{MSY} (80% CI)	–
B_{2010}/B_{MSY} (80% CI)	–
SB_{2010}/SB_{MSY}	–
B_{2010}/B_0 (80% CI)	–
SB_{2010}/SB_0	–
$B_{2010}/B_{0, F=0}$	–
$SB_{2010}/SB_{0, F=0}$	–

LITERATURE CITED

- Darvishi M, Kaymaram F, Salarpouri A, Behzadi S and Daghooghi B, 2011. Population dynamic and biological aspects of *Scomberomorus commerson* in the Persian Gulf and Oman Sea (Iranian coastal). IOTC–2011–WPNT01–23. Working paper.
- Froese R & Pauly DE, 2009. FishBase, version 02/2009, FishBase Consortium, <www.fishbase.org>.
- Grandcourt EM, Al Abdessalaam TZ, Francis F and Al Shamsi AT, 2005. Preliminary assessment of the biology and fishery for the narrow-barred Spanish mackerel, *Scomberomorus commerson* (Lacépède, 1800), in the southern Arabian Gulf. Fish. Res.76:277–290.