

Preparation of data input files for the assessments of Indian Ocean yellowfin tuna stock

Miguel Herrera¹, Julien Million²
(IOTC Secretariat)

Summary

This document describes the methods used by the IOTC Secretariat to prepare catch, effort tagging data and length-frequency tables for the assessments of the Indian Ocean yellowfin tuna stock, for the period 1950-2010, using estimates of total catch and the available catch-and-effort, size frequency data and other biological data in the IOTC database.

Using information from the IOTC database, the IOTC Secretariat prepared a flat table containing estimates of total catches, and the effort and length-frequency samples available, by fishery, assessment area, year and quarter.

The results are affected by the lack of information for some fleets, areas and years, and, in particular, by the lack of catch, effort and size data from most artisanal fleets and some industrial fleets.

Rationale

The IOTC database contains estimates of total catches by country, gear, year and IOTC Area (**Figure 1**). In addition, the IOTC database contains catch-and-effort data and size frequency data by country, gear, time-area strata and species, which generally represent a sample of the total catches estimated by country, gear, year and species.

The Secretariat used the above data to produce an input file containing estimates of total catches of yellowfin tuna, in number and weight, and the effort and length-frequency samples available by fishery, assessment area, year and quarter. This file was used for the assessments of yellowfin tuna, carried out using Multifan-CL (MF-CL), Stock Synthesis III (SS3), or other assessment models. The results of the assessments are reported in separate papers³.

The IOTC is hosting the database developed under the framework of the Regional Tuna Tagging Project in the Indian Ocean (RTTP-IO). This database contains all the information on releases and recoveries of yellowfin tuna, bigeye tuna and skipjack tuna tagged by the RTTP-IO. Tagging information from other tagging projects (*e.g.* past project in Maldives, and small-scale project of the Indian Ocean Tuna Tagging Programme) is also available but was not used for the assessments. The Secretariat prepared a file containing releases and recaptures of yellowfin tuna, to be used for the assessments.

Input Table

The Secretariat prepared a table containing the following information for the assessments of yellowfin tuna:

- a. **Total catches** of yellowfin tuna, in number of specimens and weight, by year, quarter, assessment fishery (**Table 6**), and assessment area (**Table 7** and **Figure 2**).
- b. **Effort** data by year, quarter, assessment fishery, and assessment area, including:
 - a. Industrial purse seine and longline fleets: Estimates of total effort by stratum (*i.e.* fishery, area, year and quarter) in strata for which effort data were available (no effort was estimated in strata for which no effort data were available at all)
 - b. Other fleets: The effort data were used as available.
- c. Number of yellowfin tuna specimens **sampled** by **length** interval, year, quarter, assessment fishery, and assessment area.
- d. Number of yellowfin tuna **tagged and release** by length, region of release and quarter (*i.e.* release event), and for each release event, the number of recovered yellowfin by length at release and by fishery of recovery (tagging data).

Examples of the input table can be found in **Appendix I**.

¹ Data Coordinator, IOTC Secretariat (mh@iotc.org; Miguel.Herrera@iotc.org)


² Fishery Officer, IOTC Secretariat (jm@iotc.org; Julien.Million@iotc.org)

³ Preliminary stock assessment of yellowfin tuna in the Indian Ocean using MULTIFAN-CL, A.Langley *et al.* (IOTC-2011-WPTT13-36) and Stock assessment by Stock Synthesis III (SS3) for yellowfin tuna in the Indian Ocean, T. Kitakado *et al.* (IOTC-2011-WPTT13-37)

Basic Data

Five datasets are used for the preparation of the input files for the assessments of yellowfin tuna:

- **Nominal catches:** Total catch estimates per Species, Fleet, Year, Gear and IOTC Area (**Figure 1**). The data in this dataset issues from two different sources:
 - a. Reports from the flag countries or reports from other countries on the catches of foreign vessels operating within its Economic Exclusive Zone or based in ports within its territory.
 - b. Estimates carried out by the IOTC Secretariat: this may involve changes in the catches reported by the above or the estimation of catches for non-reporting fleets (e.g. catches recorded under the NEI⁴ category), using various sources.
- **Catch-and-effort by area** (derived from the catch-and-effort table): Catches by species (in tonnes or/and in number of fish) and effort are recorded by Fleet, Year, Gear, Fishing Mode, Time Interval (month or quarter usually) and area (usually 1^o square areas for industrial purse seine fisheries, 5^o square areas for industrial longline fisheries and various regular or irregular areas for artisanal fisheries). Catches/effort by area are not available for all Nominal catches strata. When recorded, the catches/effort in these datasets might represent the total catches/effort of the species in the year for the fleet and gear concerned or represent simply a sample of those.
- **Size data:** Size frequency data (standard or processed lengths or standard or processed weights) are recorded by Species, Fleet, Year, Gear, Fishing Mode, Time Interval (month or quarter or year usually) and area (usually 5^o square areas for purse seine fisheries, 10^o latitude by 20^o longitude for longline fisheries and various regular or irregular areas for artisanal fisheries). Size data are not available for all Nominal catches strata. When recorded, the size data might represent the total catches of the species in the strata concerned (Catch-at-Size) or simply a sample of those (non-raised or partially raised samples).
- **Biological data:** includes several types of biological parameters for the yellowfin tuna, in particular:
 - a. **Conversion from non-standard measurements into fork length:** Equations (data) used to convert specimens of yellowfin tuna measured by using non-standard procedures into the standard length measurement used for these species, representing the distance from the tip of the snout to the fork of the tail (fork length).
 - b. **Conversion from fork length into live weight:** Equations (data) used to estimate sample weights from the available lengths (length-weight relationships).
- **Tagging data:**
 - a. **Release data:** number of release at length, quarter and region of release. In total during the RTTP-IO, 54,688 yellowfin were released in the western Indian Ocean.
 - b. **Recovery data:** 9,612 tagged yellowfin were recovered and reported as of December 2010 (?). Most of the recoveries are reported from the purse-seine fleet at the moment of the unloading of the fishing vessel in the ports of the Indian Ocean, in particular Seychelles. For all these recoveries, the information collected contains no specific date and location of recapture but the date of recovery (*i.e.* unloading) and the well number in which the tagged fish was recovered. This information is then linked with the logbook database of the fleet in order to extract the need information about the location, date of recapture and type of association.
 - c. **Recovery data:**



⁴ Not elsewhere identified

The type of information recorded in each case is summarized in **Table 1** below:

Table 1: Main types of fisheries statistics gathered by the IOTC				
<i>Dataset</i>	<i>Fishery Strata</i>	<i>Time Period Strata</i>	<i>Area Strata</i>	<i>Represents</i>
Nominal Catches	Fleet-Gear type (or gear aggregate)-Species (or species aggregate)	Year	IOTC Area (Western / Eastern Indian Ocean)	Total catches
Catch-and-Effort	Fleet-Gear (or gear aggregate)-Fishing Mode (purse seine only)-Species	Month (quarter or year)	1° square area (purse seine) 5° square area (longline) Other regular or irregular areas	Sample
Size frequency	Species- Fleet-Gear (or gear aggregate)-Fishing Mode (purse seine only)-Type of measurement (length or weight, standard or processed)-Size interval (between size classes)	Month (quarter or year)	5° square area (purse seine) 10° Lat.x20° Lon. area (longline) Other regular or irregular areas	Sample
Biological data	Various, depending on dataset	Various	Various, depending on dataset	Sample
Release data	Species, length, gear	Date of release	Position of tagging	Total
Recovery data	Species, length or weight, gear	Date of recovery (or estimation)	Position of recovery (or estimation)	Total

Data Processing

Summary of the estimation procedures used in the preparation of input files for the assessments of yellowfin tuna

Catch, effort and length frequency samples file:

1. Standardizing catch, effort, and size frequency tables
 - a. Nominal catches (NC): Assigning the catches not reported by species/gear by species/gear (NC→NCst)
 - b. Catch-and-effort (CE): Assigning catches/effort not recorded by quarter by quarter (CE→CEst)
 - c. Size frequency (SF→LFst):
 - i. Converting non-standard measurements into standard measurements
 - ii. Breaking the existing lengths into the standard length class intervals used for the species
 - iii. Scaling down length frequency distributions to sampled numbers
 1. EC purse seiners: scaling down CAS to the numbers of fish estimated for sampled purse seine sets
 2. Other fleets: scaling down CAS or raised length distributions to the actual numbers of fish sampled, where required
 - iv. Assigning length samples by quarter
2. Assigning catches, effort and length samples by year, quarter, fishery and assessment area (NCst/CEst/LFst → NCELfa):
 - a. Fleets operating fully within one of the yellowfin tuna assessment areas (artisanal fleets): Assigning total catches (NCst) and available effort (CEst) and length samples (LFst) by fishery, assessment area, year and quarter (NCst/CEst/LFst → NCELfa₁)
 - b. Fleets operating in two or more assessment areas (industrial fleets):
 - i. Breaking CEst by fishery, 5° grid, year and quarter (CEst→CEds)
 - ii. Breaking NCst by fishery, 5° grid, year and quarter using the CEds (NCst→NCds)
 - iii. Assigning LFst by fishery, 5° grid, year and quarter (LFst→LFds)
 - iv. Assigning total catches (NCds) by fishery, assessment area, year and quarter (NCds → NCELfa₂)
 - v. Estimating total effort (CEds) by fishery, assessment area, year and quarter for industrial purse seine and longline fleets (CEds → NCELfa₂) in strata where effort data are available
 - vi. Assigning the available length frequency samples (LFds) by fishery, assessment area, year and quarter for other fleets (LFds → NCELfa₂)
 - c. Merging NCELfa₁ and NCELfa₂ into NCELfa

3. **Yellowfin tuna MF-CL input file** (NCELfa → NCEL_{MFCL}): re-allocating strata from NCELfa into other assessment areas. The catches (in weight⁵), effort, and length data for some fisheries were reallocated to specific areas when the accumulated catches in the area concerned represented a small proportion of the total catches for the fishery concerned.

Release and recapture data file:

1. Assigning releases by length, year, quarter and assessment area (*i.e.* release event)
2. Assigning recaptures by release event, year and quarter of recapture, fishery and assessment area
 - a. Assigning date and location to the recoveries on-board purse-seiners:
 - i. Linking date of recovery (*i.e.* unloading) and well number with the logbook database
 - ii. Assigning possible sets from which the recapture might come from
3. Deriving reference catches: Deriving catches of yellowfin tuna for vessel trips sampled for tags, by gear, fishing mode, and quarter.

Standardizing catch, effort and size frequency tables

i. Estimating total catches of yellowfin tuna by fleet, gear, year and IOTC Area

The catches in the IOTC nominal catches database are not recorded by species and/or by gear in all cases. The Secretariat conducted a review aiming at estimating catches when data were not available by species or gear in the IOTC database. This process was documented in a paper presented to the WPTT in 2004 (IOTC-2004-WPTT-06). The disaggregation of catches does not affect significantly the catches of yellowfin tuna.

ii. Standardizing the available catch-and-effort data

The catches in the catch-and-effort table are recorded under different levels of aggregation. All the catches from this record were assigned by Species-Fleet-Gear-Fishing Mode-Year-Month-5° square grid-Catch in number of fish-(and/or)-Catch in metric tons.

- i. Converting from non-standard effort units into standard effort units: The effort units reported vary depending on the fleet and/or fishing period. The units of effort that were selected by fishery are shown in **Table 2** (page 4). In some cases, especially for some artisanal fisheries, the units of effort reported were assumed to be equivalent to the standard units of effort selected for those fisheries.
- ii. It is important to note that the effort recorded for industrial purse seine fisheries has not been broken down by fishing mode (LS and FS) as the actual effort exerted on free schools or associated schools is unknown. Thus, **the effort recorded under FS and LS actually represents the effort exerted by industrial purse seiners as a whole, not the effort for the individual fisheries.**

Table 2: Standard length, first length, interval and total number of size classes used for the yellowfin tuna

<i>Fishery</i>	<i>Standard Effort Units</i>	<i>Other effort Units considered to be the same as the standard effort unit</i>
LS	Number of days fishing (FDAYS)	
FS	Number of days fishing (FDAYS)	
LL	Number of hooks set (HOOKS)	
LF	Number of hooks set (HOOKS)	
BB	Number of trips (TRIPS)	Number of fishing days (FDAYS), number of days at sea (DAYS)
GI	Number of trips (TRIPS)	Number of fishing days (FDAYS), number of days at sea (DAYS)
HD	Number of trips (TRIPS)	Number of fishing days (FDAYS), number of days at sea (DAYS)
TR	Number of trips (TRIPS)	Number of fishing days (FDAYS), number of days at sea (DAYS)
OT	Number of trips (TRIPS)	Number of fishing days (FDAYS), number of days at sea (DAYS)

NOTE: Refer to Table 6 for a description of the fisheries referred to above

- iii. Area allocation: All the catches/effort not recorded by 5° square areas were assigned to 5° square areas as follows:
 - a. Allocation of catches/effort recorded under irregular areas to regular grids: The catches/effort recorded under irregular areas (e.g. port of unloading, fishing district, etc.) were assigned to the neighbouring regular grids.
 - b. Allocation of catches/effort recorded under areas that fell within a single 5° square area: all catches/effort recorded under areas that fell within a 5° square area were assigned to the corresponding 5° square areas.
 - c. Allocation of catches/effort recorded under areas overlapping two or more 5° square areas: all catches/effort recorded under areas that overlapped two or more 5° square areas were assigned proportionally by 5° square areas (*i.e.* by using the proportions obtained by dividing the amount of 1 degree square grids that fell within each 5° square area over the total amount of squares from the overlapping area).

⁵ Note that while numbers of yellowfin tuna were used in previous assessments the present assessment uses catches in metric tonnes

- iv. **Time period allocation:** The catches/effort available in the catch-and-effort file were assigned by month as follows:
- Allocation of catches/effort recorded under time period strata that fall within a single month: all catches/effort recorded under time periods that fell within a month were assigned to the corresponding months.
 - Allocation of catches/effort recorded under time period strata overlapping two or more months: all catches/effort recorded under time periods that overlapped two or more months were assigned proportionally by month (e.g. 1/3 of the catches/effort recorded under the first quarter of a year were assigned to each of the months making up that quarter).

iii. **Standardizing the available size frequency data**

The following process was used to convert the samples available for the yellowfin tuna into standard form:

- Converting from non-standard measurement types into standard length (Table 3):** The regression equations presented in **Table 3** were used to estimate the distance from the tip of the snout to the fork of the tail (fork length) for specimens of yellowfin tuna that were recorded under non-standard lengths or weights in the IOTC database:
 - Conversions from gilled-and-gutted weight into fork length: The conversion from gilled-and-gutted weight into fork length was done by using a distribution matrix for the yellowfin tuna, estimated by using individual weight (GGT)-length (FL) measurements.
 - Other measurements: any other non-standard measurements were converted into fork length by using the equations on **Table 3** (deterministic conversion).

Type Measurement	Equation	Parameters	Sample size	Size range	Variance	Covariance ab	Mean Residual	Gradient
Weight gilled and gutted ^A	aW^b	a= 44.28699 b= 0.3008591	2,361	Min:14 Max:71	a=0.00752476509 b=2.86244E-07	-4.626246E-05	4.095958	a=3.033852 b=495.6385
Length to the base of the 1 st dorsal fin ^B	aL^b	a=2.0759 b=1.1513	7,036	Min: 29 Max: 164				
Length base of first dorsal fin to fork of caudal fin	No equation available							

A: Data from IPTP Penang Sampling Programme (1992-93)
B: Data from the Indian Ocean (Marsac, F. et al in IOTC-2006-WPTT-09)

- Breaking the samples by the standard length frequency intervals used for the yellowfin tuna:** The length-frequency interval that is used for the yellowfin tuna is shown in **Table 4** (page 5).
 - Allocation of specimens recorded under length classes that fall within a single standard length class:
 - Yellowfin tuna specimens recorded under one centimetre length classes were aggregated under the corresponding length classes (e.g. specimens of yellowfin tuna recorded under the classes 16-17cm and 17-18cm were accumulated under fork length 16cm⁶).
 - Yellowfin tuna specimens recorded under two centimetre length classes that fell within standard length classes were assigned to the corresponding standard length classes (e.g. specimens recorded under length class 16-18cm were assigned to fork length 16cm)

Species	Standard Length*	First length (cm)	Interval between length classes (cm)	Total number of size classes	Maximum interval allowed (cm)
Yellowfin tuna	Fork length	10	2	150	4

NOTE: All samples in the IOTC database were assigned according to the specifications above; the samples recorded under length intervals greater than the maximum interval specified above were not used
 *Refers to the straight distance measured, to the closest lower centimetre, between the tip of the snout (upper-jaw) and the fork of the tail (FL)

- Allocation of specimens recorded under length classes overlapping two or more standard length classes: all the specimens recorded under length classes that overlap the standard classes used for the species (Table 2) were assigned proportionally to the corresponding standard length classes (e.g. 1/3 of the yellowfin tuna specimens recorded under length class 17-20cm were assigned to length 18cm, and 2/3 to length 18cm). All samples containing specimens of yellowfin tuna recorded under length intervals 5cm or higher were discarded.

⁶ Note that lengths are recorded to the lowest length interval, e.g. the length range of specimens recorded under length 16cm for a species for which 2cm intervals are used is 16cm(inclusive)-18cm(exclusive); note that medium values are used to estimate weights from the available lengths (17cm in this case).

- iii. Weighting length frequency distributions to sample numbers: The length frequency data in the IOTC database do not represent sample numbers in all cases as some countries report length frequency data that has been raised in various ways (e.g. to the catches in the stratum covered through sampling, to the total catches estimated for the country, etc.).
- EC purse seiners: The EC reports length frequency distributions for EC purse seiners that have been raised to the total catches of each 5° square grid and month; the CAS for each stratum were weighted by the numbers of fish estimated for the sampling units making up the stratum (which is the catches corresponding to all sets within fish compartments where samples were taken).
 - The actual sample numbers were used for all other fleets to weight the reported length frequencies, when required; i.e. the number of specimens recorded under each length class was multiplied by the number obtained by dividing the total number of specimens sampled (all lengths combined) by the total number of specimens in the raised length frequency (all lengths combined).
- iv. Time period allocation: The available length frequency samples were assigned by quarter as follows:
- Allocation of specimens recorded under time-periods that fall within a single quarter: all specimens from samples recorded under time periods that fell within a quarter were assigned to the corresponding quarter.
 - Allocation of specimens recorded under time-periods overlapping two or more quarters: all specimens from samples recorded under time-periods that overlapped two or more quarters were assigned proportionally by quarter (e.g. 2/3 of the specimens recorded under the time period February-April of any year were assigned to the first quarter (Jan-Mar) of that year while the remaining 1/3 specimens were assigned to the second quarter (Apr-Jun)).
- v. Estimation of sample weights: Weights by sample were calculated by adding the weights estimated for all the specimens making it. The equations used to estimate weights from the available lengths are shown in **Table 5** (note that deterministic methods were used for the conversion).

Species	Gear Type/s	From type measurement – To type measurement	Equation	Parameters	Sample size	Length
Yellowfin tuna	Purse seine Pole and Line Gillnet & other	Fork length – Round Weight(kg) ^A	$RND=a*L^b$	a= 0.00001886 b= 3.0195	6,752	Min: 29 Max: 164
	Longline Hand line	Fork length(cm) – Gilled and gutted weight(kg) ^B Gilled and gutted weight(kg) - Round Weight(kg) ^C	$GGT=a*L^b$ $RND=GGT*1.13$	a= 0.0000094007 b= 3.126843987	15,133	Min:72 Max:177

A: Data from the Indian Ocean (Marsac, F. et al in IOTC-2006- WPTT-09)
 B: Multilateral catch monitoring Benoa (2002-04)
 C: ICCAT Field Manual (Appendix 4: Population parameters for key ICCAT species. Product Conversion Factors)

Assigning catches, effort and length samples by year, quarter, fishery and assessment area

i. Allocation of assessment fishery:

Each Fleet-gear stratum in the length frequency data table was assigned to the corresponding assessment fishery. Details on the fleet-gear length frequency strata making up each fishery are shown in **Appendix III. Table 6** shows the fisheries that are used for the assessment of yellowfin tuna. It shows also total catches by fishery accumulated for the entire catch data series (1950-2009) and the contribution that the catches from each fishery made to the total accumulated catches for 1950-2009, and in recent years (2005-09).

Fishery	Description	Total Catch 50-10 (,000 t)	% 50-10	% 06-10
FS	Industrial purse seines on free-swimming schools	1,905	19	17
LS	Industrial purse seines on associated schools (FAD)	1,435	14	17
LL	Industrial longlines (freezing vessels)	2,527	25	12
LF	Industrial longlines (fresh-tuna vessels)	877	9	8
BB	Pole-and-lines	444	4	5
GI	Gillnets	1,747	17	26
HD	Hand lines	653	6	8
TR	Troll lines	512	5	6
OT	Other artisanal gears	46	1	1

ii. *Allocation of assessment area*

The total catches, effort and samples existing for the yellowfin tuna were aggregated by assessment area. The areas that were used for the assessments of the yellowfin tuna stock are shown in **Table 7** and **Figure 2**. The catches of yellowfin tuna from areas outside the five areas were assigned to the closest area, as indicated through the arrows on **Figure 2**. **Table 7** shows also total catches by area accumulated for the entire catch data series (1950-2010) and the contribution that the catches from each area made out of the total accumulated catches for 1950-2010, and in recent years (2006-10).

Table 7: Areas used for the assessments of Indian Ocean yellowfin tuna; the total catches (tonnes) accumulated for the period 1950-2010 (Total Catch 50-10), the relative importance of the catches in each area over both the entire catch series (%50-10) and in current years (%06-10), and the accumulated 1950-2010 catches by assessment fishery (Table 6, page 9) in each of the areas concerned are also shown


Area	Description	Catch (,000 t) 50-10	% 50- 10	% 06- 10	Catches by Fishery 1950-2010 (,000 t)								
					FS	LS	LL	LF	BB	GI	HD	TR	OT
1	Arabian Sea	1,987	20	23	54	67	449	12	44	871	517	39	35
2	Western IO	4,662	46	42	1701	1202	1095	3	390	7	33	44	1
3	Southwest IO	874	9	10	102	128	345	2	2		47	271	1
4	Southern IO	106	1	0			103	8	<1	<1	<1	21	<1
5	Eastern IO	2,516	25	26	48	38	536	852	8	869	56	136	9

Note that the catches from areas that were kept for the assessment are shown in bold; the catches from other areas (*italics*) were allocated to one of the areas selected for the assessment, depending on the fishery (refer to Table 8 on page 8 for details)

The following process was used to allocate the existing samples by area:

- i. Allocation of catches, effort and length samples available for selected fisheries to specific assessment areas: The catches, effort and samples available for selected fisheries were fully assigned to specific assessment areas on the assumption that the majority of the specimens caught by those fisheries comes from the areas selected. This is thought to be the case with the majority of artisanal fisheries and with a limited number of industrial fisheries. Details on fleet-gear size frequency strata that make up each area are shown in **Appendix II**.
- ii. Allocation of the catches, effort and length samples available for other fisheries:
 - a. Breaking the nominal catches by time-period and area: the nominal catches for the fisheries concerned were assigned by month and 5° square areas by using the catches by time period and 5° square grids available for the yellowfin tuna (see “Standardizing the available catch-and-effort data” on page 3). The following process was used:
 - a. Nominal catches strata for which time-area catches exist:
 - i. Deleting strata from the CE table: The CE for NEI-(deep)-freezing longliners and NEI-fresh tuna longliners were not used because they refer to very specific areas and time-periods and are not considered to be representative of the activities of these fleets.
 - ii. Breaking the nominal catches by time and area: The nominal catches were broken by time and area in years for which spatio-temporal catches were available for the fleet concerned.

Figure 2: Areas used for the assessments of yellowfin tuna using MF-CL (note that Areas R3 and R4 have been modified for the present assessment, R4b representing the part of area R3 that has been moved to area R4 in the present assessment)



- b. Nominal catches strata for which CE data do not exist:
 - i. Catches per area are available for the same fleet in years before or after the year concerned:

1. Catches for the same species are available: The catches recorded in the five years closest to the year of reference were accumulated and the average values obtained used to break the catches per area in the year concerned. Data extending to up to 25 years above or below the year concerned are used.
2. Catches for other species are available:
 - a. The catches recorded in the year of reference were accumulated and the average values obtained used to break the catches per area in the year concerned
 - b. The catches recorded in the five years closest to the year of reference were accumulated and the average values obtained used to break the catches per area in the year concerned. Data extending to up to 25 years above or below the year concerned are used.
- ii. Catches per area are not available for the same fleet in years before or after the year concerned or they are available but very far in time (more than 25 years before or after the year concerned): The catches available for other fleets (and years) are used to break the nominal catches per month and 5° square area/s.
 1. Catches per area for the alternative fleet are available for the same year: This information is used to break the nominal catches per time and area.
 2. Catches per area for the alternative fleet are not available for the same year: The same substitution scheme as the one defined in ii.a. above is used.
- b. Estimation of total effort for industrial purse seine and longline fleets: The efforts recorded for industrial fleets were raised to represent the total catches of the fleet concerned, where required.
- c. Allocation of catches, effort and length samples recorded under areas that fall within a single assessment area: all catches, effort and length samples recorded under areas that fell within one of the areas used for the assessment (**Figure 2**) were assigned to the corresponding assessment area. It is important to note that all catch and effort data falls under this category (see a. above and “*Standardizing the available catch-and-effort data*” on page 3).
- d. Estimation of total effort for longline and purse seine fisheries: The effort of longline and purse seine fleets for which no effort data were available was estimated using the catches for those fleets and catch rates from fleets for which effort data were available. No effort was estimated in strata in which effort data were not available from any fleet.
- e. Allocation of length samples recorded under areas overlapping two or more assessment areas: all specimens from samples recorded under areas that overlapped two or more assessment areas were assigned by assessment area using the proportion that the catches in the area within each assessment area made out of the total catches concerned (e.g. if the samples overlapped areas 2 and 3 and the catches in the area sampled were 1t for area 2 and 2t for area 3, 1/3 of the fish sampled by length class was assigned to area 2 and 2/3 to area 3).

The resulting data were aggregated to obtain total catches (in weight), and effort and length samples by year, quarter, assessment fishery, and assessment area.

It is important to note that no weighting procedure was used when aggregating the available length samples by assessment fishery and area.

Putting together the input file for the assessments of yellowfin tuna with MF-CL

The catches (in number and weight), effort and length frequency samples from some fisheries in some areas were reallocated to other areas, especially in cases where the catches represented a small proportion of the total catches for the fishery concerned and where no tag releases had been recorded for those fisheries and areas. Overall, around 3% of the catches of yellowfin tuna were re-allocated to other areas. The catches in italics in **Table 7** (page 6) represent the catches that were re-allocated to other areas for each fishery. The criteria used for the allocation of catches are shown on **Table 8**.

Table 8: Areas used for the assessments of the Indian Ocean yellowfin tuna stock: criteria used for the re-allocation of catches, effort and length samples from the areas where they were recorded to the areas used for the MF-CL assessment. The proportion of catches from each fishery that were re-allocated to other area/s, over the total catches for that same fishery for the entire time series (1950–2009), is also shown

Fishery	Areas used (MFCL assessment)	Area 1	Area 2	Area 3	Area 4	Area 5	% of catches re- allocated
		Catch, effort, length frequency data re-allocated to Area					
FS	2, 3, 5	2	2	3		5	2.8
LS	2, 3, 5	2	2	3		5	4.7
LL	1, 2, 3, 4, 5	1	2	3	4	5	0.0
LF	5	5	5	5	5	5	2.9
BB	2	2	2	2	2	2	12.2
GI	1, 5	1	1		5	5	0.4
HD	1	1	1	1	1	1	20.8

Fishery	Areas used (MFCL assessment)	Area 1	Area 2	Area 3	Area 4	Area 5	% of catches re- allocated
		Catch, effort, length frequency data re-allocated to Area					
TR	2, 3, 5	2	2	3	5	5	11.7
OT	1, 5	1	1	1	5	5	4.0

After the re-allocation process, the catches (in weight), effort and length frequency samples were aggregated by Fishery, Assessment Area, year, and quarter. An example of the Input Table that was built for the assessments of yellowfin tuna with MF-CL can be found in **Appendix I**.

Assigning releases by year, quarter and assessment area (i.e. release event)

In 2011, the Secretariat included in the database all the tagging data generated during the pilot and small-scale projects implemented during the Indian Ocean Tuna Tagging Programme. This includes releases in Maldives, India, Indonesia, Mayotte as well as releases in the eastern and south west Indian Ocean, made by other institutions with the support of IOTC. During the preparation of the tagging data, all the releases and recoveries from these projects have also been included.

Year and quarter of release are derived from the exact date of release. All the recoveries are then assigned in each of the assessment area described in figure 2. Yellowfin were released during 10 quarters from May 2005 to September 2007 in region 1 (3169 releases), 2 (56521 releases), 3 (707 releases), 4 (2 releases) and 5 (2182 releases).

Release Area	Release Year	Release Quarter	Nb of YFT released	
1	2005	4	29	
		5	5	
	2007	1	2056	
		2	1079	
2	2003	4	48	
		2004	2	13
			3	1264
	2005	4	8	
		2006	1	91
			2	25
			3	1409
	2007	4	3529	
		2008	1	3520
			2	13409
			3	12164
	2009	4	7364	
		2010	1	92
			2	1727
			3	7822
	2011	4	172	
1		618		
3	2002	4	1577	
		1	1385	
3	2004	2	284	
		1	15	
	2005	2	15	
		1	9	
2006	2	77		
	3	556		
4	2007	2	2	
		3	2	

		4	1
	2006	1	4
	2008	1	2
		2	26
4	2005	1	1
		3	1
5	2003	4	252
	2004	3	10
		4	359
	2005	1	704
		2	35
	2006	2	24
		3	13
		4	1
	2007	3	460
	2008	1	324

Assigning recaptures by release event, year and quarter of recapture, fishery and assessment area

i. Assigning date and location to the recoveries:

- i. Linking date of recovery (i.e. unloading) and well number with the logbook database: recoveries made during fishing can be assigned with a precise date and location of recapture. However, a large proportion of the recoveries were made on-board the purse-seine fleet at the moment of the unloading. For these, the information provided is the date of recovery (i.e. unloading) and the well, and a link has to be made with the logbook of the purse-seiner in order to be able to assign a date and position of recapture.
 - i. Assigning possible sets from which the recapture might come from: the editing of the data is done for each tag one by one through a “data editor” software that was designed to link the recovery database with the logbook database. The software extract from the logbook the data for each set that was loaded into the well where the fish was recovered.
 - ii. Assigning possible sets from which the recapture might come from: in the case of only one set was loaded in the said well, the date and position of the recovery will be the one assigned for this set. However, most of the time, sets were mixed and therefore the date and position will be an average of the different dates and positions.

Putting together release and recovery data of yellowfin tuna

Different files are produced for the users of the tagging data, however, the main client file includes **all** the individual information for the releases and the recoveries, including the estimation of the position for the recoveries on-board the purse-seine fleet.

Reporting Rates, Shedding Rates and Reference Catch

In addition to the tagging data (releases and recoveries), it is important that the users are provided with additional sets of data including Reporting Rates, Shedding Rates and the Reference Catch:

Reporting Rates: out of the recapture tagged not all are reported to the IOTC Secretariat. In order to fully account for the tagged fish that have been recaptured and not reported, it is necessary to assess the Reporting Rates. Estimating reporting rates was done by IOTC for the European purse-seine fleet based in Seychelles through on-board observer and the cooperation of skippers.

Shedding Rates: shedding rates accounts for the number of tags that fell of the fish after tagging and are assessed through double tagging. A shedding rate analysis was done in 2008 and updated in 2009.

Reference Catch: reporting rates estimated for the purse-seine fleet do not apply to the totality of the catch unloaded by the fleet, but only to the proportion of the catch that is unloaded in Seychelles. This proportion were assessed by the scientists of the EU in order to be incorporated in the final analysis of the tagging data.

- v. **Length samples referring to areas that overlap to or more assessment areas:** The majority of the samples available for the longline fishery are recorded by quarter and 10 degrees latitude by 20 degrees longitude areas⁷. This means that a significant amount of samples overlaps two or more assessment areas, samples that need to be allocated by assessment area.

In consequence, the length distributions that are derived for the above fisheries and periods are not considered to be representative of those fisheries as a whole.

iv. **Quality of release data**

There is no major problem identified in the quality of the release data from the RTTP-IO. Reliability codes were used to qualifying the data quality, *e.g.* quality of the tagging or of the tagged fish. For instance the values attained by some of these codes are as follows:

- i. Some doubt on the species identified for only 1.04% of all tuna tagged; among these fish there are 579 Unknown species (unknown) accounting for 0.34%;
- ii. 1.14% of the Fork Length are not completely reliable including 1,368 with no FL accounting for 0.81%;
- iii. 2.43% of the fish are not released in a very “good” condition, *e.g.* 1,052 dropped on deck or having hit the vessel accounting for 0.63%.

There is probably more quality issues with the release data from the pilot and small-scale operations due to the nature of these projects. However, this data has not yet been fully analysed.

v. **Quality of recovery data**

The quality of the recovery data is completely dependent on the recovery platform and one the awareness campaign that were designed and implemented. In fact, recoveries of artisanal fleet for example, are in general of a lower quality. By being host in Seychelles, based of the major tuna purse-seine fleet of the Indian Ocean, the RTTP-IO was able to design and implement a large and comprehensive tag recovery strategy in Seychelles, ensuring good quality of the recoveries. However, as the majority of the recoveries were made by the stevedores during the unloading and the transshipment of the fishing vessel, part of the recovery information had to be estimated from the logbooks, *i.e.* data and position of the recapture, type of association of the school (*e.g.* free or log school).

In the 10560 recoveries of yellowfin tuna, there are 891 for which no position of recoveries could be assigned (542 from purse-seine). One of the main issues regarding the quality of the recoveries of the purse-seine fleet is that for a large proportion of them, the type of school association is unknown, while they are defined separated fisheries in the model, *i.e.* for 34% of the recoveries, the association type is not known.

<i>Fishing mode</i>	<i>Nb</i>
Free-swimming schools	1253
Associated schools	4840
Unknown fishing mode	3144

To include these recoveries in the model, they had to be distributed among the free school and log school recoveries, keep the proportion of each type for each year.

Main changes in the preparation of data with respect to previous MF-CL assessments

There have not been important changes in the preparation of data for the assessments of yellowfin tuna, other than:

- i.
- ii. Breaking of longline fisheries, recorded as a single fishery in previous assessments, into freezing longline and fresh-tuna longline fisheries, with all catch, effort and length data for fresh-tuna longline fisheries assigned to Area 5. The reason for this is that fresh-tuna longliners have traditionally operated in tropical waters of the eastern Indian Ocean, the main fleets operating Indonesia or Taiwan, China flags. The separation of longline fisheries into freezing longline (LL) and fresh-tuna longline (LF) also allowed for a more precise estimation of total effort for the freezing longline component, as effort data are more complete in this case.
- iii. **Correction in the recovery data:** in order to better estimate the real number of tag recapture, the recoveries from the purse-seiner where separated in at-sea recoveries (recoveries made during fishing) and on-land recoveries (recoveries made during the unloading/transshipment in Seychelles), and for each the number of recoveries with an unknown set type were distributed in the free and log school type. Moreover, the recoveries on-land were corrected by the reporting rate estimated through the tag seeding experiment that took place on-board the purse-seine fleet based in Seychelles and with the proportion of the catch of the purse-seine fleet that was unloaded in Seychelles.

⁷ Note that the standards of reporting existing at the IOTC for size data from longline fisheries are month and 5 degrees square grid

Proposal for additional work to be undertaken in relation with the preparation of datasets for future assessments

The fisheries defined for the MF-CL assessment are made in some cases of a high number of fleets and gears (Appendix III). The catches and samples recorded under some fisheries may, for this reason, combine specimens of different size, depending on the fleet or time-period involved and the amount of data that are available over the time series from each fleet. In addition, the representativeness of some sets of catch-and-effort and/or size data may be compromised due to poor sampling coverage or for other reasons. A more thorough analysis of the information at hand is required in order to be able to select the fisheries and areas that represent best the yellowfin tuna fishery and the fleets that made up each fishery and area.

The following changes are suggested in the preparation of data for future assessments of yellowfin tuna with MF-CL, provided that the above analysis is undertaken:

- i. Weighting all length frequency distributions by the actual sample numbers: The length frequency distributions for purse seiners represent the total number of specimens within the sampling units selected while the length frequency distributions for other fleets represent the actual sample numbers; initially the length frequency distributions for purse seiners could be weighted by the real sample numbers in each stratum. However, the actual number of lengths measured on industrial purse seiners is not known for all fleets and time-area strata.
- ii. Weighting the existing length samples by fleet prior to the aggregation of samples under each fishery: The length distributions in the MF-CL were built by aggregating the length samples for the fleets under each fishery, irrespective of the sampling coverage levels for the fleets involved; in some cases, the estimation of length distributions could be improved by weighting the lengths sampled by fleet prior to the aggregation of lengths for the fleets making up a fishery. However, the little amount of size data that are available from some fisheries, in particular coastal fisheries, may compromise the weighting of samples for those and needs to be further explored.
- iii. Improving the estimation of the total number of recaptured tags: the process on how to distribute the unknown set type recoveries and how to adjust for the reporting rate need to be better discussed and formalized for all the analysis.

APPENDIX I

MF-CL Input Table

Year	The year where the catches were made
Quarter	The quarter where the catches were made (Jan-Mar(1), Apr-Jun(2), Jul-Sep(3), Oct-Dec(4))
FisheryCode	Type of fishery (FS; LS; LL; LF; GI; HD; TR; OT)
Assessment Area	The areas used for the assessment (1; 2; 3; 4; 5)
Catch	Total catch estimated (in metric tonnes of fish)
Effort	Fishing effort estimated (longline and purse seine fisheries) or available (other fisheries)
C001...C150	Number of yellowfin tuna measured (sampled) by length class, using 10cm as the first class(lower bound) and 2cm intervals
C001	Number of YFT specimens having a fork length between 10cm and 12cm
C002	Number of YFT specimens having a fork length between 12cm and 14cm
	And so forth

APPENDIX II

Fleets whose catches, effort and length samples were fully allocated to specific Assessment Areas

Gear	Fleet Code	Fishery	Ass. Area	Real Area
BB	AUS	BB	2	4
BBPS	AUS	BB	2	4
BB	EGY	BB	2	1
BB	ESP	BB	2	2
BB	IDN	BB	2	5
BB	IND	BB	2	99
BB	KOR	BB	2	3
BB	LKA	BB	2	5
BB	MDG	BB	2	3
BB	MDV	BB	2	2
BBM	MDV	BB	2	2
BBN	MDV	BB	2	2
BB	TZA	BB	2	2
GILL	ARE	GI	1	1
GILL	AUS	GI	5	4
GILL	BGD	GI	5	5
GILL	BHR	GI	1	1
GILL	DJI	GI	1	1
GILL	EGY	GI	1	1
GILL	ERI	GI	1	1
GILL	IDN	GI	5	5
GILL	IND	GI	99 ⁸	99
GILL	IRN	GI	1	1
GILL	ISR	GI	1	1
GILL	JOR	GI	1	1
GILL	KEN	GI	1	2
GILL	KWT	GI	1	1
G/L	LKA	GI	5	5
GILL	LKA	GI	5	5
GIOF	LKA	GI	5	5
GILL	MYS	GI	5	5
GIHA	OMN	GI	1	1
GILL	OMN	GI	1	1
GILL	PAK	GI	1	1
GILL	QAT	GI	1	1
GILL	SAU	GI	1	1
GILL	SDN	GI	1	1
GILL	SYC	GI	1	2
GILL	THA	GI	5	5
GILL	TMP	GI	5	5
GILL	TWN	GI	5	4
GILL	TZA	GI	1	2
GILL	YEM	GI	1	1
HAND	ARE	HD	1	1
HAND	AUS	HD	1	4
HAND	BGD	HD	1	5
HAND	BHR	HD	1	1
HAND	COM	HD	1	3
HAND	EGY	HD	1	1
LL	EGY	HD	1	1
HAND	FRAT	HD	1	3
HAND	GBRT	HD	1	2
HAND	IDN	HD	1	5
LLCO	IDN	HD	1	5
LLF	IDN	HD	1	5
HAND	IND	HD	1	99
HAND	IRN	HD	1	1
HAND	KEN	HD	1	2
HAND	LKA	HD	1	5
LL	LKA	HD	1	5
LLHA	LKA	HD	1	5
HAND	MDV	HD	1	2

Gear	Fleet Code	Fishery	Ass. Area	Real Area
LLCO	MDV	HD	1	2
HAND	MYS	HD	1	5
LLCO	MYS	HD	1	5
HAND	OMN	HD	1	1
HAND	SAU	HD	1	1
LLCO	SAU	HD	1	1
HAND	SYC	HD	1	2
HAND	TMP	HD	1	5
HAND	TZA	HD	1	2
GIHT	YEM	HD	1	1
HAND	YEM	HD	1	1
HATR	YEM	HD	1	1
HAND	ZAF	HD	1	3
FLL	BLZ	LF	5	4
FLL	CHN	LF	5	5
FLL	IDN	LF	5	5
FLL	IND	LF	5	5
FLL	LKA	LF	5	5
FLL	MDV	LF	5	2
FLL	MYS	LF	5	5
FLL	NEI-ICE	LF	5	5
FLL	NEI-IDN	LF	5	5
FLL	OMN	LF	5	1
FLL	THA	LF	5	5
FLL	TWN	LF	5	0
FLL	VUT	LF	5	4
UNCL	ARE	OT	1	1
HATR	AUS	OT	5	4
HOOK	AUS	OT	5	4
SEN	AUS	OT	5	4
SPOR	AUS	OT	5	4
TRAP	AUS	OT	5	4
TRAW	AUS	OT	5	4
UNCL	AUS	OT	5	4
TRAW	BGR	OT	1	2
UNCL	BGR	OT	1	2
HATR	BHR	OT	1	1
HOOK	BHR	OT	1	1
TRAW	BHR	OT	1	1
TRAW	BLZ	OT	1	2
UNCL	COM	OT	1	3
PS	EGY	OT	1	1
TRAW	EGY	OT	1	1
UNCL	EGY	OT	1	1
TRAW	ERI	OT	1	1
HATR	FRAT	OT	1	3
SPOR	GBRT	OT	1	2
PS	IDN	OT	5	5
BS	IDN	OT	5	5
DSEI	IDN	OT	5	5
FN	IDN	OT	5	5
LIFT	IDN	OT	5	5
LIGB	IDN	OT	5	5
PSS	IDN	OT	5	5
UNCL	IDN	OT	5	5
LIFT	IND	OT	99	99
HATR	IND	OT	99	99
HOOK	IND	OT	99	99
PSS	IND	OT	99	99
TRAW	IND	OT	99	99
UNCL	IND	OT	99	99
UNCL	ISR	OT	1	1
OTHER	JOR	OT	1	1
UNCL	JOR	OT	1	1
UNCL	JPN	OT	1	2
UNCL	KEN	OT	1	2
BS	LKA	OT	5	5
HARP	LKA	OT	5	5

Gear	Fleet Code	Fishery	Ass. Area	Real Area
HATR	LKA	OT	5	5
RIN	LKA	OT	5	5
UNCL	LKA	OT	5	5
FN	MDV	OT	1	2
UNCL	MDV	OT	1	2
HOOK	MYS	OT	5	5
PSS	MYS	OT	5	5
TRAP	MYS	OT	5	5
TRAW	MYS	OT	5	5
UNCL	MYS	OT	5	5
BS	OMN	OT	1	1
CN	OMN	OT	1	1
FN	OMN	OT	1	1
TRAP	OMN	OT	1	1
UNCL	OMN	OT	1	1
PSS	SAU	OT	1	1
TRAP	SAU	OT	1	1
TRAW	SAU	OT	1	1
UNCL	SAU	OT	1	1
UNCL	SDN	OT	1	1
PSS	SUN	OT	1	1
UNCL	SUN	OT	1	2
UNCL	SYC	OT	1	2
PSS	THA	OT	5	5
TRAW	THA	OT	5	5
UNCL	THA	OT	5	5
HATR	TMP	OT	5	5
UNCL	TMP	OT	5	5
OTHER	TZA	OT	1	2
PSS	TZA	OT	1	2
UNCL	TZA	OT	1	2
UNCL	YEM	OT	1	1
HOOK	ZAF	OT	1	3
SPOR	ZAF	OT	1	3
UNCL	ZAF	OT	1	3
TROL	ARE	TR	2	1
TROL	AUS	TR	5	4
TROL	BHR	TR	2	1
TROL	COM	TR	3	3
TROL	EGY	TR	2	1
TROL	FRA-REU	TR	3	4
TROL	FRAT	TR	3	3
TROL	IDN	TR	5	5
TROL	IND	TR	5	99
TROL	IRN	TR	2	1
TROL	ISR	TR	2	1
TROL	JOR	TR	2	1
TROL	KEN	TR	2	2
TROL	LKA	TR	5	5
TROL	MDG	TR	3	3
TROL	MDV	TR	2	2
TROLM	MDV	TR	2	2
TROLN	MDV	TR	2	2
TROL	MUS	TR	3	4
TROL	MYS	TR	5	5
TROL	OMN	TR	2	1
TROL	SAU	TR	2	1
TROL	SYC	TR	2	2
TROL	TMP	TR	5	5
TROL	TZA	TR	2	2
TROL	YEM	TR	2	1

⁸ Area 99: Catches, effort and length samples in the Western Indian Ocean assigned to Area 1 and those in the Eastern Indian Ocean assigned to Area 5

APPENDIX III

Fleets making up the Fisheries selected for the assessment

Purse seine (PS)

Fishery	Fleet	Gear
PS	AUS	PS
PS	BGR	PS
PS	BLZ	PS
PS	ESP	PS
PS	FRA	PS
PS	FRAT	PS
PS	IRN	PS
PS	JPN	PS
PS	MUS	PS
PS	MYS	PS
PS	NEI-OTH	PS
PS	NEI-SUN	PS
PS	SUN	PS
PS	SYC	PS
PS	THA	PS

Longline (LL)

Fishery	Fleet	Gear
LL	AUS	ELL
LL	BLZ	LL
LL	CHN	LL
LL	ESP	ELL
LL	FRA-REU	ELL
LL	FRAT	ELL
LL	GBR	ELL
LL	GIN	ELL
LL	IDN	LL
LL	IND	LL
LL	IND	LLEX
LL	IRN	LL
LL	JPN	LL
LL	KEN	ELL
LL	KOR	LL
LL	MDG	ELL
LL	MDG	LL
LL	MUS	ELL
LL	MUS	LL
LL	NEI-DFRZ	ELL
LL	NEI-DFRZ	LL
LL	NEI-DFRZ	TLL
LL	OMN	LL
LL	PAK	LL
LL	PHL	LL
LL	PRT	ELL
LL	PRT	LL
LL	PRT	LID
LL	PRT	SLL
LL	SEN	ELL
LL	SUN	LL
LL	SYC	ELL
LL	SYC	LL
LL	THA	LL
LL	TWN	LL
LL	TZA	ELL
LL	URY	ELL
LL	VUT	LL
LL	ZAF	LL
LL	ZAF	SLL
LL	ZAF	TLL

Longline fresh (FL)

Fishery	Fleet	Gear
LF	BLZ	FLL
LF	CHN	FLL
LF	IDN	FLL
LF	IND	FLL
LF	LKA	FLL
LF	MDV	FLL
LF	MYS	FLL
LF	NEI-ICE	FLL
LF	NEI-IDN	FLL
LF	OMN	FLL
LF	THA	FLL
LF	TWN	FLL
LF	VUT	FLL

Pole-and-line (BB)

Fishery	Fleet	Gear
BB	AUS	BB
BB	AUS	BBPS
BB	EGY	BB
BB	ESP	BB
BB	IDN	BB
BB	IND	BB
BB	KOR	BB
BB	LKA	BB
BB	MDG	BB
BB	MDV	BB
BB	MDV	BBM
BB	MDV	BBN
BB	TZA	BB

Gillnet (GI)

Fishery	Fleet	Gear
GI	ARE	GILL
GI	AUS	GILL
GI	BGD	GILL
GI	BHR	GILL
GI	DJI	GILL
GI	EGY	GILL
GI	ERI	GILL
GI	IDN	GILL
GI	IND	GILL
GI	IRN	GILL
GI	ISR	GILL
GI	JOR	GILL
GI	KEN	GILL
GI	KWT	GILL
GI	LKA	G/L
GI	LKA	GILL
GI	LKA	GIOF
GI	MYS	GILL
GI	OMN	GIHA
GI	OMN	GILL
GI	PAK	GILL
GI	QAT	GILL
GI	SAU	GILL
GI	SDN	GILL
GI	SYC	GILL
GI	THA	GILL
GI	TMP	GILL
GI	TWN	GILL
GI	TZA	GILL
GI	YEM	GILL

Handline (HD)

Fishery	Fleet	Gear
HD	ARE	HAND
HD	AUS	HAND
HD	BGD	HAND
HD	BHR	HAND
HD	COM	HAND
HD	EGY	HAND
HD	EGY	LL
HD	FRAT	HAND
HD	GBRT	HAND
HD	IDN	HAND
HD	IDN	LLCO
HD	IDN	LLF
HD	IND	HAND
HD	IRN	HAND
HD	KEN	HAND
HD	LKA	HAND
HD	LKA	LL
HD	LKA	LLHA
HD	MDV	HAND
HD	MDV	LLCO
HD	MYS	HAND
HD	MYS	LLCO
HD	OMN	HAND
HD	SAU	HAND
HD	SAU	LLCO
HD	SYC	HAND
HD	TMP	HAND
HD	TZA	HAND
HD	YEM	GIHT

Fishery	Fleet	Gear
HD	YEM	HAND
HD	ZAF	HAND

Trolling (TR)

Fishery	Fleet	Gear
TR	ARE	TROL
TR	AUS	TROL
TR	BHR	TROL
TR	COM	TROL
TR	EGY	TROL
TR	FRA-REU	TROL
TR	FRAT	TROL
TR	IDN	TROL
TR	IND	TROL
TR	IRN	TROL
TR	ISR	TROL
TR	JOR	TROL
TR	KEN	TROL
TR	LKA	TROL
TR	MDG	TROL
TR	MDV	TROL
TR	MDV	TROLM
TR	MDV	TROLN
TR	MUS	TROL
TR	MYS	TROL
TR	OMN	TROL
TR	SAU	TROL
TR	SYC	TROL
TR	TMP	TROL
TR	TZA	TROL
TR	YEM	TROL

Other gears (OT)

Fishery	Fleet	Gear
OT	ARE	UNCL
OT	AUS	HATR
OT	AUS	HOOK
OT	AUS	SEN
OT	AUS	SPOR
OT	AUS	TRAP
OT	AUS	TRAW
OT	AUS	UNCL
OT	BGR	TRAW
OT	BGR	UNCL
OT	BHR	HATR
OT	BHR	HOOK
OT	BHR	TRAW
OT	BLZ	TRAW
OT	COM	UNCL
OT	EGY	PS
OT	EGY	TRAW
OT	EGY	UNCL
OT	ERI	TRAW
OT	FRAT	HATR
OT	GBRT	SPOR
OT	IDN	BS
OT	IDN	DSEI
OT	IDN	FN
OT	IDN	LIFT
OT	IDN	LIGB
OT	IDN	PSS
OT	IDN	UNCL
OT	IND	LIFT
OT	IND	HATR
OT	IND	HOOK
OT	IND	PSS
OT	IND	TRAW
OT	IND	UNCL
OT	ISR	UNCL
OT	JOR	OTHER
OT	JOR	UNCL
OT	JPN	UNCL
OT	KEN	UNCL
OT	LKA	BS
OT	LKA	HARP
OT	LKA	HATR
OT	LKA	RIN
OT	LKA	UNCL
OT	MDV	FN

Fishery	Fleet	Gear
OT	MDV	UNCL
OT	MYS	HOOK
OT	MYS	PSS
OT	MYS	TRAP
OT	MYS	TRAW
OT	MYS	UNCL
OT	OMN	BS
OT	OMN	CN
OT	OMN	FN
OT	OMN	TRAP
OT	OMN	UNCL
OT	SAU	PSS
OT	SAU	TRAP
OT	SAU	TRAW
OT	SAU	UNCL
OT	SDN	UNCL
OT	SUN	PSS
OT	SUN	UNCL
OT	SYC	SUPP
OT	SYC	UNCL
OT	THA	PSS
OT	THA	TRAW
OT	THA	UNCL
OT	TMP	HATR
OT	TMP	UNCL
OT	TZA	OTHER
OT	TZA	PSS
OT	TZA	UNCL
OT	YEM	UNCL
OT	ZAF	HOOK
OT	ZAF	SPOR
OT	ZAF	UNCL

