

Sampling strategies as a function of the IOTC fishery

Category: Fishing Gear

[IOTC ROS SFO TR16]

Sampling strategies as a function of the IOTC fishery

Descriptor

This module aims to familiarize Observers with sampling requirements, procedures and methods to be used in the collection of scientific fisheries data with IOTC fisheries (industrial tuna purse-seine; pelagic longline; pelagic gillnet; pole-and-line) as these will be used daily in their routine work.

Category: Fishing Gear

[IOTC ROS SFO TR16]

IOTC fisheries: Pelagic gillnet fishery

Trainee performance is evaluated against the following agreed IOTC ROS competency standards:

- Capable of collecting and estimating catch weight, volumes and ratios according to ROS standard procedures.

The achieving of these standards is demonstrated by candidate capacity to:

- Correctly select sampling strategies to use as a function of the fisheries, vessel configuration and set type (for the purse-seine fisheries);
- Correctly estimate weights, volumes and ratios with the industrial tuna purse-seiners, pelagic longliners, pole and line vessels and gillnetters.

Sampling strategies

- Sampling objectives include:
 - CPUE recording catch and effort
 - catch composition (target species and by-catch)
 - biological biometrics (*length frequencies, sex and maturity, diet and collecting DNA material*)

Sampling strategies

- Fishing operation and gear information can be obtained from Fishing Masters Logbook
- Observer's responsibility to verify information through
 - observations
 - measuring gear and recording operations
 - sampling

Sampling strategies

- CPUE Data (FORM 3– GIL)
- Gear information for gillnets are
 - net depth
 - length of the net set;
 - soak time (time the net is in the water
 - set depth (vertical position in the water column)
 - mesh size

Sampling strategies

- Recording Catch composition
- catch composition can be obtained two sources
 1. vessels logbook (fish retained)
 2. observer monitoring (fish retained and by-catch discarded)
- Observe position requires view of net coming out of water and onto the deck*
- Two possibilities
 1. Total observed catch – *(monitoring whole net [100%] hauled back)*
 2. Percentage of net hauled *(length of net observed vs total length let set – raised figures)*

Sampling strategies

- Recording Catch composition
- Percentage of net hauled equals

•*(length of net observed vs total length let set)*

Number of panels observed / total length of net (m)

Sampling strategies

- Example:** Percentage of net observed

- net panel dimensions 30 m long and 24 m deep
- number of panels set 250
- number of panels observed 150

- Calculation *Total length of net set* = (30×250) = 7500 m

- *Length observed hauled* = (30×150) = 4500 m

Number of panels observed / total length of net (m)

Sampling strategies

- Example:** Calculated total catch raised by percentage observed:

number of each species observed divided by % of net observed

Species Observed	YFT	SKJ	KAW	LOT	GUT	SFA
No. Obs	20	37	145	200	89	23
% of total	0.6	0.6	0.6	0.6	0.6	0.6
Calculated total catch per species	33	62	242	333	148	38

Sampling strategies

- Length measurements
- factors influencing size range of the catches in gillnets
 - net selectivity (mesh size and net tension (hanging ratio))
 - shoaling fish
 - variable catch rates

Sampling strategies

- Length measurements
- Sampling position and sample selection
- Large industrial vessels

factory deck where fish are processed for freezing. sampling station close to where the fish come into the factory

- Smaller open deck vessels

Where fish temporarily held before transfer to hold (*This could be in bins or baskets or sections of the deck partitioned off with planks*)

Sampling strategies

- Length measurements
- Stratified Sampling Protocol *allows for the varied catch rates*
- Measure fixed number of target species in different size categories to get a representative sample for length
- Practical stratified sampling
- Select a number of fish of same species BUT randomly for length
- Measure until a clear length mode is achieved (up to 100 fish)
- If this is not evident on either of the extreme sizes then a higher number of fish should be measured.*

Sampling strategies

- Length measurements
- For Example:*
- 100 Kawakawa and 100 Longtail tuna
- The length frequency shows a narrow length mode*
- Notably this would indicate a highly selective net with a high hanging ratio.*

Length (cm) (FL)	KAW	LOT
33	0	
34	0	
35	0	
36	2	
37	12	0
38	38	1
39	41	2
40	3	12
41	4	38
42	0	44
43	0	3
44		0
45		
46		
Total measured	100	100

Sampling strategies

- Length and Catch Composition (mixed species and proportional sampling)
 - sample a fixed number (or volume) at intervals
 - samples must be collected before its sorted if this is possible
- Sort the sample to relative species level and then record length frequencies per species
- The number of species and number per species sampled should provide indication of
 - the relative catch composition (relative number per species)
 - the relative length frequency of the different species.

END - *IOTC SFO TR 16*

THANK YOU

[Insert here logos from partners other than CCP]